

λ Boo stars among the γ Dor-type pulsators: the cases of HD 218427 and HD 239276

E. Rodríguez,¹ J. C. Suárez,^{1,2} A. Moya,^{1,2} M. A. Dupret,² A. Grigahcène,³ V. Costa,¹ M. J. López-González,¹ A.-Y. Zhou,⁴ P. J. Amado,¹ E. Poretti,⁵ J.-Y. Wei,⁴ Y. Fan⁴

¹ Instituto de Astrofísica de Andalucía, CSIC, P.O. Box 3004, E-18080 Granada, Spain, E-mail: eloy@iaa.es

² LESIA, Observatoire de Paris-Meudon, UMR 8109, 92190 Meudon, France

³ CRAAG, Algiers Observatory, BP 63 Bouzareah 16340, Algiers, Algeria

⁴ National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

⁵ INAF-Osservatorio Astronomico di Brera, Via E. Bianchi 46, 23807 Merate, Italy

The γ Dor-type variables constitute a relatively recently recognized class of pulsating variables in the zone where the red edge of the δ Sct region intersects with the main sequence. λ Boo stars are metal-poor Population I stars that show significant underabundances of metals, except for the elements C, N, O and S. These stars are also characterized by showing broad, but often shallow, hydrogen-line wings and weak MgII λ 4481 lines. A number of λ Boo stars are known to be δ Sct pulsators, but this question is still open concerning the γ Dor-type pulsators. To date, HR 8799 is the unique case known of a γ Dor-type variable being a λ Boo star too. However, some of these variables seem to be metal-deficient. This has important implications in asteroseismology of γ Dor stars concerning the distinction between the two possibilities or, vice versa, if a star is already known to be of the λ Boo-type, this can be used to constrain asteroseismic models.

In this work, we study the cases of HD 218427 and HD 239276. Both variables were discovered as multiperiodic γ Dor-type pulsators (Rodríguez et al. 2006a,b), by means of simultaneous $uvby\beta$ photometry, while they were used as check stars for observations devoted to other already well-known pulsators in the Lower Instability Strip, AC And and XX Cyg.

HD 218427 and HD 239276 present very similar photometric characteristics to the multiperiodic γ Dor HR 8799, including a slight deficiency in metal content. This could be a sign of a λ Boo nature as was already found for HR 8799 by Gray & Kaye (1999). Indeed, the three stars are located inside the λ Boo region of both $(m_1, b - y)$ and $([m_1], \beta)$ diagrams (Gray 1988, Gray & Corbally 1993).

The Time-Dependent Convection (TDC) treatment for multicolour photometry (Dupret et al. 2005, Grigahcène et al. 2005) and the Frequency Ratio Method (FRM) are used to discriminate the angular orders ℓ of the three main modes excited in these two stars. However, no definitive conclusions are obtained concerning the true nature of the observed metal deficiency, such as: (a) they really are metal-deficient stars or (b) they are λ Boo stars.

In the case of HD 239276, by means of the TDC study, the two main modes are identified as $\ell = 1$ and the third mode is suggested to be $\ell = 1$ or 2. However, our results do not allow us to discriminate between a λ Boo or a truly metal-poor nature for this star. On the other hand, the FRM suggests low metallicity for this star, but a λ Boo nature cannot be ruled out.

References

Dupret M.-A., Rodríguez E., Garrido R., et al., 2005, MNRAS 360, 1143
 Gray R. O., 1988, AJ, 95, 220
 Gray R. O., Corbally C. J., 1993, AJ, 106, 632
 Gray R. O., Kaye A. B., 1999, AJ, 118, 2993
 Grigahcène A., Dupret M.-A., Gabriel M., Garrido R., Scuflaire R., 2005, A&A, 434, 1055
 Moya A., Suárez J. C., Amado P. J., Martín-Ruiz S., Garrido R., 2005, A&A, 432, 189
 Rodríguez E., Amado P. J., Suárez J. C., et al., 2006a, A&A, 450, 715
 Rodríguez E., Costa V., Zhou A.-Y., et al., 2006b, A&A, 456, 261




Figure 1: Strömgren photometric amplitude ratios obtained for HD 239276 with the TDC treatment for a model with $M=1.3 M_{\odot}$, $Z=0.01$, $\log T_e=3.8569$, $\alpha=2$.



Juan Carlos Suárez, Pedro Amado, Rafa Garrido and Eric Michel.