

Is the western population of the European mink, (*Mustela lutreola*), a distinct Management Unit for conservation?

J.R. Michaux^{a,*}, R. Libois^a, A. Davison^c, P. Chevret^b, R. Rosoux^d

^aUnité de Recherches Zoogéographiques, Institut de Zoologie, Quai Van Beneden, 22, 4020 Liège, Belgium

^bLaboratoire de Paléontologie—cc064, Institut des Sciences de l'Evolution (UMR 5554-CNRS), Université Montpellier II, Place E. Bataillon, 34095 Montpellier Cédex 05, France

^cDepartment of Ecology and Evolutionary Biology, Biological Institute, Graduate School of Science, Tohoku University, Aramaki-Aza-Aoba, Aoba-ku, Sendai 980 8578, Japan and ICAPB, Ashworth Laboratories, West Mains Road, University of Edinburgh, Edinburgh EH9 3JT, UK

^dMuseum des Sciences naturelles, rue Marcel Proust, 6 F 45000 Orléans, France

Received 20 November 2002; received in revised form 6 March 2003; accepted 6 March 2003

Abstract

The European mink (*Mustela lutreola*) is one of the most threatened carnivores in Europe, with fragmented populations in Belarus, Russia and Romania, as well in south-western France and northern Spain. Many populations have become extinct recently, or are declining. We investigated mitochondrial DNA variation, using the complete D-loop region, and concentrating on the west European population. The aim was two-fold: to use the genetic information to advise on the conservation of European mink, and to begin to understand their history through the Pleistocene. Captive breeding and re-introduction programmes are underway, so it is particularly vital to know whether the West European population should be treated separately. We find that European mink probably colonised from a single refugium after the last glaciation. West European populations may be fixed for a single haplotype, also suggesting a common origin. Despite this evidence for gene flow, following the precautionary principle we suggest that mink from the three geographically separate populations (Romania, Eastern and Western Europe) should be managed separately, for the moment.

© 2003 Elsevier Ltd. All rights reserved.

Keywords: *Mustela lutreola*; mtDNA; Conservation biology; Management unit; Phylogeography

1. Introduction

The European mink (*Mustela lutreola*) is one of the most threatened carnivores (Baillie and Groombridge, 1996). In the latter part of the 20th century, its distribution has fragmented, and populations continue to decline (Van Bree and Saint Girons, 1966; Camby, 1990; Maran, 1992; Tumanov, 1992; Sidorovich, 2000). Now, the species is extant in the eastern part of Belarus, in parts of Russia (Sidorovich, 2000; Wolsan, 1993) and in Romania, namely in the Danube delta (Gotea and Kranz, 1999). In contrast, in the west, it is reported in only seven departments of the south-westernmost part of France (Maizeret et al., 1995), as well as in the high valley of the river Ebra (Spain) (Ruiz-Olmo and Palazón, 1991).

In fact, it is possible that European mink were never distributed across most of western Europe during the Holocene. The fossil record is sparse (Davison et al., 2000), with the only confirmed records being an undated, probably Holocene, skull from Moscow District, another Holocene specimen from the Netherlands, material from the Polish site of Biskupin, and from the Romanian site of the 'La Adam' Cave (references in Davison et al., 2000). Although there is no direct evidence, there has also been some debate as to whether the western population is a recent introduction. The species was not recorded in France until the first half of the 19th century, when mink were already declining in Central Europe (de Bellefroid, 1999). One explanation for this is that local hunters or early naturalists did not distinguish between European mink and polecats (*M. putorius*).

Recently, many European species of mammals have been investigated using phylogeographic methods

* Corresponding author. Fax: +32-4-366-5010.

E-mail address: johan.michaux@ulg.ac.be (J.R. Michaux).

(Avisé, 2000). In most cases, a strong geographic structure of the genetic variability has been found throughout Europe, with different lineages arising from putative refugia in Iberia (Spain and Portugal), Italy, the Balkans and the Caucasus (Michaux et al., 1996, 1998, 2003; Santucci et al., 1998; Taberlet et al., 1998; Libois et al., 2001; Hewitt, 1999). Yet some species, especially wide-ranging carnivores, exhibit little geographic structure between the proposed refugia. This seems to be true for mustelids especially, with low variation and few lineages reported in polecats, pine martens (*Martes martes*; Davison et al., 2001), otters (*Lutra lutra*, Cassens et al., 2000), and wolverines (*Gulo gulo*, Walker et al., 2001), though no mustelid has been intensively sampled across its whole range.

Understanding the population history of European mink is a key part in the conservation effort, with studies ever more urgent because of the continuing decline of the species, and captive breeding/reintroduction programmes that are already underway. Locally, the French restoration plan (Anonymous, 1999) has proposed that captive-bred individuals are released into the wild as a reinforcement measure. In this circumstance, the choice of the animals to be bred is of a great importance. Indeed, if the western population is genetically distinct from the eastern ones, and if the animals are locally adapted, outbreeding depression (Lynch, 1991) could result. The definition of precise “management units” for European mink (Avisé, 2000), based on genetic markers is therefore of prime importance for the conservation of the species. On a European scale, the captive breeding programme has the stated aim to “maintain in European Zoos and other breeding facilities a population capable to maintain 90% of its heterozygosity for 100 years” (www.lutreola.ee/index.html). Captive-bred mink have already been released on Hiiumaa Island (Estonia). Thus, it is imperative that informed decisions are made regarding their management (in this case: restocking or reintroduction), based at least in part on genetic data.

A few studies have begun to investigate genetic variation in European mink, confirming that their superficial resemblance to American mink (*M. vison*) is a result of convergent evolution (Davison et al., 1999, 2000; Kurose et al., 2000; Hosoda et al., 2000). Mitochondrial studies have suggested that European mink is most closely related to the polecat (*Mustela putorius*) or to the steppe polecat (*Mustela eversmanni*), though the similarity may be a consequence of hybridization, in the late Pleistocene or Holocene (Davison et al., 2000). A nuclear DNA study seems to confirm this (Sato et al. in press).

For population genetic analysis, a cytochrome *b* fragment from 30 eastern European mink and seven Spanish animals has been sequenced previously, along with a more restricted D-loop fragment sample (Davison et al., 2000). The resulting phylogeny did not resolve the rela-

tionships of the species, probably due to hybridization and also because of low sequence variation. Whereas cytochrome *b* is useful in resolving some taxonomic groupings (Koepfli and Wayne, 1998; Kurose et al., 2000), its evolution is too slow (Flynn and Nédball, 1998) for intra-specific studies of mustelids.

In this study, we use the complete mitochondrial D-Loop region to investigate variation across a large part of the extant range of European mink, including the first samples from France and Romania. There were two main aims: to use the genetic information to advise on the conservation of European mink, and to better understand their history through the Pleistocene. Specifically, we would like to know whether the French/Spanish population should be managed separately from the Eastern populations.

2. Methods

2.1. Samples

A total of 43 European mink were studied, 23 from France, four from Spain, two from the Danube delta, three from Estonia, two from Belarus (Vitebsk) and nine from Russia (Tver and Pskov). The references and the geographic origin of these specimens are given in Table 1. They were compared with 10 polecats (*M. putorius*), two steppe polecats (*M. eversmanni*) and two black-footed ferrets (*M. nigripes*).

2.2. DNA methods

DNA was extracted from ethanol-preserved tissue as described by Sambrook et al. (1989). French samples were taken from the *Mustela* tissue collection of Dr. R. Rosoux, the GREGE samples (Groupe de recherches et d'études pour la gestion de l'environnement), and Romanian samples were provided by Dr. A. Toman. The remaining samples were described previously in Davison et al. (2000). All samples were taken either from road-killed European mink (muscle) or from live specimens (ear piece) caught, marked and released in the wild.

The complete D-Loop was amplified using specific primers L0ML (5'-TAT TCT AAC TAA ACT ATT CCC TG-3') and EML (5'-CTA TAG ATG TRT TTA TAA CCC-3') designed by J.R. Michaux. A portion of cytochrome *b* (450 bp of the 5' region) was also amplified by modifying the Universal PCR primers L7 (5'-ACC AAT GAC ATG AAA AAT CAT CGT T-3') and H8 (5'-ACA TGA ATY GGA GGY CAA CCW G-3') originally described by Kocher et al. (1989). Amplification reactions were carried out in 2 × 50 µl volumes including 25 µl of each 2 µM primer, 20 µl of 1 mM dNTP, 10 µl of 10× reaction buffer, 10 µl of purified

Table 1
Geographic distribution and references of *Mustela* tissues used for the experiments

| Geographic origin: | Total No. of animals | (see Figs. 1 and 2) | Tissue sample numbers or Genbank access (for sequenced samples) | |
|----------------------------|----------------------|---------------------|---|-----------------------------|
| <i>Mustela lutreola</i> | | | | |
| France | Bruges | 2 | F1 | 020500, 001G1 |
| | Audenge | 1 | F2 | 60400 |
| | Roquebrune | 1 | F3 | 003D |
| | St Martial sur le Né | 2 | F4 | V1, V2 |
| | Pont de Martrou | 1 | F5 | VF4 |
| | La Clisse | 1 | F6 | VF5 |
| | Crazannes | 1 | F7 | VM8 |
| | Orx | 2 | F8 | 003Orx, 004Orx |
| | Uzeste | 1 | F9 | 004C |
| | St Léger de Balson | 1 | F10 | 006C |
| | Ambleville | 2 | F11 | 003CHA, 004CHA |
| | Lachaise | 2 | F12 | 005CHA, 006CHA |
| | Tocanne St Apre | 1 | F13 | B |
| | Villefranque | 1 | F14 | C |
| | Belin-Beliet | 1 | F15 | D |
| | St Médard en Jalles | 1 | F16 | E |
| | Pont de Martrou | 1 | F17 | VM6 |
| | Blaye | 1 | F18 | VF7 |
| Spain | La Rioja | 2 | Sp1 | Mulp 55, Mulp 56 |
| | Navarra | 1 | Sp2 | Mulp 50 |
| | Gipuzkoa | 1 | Sp3 | AF207725 |
| Romania | Danube delta | 2 | Ro1 | JRM-698, JRM-699 |
| Estonia | Tallin | 2 | Est1 | JRM-659, JRM-660 |
| | Unknown locality | 1 | Est | AF207723 |
| Bielorussia | Vitebsk | 2 | Bel | Mulu 3, AF207724 |
| Russia | Tver | 7 | Rus1 | AF207720, Mulp 8 to Mulp 13 |
| | Pskov | 2 | Ru2 | Mulp 6, Mulp 7 |
| <i>Mustela putorius</i> | | | | |
| Spain | La rioja | 1 | Sp1 | AF207726 |
| | Gipuzkoa | 1 | Sp3 | Mulp 53 |
| Portugal | Unkown locality | 1 | Por | MPU 27 |
| Estonia | Unknown locality | 2 | Est | Mulp 2, Mulp 3 |
| Russia | Unknown locality | 1 | Rus | Mulp 41 |
| | Tver | 1 | Rus2 | Mulp 35 |
| ? | | 1 | | AF207717 |
| ? | | 1 | | AF207718 |
| ? | | 1 | | AF068570 |
| <i>Mustela eversmannii</i> | | | | |
| Mongolia | E. Inner | 1 | | M. Evers. 1 |
| Serbia | North West region | 1 | | M. Evers. 2 |
| <i>Mustela nigripes</i> | | | | |
| USA | Unknown locality | 2 | | M. nigripes 1 and 2 |

water and 0.2 µl of 5 U/µ Promega Taq DNA polymerase. Approximately 200 ng of DNA extract (10 µl) was used per PCR amplification. PCR was performed using an MJ Research PTC100 thermal cycler, employing 33 cycles (20 s at 94 °C, 30 s at 50 °C and 1 min 30 s at 68 °C) with a final extension cycle of 10 min at 68 °C. PCR products were then purified using the Ultra-free DA Amicon kit (Millipore) and directly sequenced. Both strands were sequenced using a BigDye terminator

(Applied Biosystems) sequencing kit on an ABI 310 (Applied Biosystems) automated sequencer.

The newly determined sequences were compared with five European mink (AF207720, AF207721, AF207723, AF207724 and AF207725) and four polecat (AF068570, AF207717, AF207718 and AF207726) partial D-Loop sequences available in Genbank (Table 1) using the ED editor (MUST package; Philippe, 1993). The hypervariable C_nT_n region (Davison et al., 2000) and an 11 bp

minisatellite were excluded from phylogenetic analysis, because it was not possible to unambiguously align all individuals.

2.3. Analyses

The aligned sequences were analysed by distance (neighbour joining, NJ; Saitou and Nei, 1987), maximum parsimony (MP) (Fitch, 1971) and maximum likelihood methods. The General Time Reversible (GTR) model and Kimura two-parameter (K2P) estimator were used for the calculation of genetic distances. The GTR estimator was chosen as it is the more general model of sequence evolution which consider six parameters for its probability matrix corresponding to each possible substitution. To take into account differences of substitution rates across sites, the GTR analysis was performed assuming a gamma distribution at eight categories. The alpha parameter (Yang, 1996) and the proportion of invariant sites (I) were estimated with the maximum-likelihood method in PAUP 4.0b8. Maximum parsimony (MP: heuristic search; TBR branch swapping option) and maximum likelihood (ML: GTR model of sequence evolution) analyses were also conducted using PAUP 4.0b8 (Swofford, 1998). The robustness of inferences was assessed by bootstrap resampling (BP) (1000 random repetitions for MP and distance analyses, and 100 for ML).

A Bayesian approach to phylogeny reconstruction (Yang and Rannala, 1997; Huelsenbeck et al., 2001) was also used, implemented in MrBayes 2.01 (Huelsenbeck and Ronquist, 2001). Metropolis-coupled Markov chain Monte Carlo sampling was performed with four chains that were ran for 500,000 generations, using default model parameters as starting values. Bayesian posterior probabilities were picked from the 50% majority rules consensus of trees sampled every 20 generations, after removing trees obtained before chains reached apparent stationarity (“burn in” determined by empirical checking of likelihood values).

A minimum spanning network was constructed using the MINSPNET algorithm available in the Arlequin 2.0 program (Schneider et al., 2000) as this method is very useful when sequences are closely related.

Nucleotide (π) and Haplotype (h) diversities, were estimated using the DNASP program (Rozas and Rozas, 1997). Calculations were performed on the main data matrix, including 57 animals.

3. Results

3.1. Sequences

The complete sequence of the mitochondrial D-Loop region was obtained for 38 European mink from 25

different localities, six European polecats from five regions, two steppe polecats and two black-footed ferrets (*M. nigripes*), used as outgroup (Table 1). These sequences have been deposited in the EMBL gene bank under accession numbers AJ548474 to AJ548496. They were aligned with the D-Loop sequences already available in GenBank. A 450 base pair fragment of the mitochondrial cytochrome *b* gene was also sequenced from 26 European mink from France and Spain.

3.2. Phylogenetic analyses

Cytochrome *b* sequences of different Mustelidae species: *Martes foina* (AF448245, ABO51250), *Martes martes* (AF448241, ABO51253), *Mustela vison* (ABO26106), *Mustela sibirica* (ABO51289, ABO51288), *Mustela putorius* (AF207716, AF207715), *Mustela nivalis* (ABO51272, ABO51271), *Mustela nigripes* (ARF068543), *Mustela lutreola* (ABO51263, ABO26105), *Mustela eversmannii* (ABO51261, ABO26102) and *Mustela erminea* (ABO51266, ABO51259) were used to determine the most suitable outgroup for the D-Loop analyses. Phylogenetic analyses (data not shown) performed on the basis of these sequences showed that *M. nigripes* is the most suitable outgroup to study mtDNA relationships within *M. lutreola* and *M. putorius*, as expected from earlier studies (Davison et al., 2000).

A first set of analyses was performed using partial D-Loop sequences, including those from Genbank. The aligned data matrix includes 57 *Mustela* (*lutreola*, *putorius*, *eversmannii* and *nigripes*) specimens and 254 sites, 40 of them variable and 27 phylogenetically informative. Thirty-one D-Loop haplotypes remained after removal of the C_nT_n array. The mean transition to transversion ratio is 2.29 and the nucleotide frequencies are: C 26.3%, T 30.5%, A 30.7%, G 12.5%. The Kimura two-parameter phylogeny (Fig. 1) shows two major clades: the first one corresponding to the polecat, *M. putorius*, and the second subdividing into two monophyletic groups of *M. lutreola* and *M. eversmannii*. The Bayesian, ML and MP (one most parsimonious tree, L = 60 steps; CI = 0.65; RI = 0.85) analyses yielded phylogenies of identical structure. Bootstrap values and Bayesian probabilities (BaP) resulting from these analyses are indicated in Fig. 1. Unfortunately, the major groups are not well supported, with bootstrap support (BP) around 50% (Fig. 1).

The D-loop sequences of French and Spanish mink are nearly identical, with variation restricted to the hypervariable C_nT_n array (either seven or eight thymine residues).

To confirm these results, a second set of analyses was performed, using the complete mitochondrial D-Loop region, with two *M. nigripes*, two *M. eversmannii*, six *M. putorius* and 38 *M. lutreola*. Again, all the French and Spanish European animals appear extremely similar.




Fig. 1. Consensus tree derived from the analysis of a partial sequence of the mitochondrial control region for 57 European minks, polecats and black footed ferrets (used as outgroup). Each haplotype is identified by its accession number in GenBank or by the letters DL + a specific number. The locality codes (see Table 1) are also given for each haplotype. For each node supported by a least 50% of bootstrap value, the different robustness are indicated as followed: $\frac{\text{Neighbour joining/maximum likelihood}}{\text{Maximum parsimony/Bayesian probability}}$.

They shared the same haplotype with the exception of the C_nT_n array (see earlier) and a variable (from 11 to at least 23) number of copies of an 11 bp minisatellite in the hypervariable region R. It was impossible to determine the exact number of minisatellite copies, when $n > 23$, due to sequencing difficulties. This region was not considered in any further phylogenetic analysis.

After removal of the C_nT_n array and the minisatellite repeats, the final data matrix involved 729 sites, of which 63 were variable and 41 phylogenetically infor-

mative. The neighbour joining tree is shown in Fig. 2, obviously similar in structure to the tree in Fig. 1. The group corresponding to *M. putorius* is well supported (BP values for NJ: 79%, ML: 59, MP: 80% and BaP: 0.99). The second group of *M. lutreola* and *M. eversmannii* appears less robust (BP values for NJ: 53%, ML: 50% and BaP: 0.55). Within this group, *M. lutreola* and *M. eversmannii* are monophyletic, with stronger support compared with the first analysis (Fig. 1).




Fig. 2. Consensus tree derived from the analysis of the complete mitochondrial control region sequences for 37 European minks, polecats and black footed ferrets (used as outgroup). Each haplotype is identified by the letters DL + a specific number. The locality codes (see table 1) are also given for each haplotype. For each node supported by a least 50% of bootstrap value, the different robustness are indicated as followed:

Neighbour joining/maximum likelihood
Maximum parsimony/Bayesian probability

To confirm absolutely the lack of variation among all the French/Spanish European mink, we sequenced a 450 bp fragment of the mitochondrial cytochrome *b* gene in 23 French and three Spanish animals. All the sequences obtained were identical to a shorter sequence available in Genbank (AF207725; haplotype C11; Davison et al., 2000).

3.3. Genetic distances between the mustelids

The levels of genetic divergence (distance analysis, K2P estimator) observed between the four *Mustela* species are summarised in Table 2. The values range from 3.2 to 4.7% K2P distance.

3.4. Genetic structure

Nucleotide (π) and haplotype (*h*) diversities were calculated for each population of *M. lutreola* (Table 3)

Table 2
Mean levels of genetic divergence (in%) between the studied *Mustela* species, obtained in the distance analysis (K2P estimator)

| | <i>Mustela nigripes</i> | <i>Mustela eversmannii</i> | <i>Mustela putorius</i> |
|----------------------------|-------------------------|----------------------------|-------------------------|
| <i>Mustela eversmannii</i> | 4.7 | | |
| <i>Mustela putorius</i> | 3.4 | 4.1 | |
| <i>Mustela lutreola</i> | 3.9 | 3.5 | 3.2 |

Table 3
Genetic variability observed within the three populations of *M. lutreola* and the one of *M. putorius*

| | Sample size | Number of haplotypes | Genetic divergence within each group (% K2P distance) | Nucleotide diversity (Pi) | Standard deviation | Haplotype diversity (Pi) | Standard deviation |
|--------------------|-------------------------------------|----------------------|---|---------------------------|--------------------|--------------------------|--------------------|
| <i>M. putorius</i> | 10 | 10 | 2.4 | 0.0108 | 0.0015 | 0.978 | 0.054 |
| <i>M. lutreola</i> | Russian + bielorrussian populations | 14 | 1.5 | 0.0197 | 0.0025 | 0.978 | 0.035 |
| | Romanian population | 2 | 0.4 | 0.0039 | 0.0019 | 1 | / |
| | Western european population | 27 | 1 | 0 | 0 | 1 | / |

Animals from Russia and Belarus have a high nucleotide and haplotype diversity, compared to the French and Spanish samples, which share a single mitochondrial type. The mismatch distribution of pairwise differences shows a signature of population growth (Luikart et al., 2001), a bell-shape, similar to that found in *M. putorius* (Fig. 4). The network analysis (Fig. 3) shows a similar distribution of the samples.

4. Discussion

4.1. The phylogeography of European mink

As previously observed by Davison et al. (2000), the intraspecific genetic variability of *M. lutreola* is low (see Table 3). This is consistent with other European carnivores, especially mustelids, like wolverines (Walker et al., 2001), otters (Cassens et al., 2000; Morales, 2002), polecats, pine martens (Davison et al., 2001) and wolves (Vila et al., 1999). However, the low variation contrasts with the situation in other mammals, such as brown bears (*Ursus arctos*) (Taberlet et al., 1998), hedgehogs (*Erinaceus europaeus*) (Seddon et al., 2001), shrews (*Sorex araneus*, *S. minutus* and *Crocidura suaveolens*) (Taberlet et al., 1994, 1998; Bilton et al., 1998), and the rodents *Apodemus sylvaticus* (Michaux et al., 2003; Libois et al., 2001), *Myopus schisticolor* (Federov et al., 1996), *Arvicola* sp. (Taberlet et al., 1998), *Microtus agrestis* and *Clethrionomys glareolus* (Jaarola and Tegelström, 1995; Bilton et al., 1998). All of these species are characterised by divergent mtDNA lineages which are distributed over distinct geographic areas.

The absence of highly divergent lineages in the European mink, as well as a mismatch distribution that is consistent with an expanding population (Fig. 4), is an indication that the present-day population is the result of a colonisation from a single refugium, after the last glaciation. Unfortunately, the majority of the southern central European populations of the mink have long since become extinct (Youngman, 1982; de Bellefroid and Rosoux, 1998), and the analysis was limited to two Romanian animals. This, coupled with a poor fossil record, means that a more accurate reconstruction of the postglacial recolonisation is not possible.

However, the phylogeny supports earlier work (Davison et al., 2000), suggesting that European mink and steppe polecat are closely related, based on the mitochondrial D-loop region. This is in contrast to a recent nuclear gene phylogeny. Sato et al. (in press) compared mustelid cytochrome *b* (complete sequence) and nuclear interphotoreceptor retinoid binding protein phylogenies. Overall, the three phylogenies are congruent (including D-loop also), with the exception of the placement of European mink, which appears more closely related to a *M. itatsi*–*M. sibirica*–*M. putorius* clade in the nuclear phylogeny.

4.2. The history of French and Spanish European mink

In contrast to the situation in other species of mammals, where the southern-most refugial populations have the greatest diversity (Michaux et al., 2003), European mink mtDNA diversity is highest in East European populations, with an almost complete lack of variation in French and Spanish animals. While this needs to be confirmed using nuclear markers, such as microsatellites, it suggests that a few individuals (possibly a single female) established this population, possibly as an anthropogenic introduction. However, an early Holocene origin is equally conceivable, not least because the D-loop mutation rate is completely unknown in this group. Leptokurtic dispersal from a refugium, where a few long-distance migrant individuals establish populations in advance of a colonisation wave, is expected to produce patches of relative genetic homogeneity such as those reported here (Ibrahim et al., 1996).

4.3. Implications for the conservation of European mink

Awise (2000) defined the concept of a Management Unit as “any population that exchanges so few migrants with others as to be genetically distinct from them normally will be demographically independent at the present time”. In this way, he noted that “even shallow matrilineal subdivisions can be relevant to conservation efforts”. Certainly, specific haplotypes are found in the East European and French and Spanish populations at different frequencies (insufficient samples were investigated in Romania), with a single haplotype predominating in France and Spain.

Ideally, this mtDNA study should be augmented by an analysis of nuclear markers and fitness related phenotypic differences (Hedrick, 1999), before defining Management Units. In the meantime, the captive breeding program is

proceeding. Following the precautionary principle, we recommend that mink from the three geographically separate populations (France/Spain, Romania, Eastern Europe) should be managed separately, at least for the





Fig. 3. A minimum spanning network constructed using mitochondrial control region sequences. The identity of the haplotypes (see Fig.s 1 and 2) and their geographic origin (see Table 1) are indicated. Numbers in italics correspond to the mutational steps observed between haplotypes.

Eastern lineage of *Mustela lutreola*



Mustela putorius lineage




Fig. 4. Mismatch distribution for mtDNA types from the eastern genetic lineage of *Mustela lutreola* and the *M. putorius* group. The expected frequency is based on a population growth-decline model (for the groups, respectively: θ initial = 0, 2.4, 0 and 0, θ final = 1000, τ = 4.4, 9.4, 2.7 and 1.6), determined using the DNASP v3.5 program (Rozas and Rozas, 1997).

moment. It is also possible that the animals are locally adapted, so that outbreeding depression (Lynch, 1991) could result from some mating combinations, especially once the animals were released into the wild.

Acknowledgements

We thank the Natural History Museum of La Rochelle as well as Dr. Y. De Beaulieu (DIREN Aquitaine) for their help in funding this study. Tissue samples were provided by Dr. Pascal Fournier (GREGE), Dr Ales Toman (Pavlov Biological Station, Czech Republic) Mrs C. de la Panouse, (Zoological park of Thoiry (F-77), Dr A. Kranz, Dr T. Maran (Tallin Zoo), the department services of the Office National de la Chasse et de la Faune Sauvage of Charente, Charente-Maritime, Gironde and Landes, Nature-Environnement 17, and the League for bird protection (LPO). Spanish samples were provided by Iñaki Irizar, Idoia Villate, Jorge González-Esteban, Juan Carlos Ceña, Alfonso Ceña, Ivan Moya, and Santiago Palazón Miñano. This conservation genetics project would not have been possible without their valuable assistance. It was initiated at the Euro-American mammal conference in Santiago de Compostela (1998) by Huw Griffiths. This paper is dedicated to his memory.

This work was supported by a Belgian FNRS fellowship to J.R. Michaux (mandat “Chargé de Recherches”), a financial grant of the Belgian FNRS (crédits aux chercheurs), and financial support of the University of Liège (Fonds spéciaux de recherche 2000–2002), of the Natural History Museum of La Rochelle as well as of the DIREN Aquitaine.

References

- Anonymous, 1999. Plan de restauration du vison d'Europe en France. Ministère de l'Aménagement du Territoire et de l'Environnement, Paris.
- Avise, J.C., 2000. Phylogeography. The History and Formation of Species. Harvard University Press, Cambridge, MA.
- Baillie, N., Groombridge, C., 1996. 1996 IUCN Red List of Threatened Animals. IUCN, Gland.
- Bilton, D.T., Mirol, P.M., Mascheretti, S., Fredga, K., Zima, J., Searle, J.B., 1998. Mediterranean Europe as an area of endemism for small mammals rather than a source for northwards postglacial colonization. Proceedings Royal Society London Biological Sciences 265, 1219–1226.
- Camby A., 1990. Le vison d'Europe (*Mustela lutreola* (Linnaeus, 1761)). Encyclopédie des Carnivores de France, n° 13. Société française pour l'Etude et la Protection des Mammifères, Nort-sur-Erdre.
- Cassens, I., Tiedemann, R., Suchentrunk, F., Hartl, G.B., 2000. Mitochondrial DNA variation in the European otter (*Lutra lutra*) and the use of spatial autocorrelation analysis in conservation. Journal of Heredity 91, 31–35.
- Davison, A., Birks, J.D.S., Griffiths, H.I., Kitchener, A.C., Biggins, D., Butlin, R.K., 1999. Hybridization and the phylogenetic relationship between polecats and domestic ferrets in Britain. Biological Conservation 87, 155–161.
- Davison, A., Griffiths, H.I., Brookes, R.C., Maran, T., MacDonald, D., Sidorovich, V.E., Kitchener, A.C., Irizar, I., Villate, I., Gonzalez-Esteban, J., Cena, J.C., Cena, A., Moya, I., Palazon Minanon, S., 2000. Mitochondrial DNA and palaeontological evidence for the origins of the endangered European mink, *Mustela lutreola*. Animal Conservation 4.
- Davison, A., Birks, J.D., Brookes, R.C., Messenger, J.E., Griffiths, H.I., 2001. Mitochondrial phylogeography and population history of pine martens *Martes martes* compared with polecats *Mustela putorius*. Molecular Ecology 10, 2479–2488.
- de Bellefroid, M.N., 1999. Étude bio-géographique de l'évolution de la population de vison européen, *Mustela lutreola*, en France. Thèse DDR, Univ. Rennes.
- de Bellefroid, M.N., Rosoux, R., 1998. Le “vison du Poitou”, un hôte des zones humides menacé dans le Centre-Ouest atlantique. Annales de la société des Sciences Naturelles Charente-Maritime 8, 865–879.
- Federov, V., Jaarola, M., Fredga, K., 1996. Low mitochondrial DNA variation and recent colonization of Scandinavia by the wood lemming *Myopus schisticolor*. Molecular Ecology 5, 577–581.
- Fitch, W.M., 1971. Towards defining the course of evolution: Minimum change for a specific tree topology. Systematic Zoology 20, 406–416.
- Flynn, J.J., Nedbal, M.A., 1998. Phylogeny of the Carnivora (Mammalia): congruence vs incompatibility among multiple data sets. Molecular Phylogenetics Evolution 9, 414–426.
- Gotea, V., Kranz, A., 1999. The European mink (*Mustela lutreola*) in the Danube delta. Small Carnivore Conservation 21, 23–25.
- Hedrick, P.W., 1999. Perspective: highly variable loci and their interpretation in evolution and conservation. Evolution 53, 313–318.
- Hewitt, G.M., 1999. Post-glacial recolonisation of European biota. Biological Journal of the Linnean Society 58, 87–112.
- Hosoda, T., Suzuki, H., Harada, M., Tsuchiya, K., Han, S.H., Zhang, Y.-P., Kryukov, A.P., Lin, L.K., 2000. Evolutionary trends of the mitochondrial lineage differentiation in species of genera *Martes* and *Mustela*. Genes Genetics Systematics 75, 259–267.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17, 754–755.
- Huelsenbeck, J.P., Ronquist, F., Nielsen, R., Bollback, J.P., 2001. Bayesian inference of phylogeny and its impact on evolutionary biology. Science 294, 2310–2314.
- Ibrahim, K.M., Nichols, R.A., Hewitt, G.M., 1996. Spatial patterns of genetic variation generated by different forms of dispersal during range expansion. Heredity 77, 282–291.
- Jaarola, M., Tegelstrom, H., 1995. Colonization history of north European field voles (*Microtus agrestis*) revealed by mitochondrial DNA. Molecular Ecology 4, 299–310.
- Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Pääbo, S., Villablanca, F.X., 1989. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. Proceedings of the National Academy of Sciences, USA 86, 6196–6200.
- Koepfli, K.P., Wayne, P., 1998. Phylogenetic relationships of otters (Carnivora :Mustelidae) based on mitochondrial cytochrome b sequences. Journal Zoology London 246, 401–416.
- Kurose, N., Abramov, A.V., Masuda, R., 2000. Intrageneric diversity of the cytochrome b gene and phylogeny of Eurasian species of the genus *Mustela* (Mustelidae, Carnivora). Zoological Sciences 17, 673–679.
- Libois, R.M., Michaux, J.R., Ramalhinho, M.G., Maurois, C., Sara, M., 2001. On the origin and systematics of the northern African wood mouse (*Apodemus sylvaticus*) populations: a comparative study of mtDNA restriction patterns. Canadian Journal of Zoology 79, 1–9.
- Luikart, G., Gellie, L., Excoffier, L., Vigne, J.-D., Bouvet, J., Taberlet, P., 2001. Multiple maternal origins and weak phylogeographic structure in domestic goats. Proceedings of the National Academy of Sciences, USA 98, 5927–5932.

- Lynch, M., 1991. The genetic interpretation of inbreeding depression and outbreeding depression. *Evolution* 45, 622–629.
- Maizeret, C., Maurin, H., Migot, P., Lafontaine, L., 1995. État d'avancement du programme d'étude de la répartition du vison d'Europe en France. *Cahiers Éthologie* 15, 419–424.
- Maran, T., 1992. The European mink, *Mustela lutreola*, in protected areas in the former Soviet Union. *Small Carnivore Conservation* 7, 10–12.
- Michaux, J.R., Filippucci, M.G., Libois, R.M., Fons, R., Matagne, R.F., 1996. Biogeography and taxonomy of *Apodemus sylvaticus* (the woodmouse) in the Tyrrhenian region: enzymatic variations and mitochondrial dna restriction pattern analysis. *Heredity* 76, 267–277.
- Michaux, J.R., Libois, R., Ramalhinho, M.G., Maurois, C., 1998. On the mtDNA restriction patterns variation of the iberian wood mouse (*Apodemus sylvaticus*). comparison with other west mediterranean populations. *Hereditas* 129, 187–194.
- Michaux, J.R., Magnanou, E., Paradis, E., Nieberding, C., Libois, R.M., 2003. Mitochondrial phylogeography of the Woodmouse (*Apodemus sylvaticus*) in the Western Palearctic region. *Molecular Ecology* 12, 685–697.
- Morales, C.F., 2002. Phylogéographie de la loutre d'Europe (*Lutra lutra*) dans la partie occidentale de son aire de répartition. Mém. DES Sciences Pays en Développement, Univ. Liège.
- Philippe, H., 1993. MUST, a computer package for management utilities for sequences and trees. *Nucleic Acids Research* 21, 5264–5272.
- Rozas, J., Rozas, R., 1997. DNASP, version 2. 0: a novel software package for extensive molecular population genetic analysis. *Computer Application in the Biosciences* 13, 307–311.
- Ruiz-Olmo, J., Palazón, S., 1991. New information on the European and American mink in the Iberian Peninsula. *Mustelid Viverrid Conserv* 5, 13.
- Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4, 406–425.
- Sambrook, J., Fritsch, E.F., Maniatis, T., 1989. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Santucci, F., Emerson, B.C., Hewitt, G.M., 1998. Mitochondrial DNA phylogeography of European hedgehogs. *Molecular Ecology* 7, 1163–1172.
- Sato, J.J., Hosoda, T., Wolsan, M., Tsuchiya, K., Yamamoto, M., Suzuki, H. Phylogenetic relationships and divergence times among mustelids (Mammalia: Carnivora) based on nucleotide sequences of the nuclear interphotoreceptor retinoid binding protein and mitochondrial cytochrome *b* genes. *Zoological Science* (in press).
- Schneider, S., Roessli, D., Excoffier, L., 2000. Arlequin, Version 2.0: a Software for Population Genetic Data Analysis. Genetics and Biometry Laboratory, University of Geneva, Geneva.
- Seddon, J.M., Santucci, F., Reeve, N.J., Hewitt, G.M., 2001. DNA footprints of European hedgehogs, *Erinaceus europaeus* and *E. concolor*: Pleistocene refugia, postglacial expansion and colonization routes. *Molecular Ecology* 10, 2187–2198.
- Sidorovich, V.E., 2000. The ongoing decline of riparian mustelids (European mink, *Mustela lutreola*, polecat, *Mustela putorius* and stoat, *Mustela erminea*) in eastern europe: a review of the results to date and na hypothesis. In: Griffiths, H.I. (Ed.), *The Mustelids in a Modern World. Management and Conservation Aspects of Small Carnivore: Human Interactions*. Bachuys, Leiden, pp. 295–317.
- Swofford, D.L., 1998. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4.0b. Sinauer Associates, Sunderland, Massachusetts.
- Taberlet, P., Fumagalli, L., Hausser, J., 1994. Chromosomal versus mitochondrial DNA evolution: tracking the evolutionary history of the southwestern European populations of the *Sorex araneus* group (Mammalia, Insectivora). *Evolution* 48, 623–636.
- Taberlet, P., Fumagalli, L., Wust-Saucy, A.G., Cosson, J.F., 1998. Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* 7, 453–464.
- Vilá, C., Savolainen, P., Maldonado, J.E., Amorim, I.R., Rice, J.E., Honeycutt, R.L., 1997. Multiple and ancient origins of the domestic dog. *Science* 276, 1687–1689.
- Tumanov, I.L., 1992. The number of European mink (*Mustela lutreola*) in the eastern area and its relation to American mink. In: Schröpfer, R. et al. (Eds.), *Semi-aquatische Säugetiere. Materialien des 2. Internat. Symposium*. Martin Luther Universität, Halle-Wittenberg, pp. 329–336.
- Van Bree, P.J.H., Saint Girons, M.C., 1966. Données sur la répartition et la taxonomie de *Mustela lutreola* (Linnaeus, 1761) en France. *Mammalia* 30, 270–291.
- Walker, W.K., Vila, C., Landa, A., Linden, M., Ellegren, H., 2001. Genetic variation and population structure in Scandinavian wolverine (*Gulo gulo*) populations. *Molecular Ecology* 10, 53–63.
- Wolsan, M., 1993. Evolution des carnivores Quaternaires en Europe centrale dans leur contexte stratigraphique et paléoclimatique. *L'Anthropologie* 97, 203–222.
- Yang, Z., 1996. Among-site rate variation and its impact on phylogenetic analyses. *Trends Ecology Evolution* 11, 367–372.
- Yang, Z., Rannala, B., 1997. Bayesian phylogenetic inference using DNA sequences: a Markov chain Monte carlo method. *Molecular Biology Evolution* 14, 717–724.
- Youngman, P.M., 1982. Distribution and systematics of the European mink, *Mustela lutreola*. *Acta Zoologica Fennascandia* 166, 1–48.