

# Zooming into the broad line region of the gravitationally lensed quasar QSO 2237+0305 $\equiv$ the Einstein Cross $\star$

## III. Determination of the size and structure of the C IV and C III] emitting regions using microlensing

D. Sluse<sup>1</sup>, R. Schmidt<sup>1</sup>, F. Courbin<sup>2</sup>, D. Hutsemékers<sup>3\*\*</sup>, G. Meylan<sup>2</sup>, A. Eigenbrod<sup>2</sup>, T. Anguita<sup>4,5</sup>, E. Agol<sup>6</sup>, and J. Wambsganss<sup>1</sup>

<sup>1</sup> Astronomisches Rechen-Institut am Zentrum fuer Astronomie der Universitaet Heidelberg Moenchhofstrasse 12-14, 69120 Heidelberg, Germany

<sup>2</sup> Laboratoire d'Astrophysique, Ecole Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, 1290 Versoix, Switzerland

<sup>3</sup> Institut d'Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août 17, B5c, 4000 Liège, Belgium

<sup>4</sup> Centro de Astro-Ingeniería, Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Casilla 306, Santiago, Chile

<sup>5</sup> Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany

<sup>6</sup> Astronomy Department, University of Washington, Box 351580, Seattle, WA 98195, USA

Received ... ; accepted ...

### ABSTRACT

**Aims.** We aim to use microlensing taking place in the lensed quasar QSO 2237+0305 to study the structure of the broad line region and measure the size of the region emitting the C IV and C III] lines.

**Methods.** Based on 39 spectrophotometric monitoring data points obtained between Oct. 2004 and Dec. 2007, we derive lightcurves for the C IV and C III] emission lines. We use three different techniques to analyse the microlensing signal. Different components of the lines (narrow, broad and very broad) are identified and studied. We build a library of simulated microlensing lightcurves which reproduce the signal observed in the continuum and in the lines provided only the source size is changed. A Bayesian analysis scheme is then developed to derive the size of the various components of the BLR.

**Results.** 1. The half-light radius of the region emitting the C IV line is found to be  $R_{\text{CIV}} \sim 66^{+110}_{-46}$  lt-days =  $0.06^{+0.09}_{-0.04}$  pc =  $1.7^{+2.8}_{-1.1} 10^{17}$  cm (at 68.3% CI). Similar values are obtained for C III]. Relative sizes of the V-band continuum and of the carbon line emitting regions are also derived with median values of  $R^{\text{line}}/R^{\text{cont}}$  in the range [4,29], depending of the FWHM of the line component.

2. The size of the C IV emitting region agrees with the Radius-Luminosity relationship derived from reverberation mapping. Using the virial theorem we derive the mass of the black hole in QSO 2237+0305 to be  $M_{\text{BH}} \sim 10^{8.3 \pm 0.3} M_{\odot}$ .

3. We find that the C IV and C III] lines are produced in at least 2 spatially distinct regions, the most compact one giving rise to the broadest component of the line. The broad and narrow line profiles are slightly different for C IV and C III].

4. Our analysis suggests a different structure of the C IV and Fe II+III emitting regions, with the latter being produced in the inner part of the BLR or in a less extended emitting region than C IV.

**Key words.** Gravitational lensing: micro, strong, quasars: general, quasars: emission lines, quasars: individual QSO 2237+0305, line: formation

## 1. Introduction

We know that quasars and active galactic nuclei (AGN) are powered by matter accreted onto a supermassive black hole. The accretion of material in the direct vicinity of the central black hole releases most of the quasar energy, in form of power-law continuum emission. Ionised gas surrounds the central accretion disc and gives rise to broad emission lines, which are used as footprints allowing the identification and classification

$\star$  Based on observations made with the ESO-VLT Unit Telescope # 2 Kueyen (Cerro Paranal, Chile; Proposals 073.B-0243(A&B), 074.B-0270(A), 075.B-0350(A), 076.B-0197(A), 177.B-0615(A&B), PI: F. Courbin).

\*\* Maître de Recherches du FNRS

of quasars. Our knowledge of the kinematics and physical conditions prevailing in the BLR gas remain elusive, especially because the nuclear region of quasars is yet spatially unresolved with existing instrumentation.

Current insights on the BLR come from various kind of studies: empirical modeling of the line shape with kinematical models, use of photo-ionization codes to reproduce the observed flux ratios between spectral lines, spectropolarimetric observations, statistical study of the width and asymmetry of the lines, use of the principal component analysis technique, and velocity resolved reverberation mapping (e.g. Boroson & Green 1992, Sulentic et al. 2000, Smith et al. 2005, Marziani et al. 2006, Zamfir et al. 2008, Gaskell 2009, 2010b, Bentz et al. 2010). Despite the development and many successes of these methods, as briefly summarized hereafter, we still do not completely understand the structure and kinematics of the BLR. The microlensing of the broad emission lines in multiply-imaged lensed AGNs provides us with a powerful alternative technique to look at the BLR, measure its size, even in high luminosity distant quasars, and get hints of the structure and geometry of both emission and intrinsic absorption within the BLR (e.g. Schneider & Wambsganss 1990, Hutsemékers et al. 1994, Lewis & Belle 1998, Abajas et al. 2002, Popović et al. 2003, Sluse et al. 2007, 2008, Hutsemékers et al. 2010).

### 1.1. Phenomenology of the line profiles

Our primary clue on the BLR comes from the shape of the broad emission lines. The largest (and highest S/N) studies of broad emission lines have focused on two lines:  $H\beta$   $\lambda 4863$  and C IV  $\lambda 1549$  (Sulentic et al. 2000 for a review). Of direct interest for the present work is C IV  $\lambda 1549$ . The C IV profile shows a broad variety of shapes from strongly asymmetric to symmetric (e.g. Wills et al. 1993, Baskin & Laor 2005), has an equivalent width anti-correlated with its intensity (Francis et al. 1992), and shows larger variability in the wings than in the core (Wilhite et al. 2006). In addition, the C IV line is also found to be systematically blueshifted by several hundred to a few thousand kilometers per second compared to the low ionisation lines (Gaskell 1982, Corbin 1990, Vanden Berk et al. 2000). The analysis of about 4000 SDSS quasars by Richards et al. (2002) suggests that this shift is caused by a lack of flux in the red wing of the line profile and correlates with the quasar orientation. Variability studies and emission line decomposition techniques (principal component analysis and analytical fitting) indicate that the region emitting C IV could be built up with 2 components, a “narrow” emission core of  $\text{FWHM} \sim 2000 \text{ km s}^{-1}$  emitted in an intermediate line region (ILR) possibly corresponding to the inner part of the narrow line region, and a very broad component (VBC) with  $\text{FWHM} \sim 7000 \text{ km s}^{-1}$  producing the line wings (Wills et al. 1993, Brotherton et al. 1994a, Sulentic 2000, Wilhite et al. 2006). In radio-quiet objects, the VBC is observed to be systematically blue-shifted by thousands of  $\text{km s}^{-1}$  w.r.t. to the narrow core (Brotherton et al. 1994a, Corbin 1995) suggesting it is associated with outflowing material. The ILR component disclosed in C IV is probably different from the one recently uncovered in  $H\beta$  (Hu

et al. 2008) but its exact nature and the physical conditions in this region are still debated (Brotherton et al. 1994b, Sulentic & Marziani 1999, Marziani et al. 2006).

The properties of the C III] emission line have received less attention in the literature, probably mostly because the line is blended with Al III  $\lambda 1857$ , Si III]  $\lambda 1892$  and with a Fe II+III  $\lambda 1914$  complex. In their study of the C IV and C III] emission, Brotherton et al. (1994a) find that the 2 lines often have different profiles and that, for some objects, a 2 components decomposition (i.e. narrow core+very broad wings) does not provide a good model of C III], a third component is required. They also find that the VBC needed to reproduce C III] has to be larger than the corresponding component of C IV. This supports the idea that the region emitting C III] is different from the one emitting C IV (Snedden & Gaskell 1999).

Recently, Marziani et al. (2010), based on the line profile decomposition of a small sample of AGNs selected in the 4D Eigenvector 1 context (e.g. Boroson & Green 1992, Zamfir et al. 2008), suggest that all the broad emission line profiles are composed of 3 components of variable relative intensity and centroid shift (from line to line in a given object and between objects). They suggest a classical unshifted broad component ( $\text{FWHM}=600\text{-}5000 \text{ km s}^{-1}$ ), a redshifted very broad component and a blue-shifted component mostly visible in the so-called Population A objects (i.e. objects with  $\text{FWHM} < 4000 \text{ km s}^{-1}$ ). Based on line ratios, they also tentatively infer that these components arise from different emitting regions.

The smoothness of the line profiles and the physical conditions derived from photo-ionization models allowed several authors to put constraints on the “structure” of the BLR gas. Two popular models remain. The first one considers that the BLR is a clumpy flow composed of small gas clouds, and the second one assumes a smooth gas outflow originating from an accretion disc (Elvis 2000, Laor 2007 and ref. therein). Several geometries have been considered for the BLR gas, the most popular ones are disc-like, spherical and biconical models (e.g. Chen & Halpern 1989, Robinson 1995). Murray and Chiang (1997) demonstrated that a continuous wind of gas originating from an accretion disc can successfully reproduce the profile and systematic blue-shift of the C IV emission line. Other indications that a fraction of the BLR material has a disc-like geometry comes from statistical studies of the broad line profiles in samples of (mostly radio-loud) AGNs (Vestergaard et al. 2000, McLure & Dunlop 2002, Jarvis & McLure 2006, Decarli et al. 2008, Risaliti et al. 2010), and from spectro-polarisation observations of Balmer lines in AGNs (Smith et al. 2005). Some of these studies also suggest the existence of a second, spherically symmetric component with keplerian motion. The spectropolarimetric observations of PG 1700+518 by Young et al. (2007) support a disc+wind model for the  $H\alpha$  emission. On the other hand, biconical models of the BLR seem needed to explain the variability of the rare double-peaked AGNs (i.e. AGNs showing emission lines with two peaks) and their polarisation properties (Sulentic et al. 1995, Corbett et al. 1998). These studies show that there is no consensus on the geometry of the BLR. From an observational and theoretical perspective, it seems nevertheless to depend on the ionisation degree of the line and on the radio properties of the object.

## 1.2. The Radius-Luminosity relationship

The “size” of the the broad line region  $R_{BLR}$  has been measured in about 40 AGNs by use of the reverberation mapping technique (e.g. Krolik et al. 1991, Horne et al. 2004). The empirical relation  $R_{BLR} \propto L^\alpha$  ( $\alpha \sim 0.5-0.6$ , Kaspi et al. 2005, Bentz et al. 2006), combined with the virial theorem, allows one to derive a relation linking the black hole mass  $M_{BH}$ , the AGN luminosity  $L$  and the FWHM of the emission line. This relation is one of the most popular methods used to measure black hole masses based on single epoch spectroscopic data and study their growth, evolution and correlation to other AGN properties.

The  $R_{BLR} \propto L^\alpha$  relation has been derived quite accurately for the broad component of the  $H\beta$  line, but only a few objects have  $R_{BLR}$  measurements for high ionisation lines like C IV (e.g. Kaspi et al. 2007). This is a severe problem for black hole mass measurements of high redshift objects. The use of the C IV line to derive black hole masses is desirable but faces several problems related to our understanding of the structure and geometry of the region emitting this line (see e.g. Marziani et al. 2006 for a review). The existence and contamination of a narrow emission component in C IV which could bias  $FWHM_{CIV}$  (Bachev et al. 2004) and the possible absence of virial equilibrium, especially in sources showing large blueshifts (Richards et al. 2002), are major concerns.

Velocity resolved reverberation mapping (Horne et al. 2004) should allow one to get insights on these problems, but the technique is still under development and has not yet been applied to the C IV line. Most of the recent advances in velocity resolved echo-mapping are based on the analysis of the  $H\beta$  emission line (Denney et al. 2009, Bentz et al. 2009, 2010). Although the kinematics of the Balmer gas was thought to be relatively simple, the technique provided puzzling results as both keplerian rotation, inflow and outflow signatures appear in various objects. It is still unclear whether this reveals a large variety of kinematics, a biased interpretation of the observed signal or a spurious signal introduced by observational artifacts. The past evidence of inflow and outflow signatures in NGC 5548 (Clavel et al. 1991, Peterson et al. 1991, Kollatschny & Dietrich 1996) might indicate that inflow/outflow signatures are not unambiguous. A possible explanation to the variety of observed signals might be off-axis illumination of the BLR (Gaskell 2010a, 2010b and references therein).

## 1.3. Microlensing in QSO 2237+0305


The many open questions concerning the BLR we outlined above motivate the interest in developing new techniques to probe this region. In this paper we investigate the constraints on the broad lines provided by means of the microlensing study of the lensed quasar QSO 2237+0305, for which long term spectro-photometric monitoring has been carried out (Eigenbrod et al. 2007, hereafter *Paper I*). The gravitational lens QSO 2237+0305, also known as “Huchra’s lens” or the “Einstein Cross”, was discovered by Huchra et al. (1985) during the Center for Astrophysics Redshift Survey. It consists of a  $z_s = 1.695$  quasar gravitationally lensed into

four images arranged in a cross-like pattern around the nucleus of a  $z_l = 0.0394$  barred Sab galaxy. The average projected distance of the images from the lens center is only  $700 \text{ pc} \sim 0.9''$ , such that the matter along the line of sight to the lensed images is mostly composed of stars. The symmetric configuration of the lensed images around the lens-galaxy bulge ensures a time-delay of less than a day between the lensed images, such that intrinsic flux variations should be seen quasi-simultaneously in the four images (Rix et al. 1992, Wambsganss & Paczyński 1994). Since the low redshift of the lens galaxy leads to a large relative transverse velocity between the observer, the lens and the source, the lensed images of QSO 2237+0305 are continuously flickering due to microlensing produced by the stars in the lens galaxy, at smaller timescales than in any other lens. The microlensing affecting the images of QSO 2237+0305 leads to large amplitude variations (Udalski et al. 2006), reaching up  $> 1$  mag, often accompanied by chromatic microlensing of the quasar continuum (Wambsganss & Paczyński 1991), used to study the accretion disc temperature profile (e.g. Kochanek 2004, Anguita et al. 2008, Eigenbrod et al. 2008=*Paper II*). Several studies have shown that the broad emission lines are also significantly affected by microlensing (Wayth et al. 2005, Metcalf et al. 2004, *Paper I*). Based on microlensing of C III] observed at one epoch in 2002, Wayth et al. (2005) derived a most likely size of the region emitting this line of  $0.06 h^{1/2} \text{ pc}$ . The work presented here aims at improving the measurement of the BLR size by using monitoring data instead of a single epoch measurement, and constraining the structure of the BLR.

The structure of the Paper is the following. In Sect. 2, we briefly summarize the data we use and explain the three different methods we apply to analyse the spectra. In Sect. 3, we apply these techniques to our data to understand how microlensing deforms the emission lines and to measure the flux ratios in several portions of the C IV and C III] emission lines. In Sect. 4, we explain the microlensing simulations we developed to derive the size of the broad emission lines and continuum emission region. In Sect. 5 we present our results: measurement of the size of the BLR and of the continuum region, comparison of our BLR size with reverberation mapping, estimate of the black hole mass, and finally constraints on the BLR structure. Finally, we summarize our main findings in Sect. 6.

## 2. Observations and analysis techniques

We compare emission line fluxes in images A & D of QSO 2237+0305. A comprehensive description of the observations and of the observational setup can be found in Papers I & II. The full data set consists of a series of spectra obtained at 39 different epochs between October 2004 and December 2007 with the FORS1 instrument at the ESO Very Large Telescope. All the observations were carried out in MOS (Multi Object Spectroscopy) mode. The multi-object mask used to obtain the data presented here included the quasar images A and D as well as several field stars which are used for simultaneous flux calibration and deconvolution. The spectra of the individual lensed quasar images were de-blended (from each other and from the lensing galaxy) using the MCS de-



**Fig. 1.** Comparison of V-band OGLE lightcurves of images A & D (points) of QSO 2237+0305 with the lightcurves derived from our spectra (diamonds; crosses are used for epochs deviating from the OGLE lightcurves). Error bars (not displayed) are typically of the order of 0.02 mag (A) and 0.04 mag (D).

convolution technique (Magain et al. 1998). A slit width of  $0.6''$  was used together with grism GRIS\_300V and blocking filter GG375, leading to a resolving power  $R = 400$  at  $5900 \text{ \AA}$ . For the data obtained in 2007, the FORS CCD had been replaced by a new camera more sensitive in the blue wavelength range. With this new camera, the GG375+30 blocking filter was not used. The spectral resolution was improved by about 10%, but the observations obtained with the new setup have a signal to noise ratio  $S/N \lesssim 10$  in the red wavelength range ( $\lambda > 6700 \text{ \AA}$ ) due to fringing. Table 1 provides a log of the data sample we used. Data flagged with a  $\dagger$  are systematically removed as the C IV emission is not covered by our spectra.

We show in Fig. 1 a comparison of the V-band lightcurves of images A & D provided by the OGLE collaboration (Udalski et al. 2006) with the lightcurves derived by applying V-band response curve to the spectro-photometric monitoring data. Throughout this paper, we shift the magnitude of image D from the OGLE-III lightcurve by  $-0.5 \text{ mag}$  w.r.t. the published values to account for different flux calibration of that image in the OGLE-III data (see Paper I for discussion). Although the overall agreement is very good, several epochs identified with an 'X' symbol deviate from the OGLE lightcurve. These data are flagged with an asterisk (\*) symbol in Table 1. We test how they affect our results in Sect. 5.2.

In order to study how microlensing deforms our spectra and how the broad lines are affected, we use three different techniques explained hereafter. Firstly, we apply the multi-component decomposition technique (MCD), i.e. we assume multiple components cause the quasar emission and measure the flux in these individual components (Sect. 2.1). Secondly, we use the macro-micro decomposition technique (MmD), i.e. we follow the prescription of Sluse et al. (2007) and Hutsemékers et al. (2010) to decompose the quasar spectrum

**Table 1.** Journal of the observations used in this paper. The observation ID refers to the whole data set presented in Papers I and II. Data flagged with a  $\dagger$  are not used. Data flagged with an \* are those deviating from the OGLE lightcurve in Fig. 1. As in Paper II, epochs #7, 8, 10, 30 have been excluded from the analysis.

	ID	Civil Date	JD-2450000	Seeing ["]	Airmass
Period P1	1*	13-10-2004	3292	0.86	1.204
	2	14-11-2004	3324	0.75	1.184
	3	01-12-2004	3341	0.88	1.355
	4*	15-12-2004	3355	0.99	1.712
	5	11-05-2005	3502	0.87	1.568
	6	01-06-2005	3523	0.63	1.342
	9	06-08-2005	3589	0.51	1.135
	11	25-08-2005	3608	0.49	1.261
	12	12-09-2005	3626	0.70	1.535
	13*	27-09-2005	3641	0.92	1.480
	14*	01-10-2005	3645	0.78	1.281
	15	11-10-2005	3655	0.57	1.140
	16*	21-10-2005	3665	0.70	1.215
	17*	11-11-2005	3686	0.90	1.137
	18*	24-11-2005	3699	0.78	1.265
	19*	06-12-2005	3711	1.10	1.720
Period P2	20 $\dagger$	24-05-2006	3880	0.87	1.709
	21 $\dagger$	16-06-2006	3903	0.66	1.213
	22 $\dagger$ *	20-06-2006	3907	0.64	1.286
	23 $\dagger$	27-06-2006	3914	0.41	1.145
	24	27-07-2006	3944	0.74	1.316
	25*	03-08-2006	3951	0.73	1.246
	26	13-10-2006	4022	0.59	1.176
	27	28-10-2006	4037	0.57	1.148
	28	10-11-2006	4050	0.89	1.515
	29	27-11-2006	4067	0.87	1.255
	31	22-12-2006	4092	0.80	2.018
Period P3	32*	10-07-2007	4292	0.63	1.153
	33*	15-07-2007	4297	0.57	1.158
	34	25-07-2007	4307	0.79	1.231
	35	03-08-2007	4316	0.68	1.412
	36	27-08-2007	4340	0.88	1.133
	37*	06-09-2007	4350	0.67	1.396
	38	20-09-2007	4364	0.73	1.230
	39*	23-09-2007	4367	1.23	1.530
	40	05-10-2007	4379	0.59	1.153
	41	10-10-2007	4384	0.76	1.283
	42	15-11-2007	4420	1.07	1.189
	43*	01-12-2007	4436	1.00	1.502

into a component  $F_M$ , unaffected by microlensing and another one,  $F_{M\mu}$ , which is microlensed (Sect. 2.2). Thirdly, we use the narrow band technique (NBD), i.e. we cut the emission lines into different velocity slices and measure the flux in each slice to look for microlensing within the emission line (Sect. 2.3).

### 2.1. The multi component decomposition (MCD)

Our most comprehensive analysis technique is a multi-component spectral decomposition of the quasar spectrum

(MCD). With this technique, we decompose the quasar spectra into several components likely associated with different emission regions (Dietrich et al. 2003). As in Papers I & II, the spectrum is modeled as a sum of three components: i) a power law for the quasar continuum emission; ii) an empirical iron template; iii) a sum of Gaussian profiles to model the emission lines. The exact decomposition of each spectrum differs in several respects from the one presented in Papers I & II:

- First, the quasar emission in image *D* is corrected for dust reddening produced by the lensing galaxy using the differential extinction values derived in Paper I. We adopt a Cardelli (1989) Milky-way like extinction law with  $(R_V, A_V) = (3.1, 0.2 \text{ mag})$ .
- Because the UV Fe II+III template constructed by Vestergaard & Wilkes (2001) from I Zw1 does not perfectly reproduce the Fe II+III blended emission observed in QSO 2237+0305 we construct the iron template empirically. In order to take into account possible intrinsic flux variations and/or systematic errors, we proceed independently for data obtained in 2004–2005 (period **P1**), 2006 (period **P2**) and 2007 (period **P3**). This template is constructed iteratively based on the residuals of the fitting procedure. In a first step, we only fit a power-law to the spectra, excluding the regions where Fe II+III is known to be significant (Vestergaard & Wilkes 2001) and where there are noticeable quasar broad emission lines. The subsequent fits are performed on the whole quasar spectrum (excluding atmospheric absorption windows), fixing the width and centroid of the quasar emission lines. A new fit is then performed using the residuals as the iron template. The procedure is repeated until a good fit of the whole quasar spectrum is obtained.
- In order to better reproduce the detailed shape of the C IV and C III] line profiles, we refine their modeling compared to Paper I by decomposing each line profile into a sum of three Gaussian profiles. The C IV line is composed of two emission (a narrow component NC and a broad component BC) and of one absorption component (AC). The faint 1600 Å blend of emission sometimes observed in AGNs, and caused mostly by He II λ1640, O III] λ1663 (Fine et al. 2010 and ref. therein), is included by construction in our Fe II+III template. To reproduce the C III] line we need to use three emission components of increasing width (abbreviated NC, BC and VBC). With the exception of C III](VBC), the central wavelength  $\lambda_c$  of the Gaussians are allowed to vary from epoch to epoch as well as the amplitude of each component. Table 2 provides for each image, the central wavelength  $\lambda_c$  and the FWHM of the different components of the line. The average value and standard deviation between epochs are displayed. When the standard deviation is not displayed, it means that the quantity is fixed for all epochs (see Appendix A for details). Figure 2 shows the C IV and C III] emission line on 2004–11–14 and the corresponding profile decomposition. As in Paper I, the other emission line profiles (e.g. Al III, He II, Si III], O III], Mg II) are fitted with single Gaussian profiles of fixed width.

**Table 2.** Decomposition of the C IV and C III] emission lines (Average value and scatter between epochs when the parameter is fitted). The central wavelength  $\lambda_c$  is expressed in Å and the FWHM in  $\text{km s}^{-1}$ .

ID	A		D	
	$\lambda_c$	FWHM	$\lambda_c$	FWHM
C IV				
BC	$1547.8 \pm 1.0$	$6332 \pm 404$	$1546.9 \pm 0.9$	$6128 \pm 557$
AC	$1548.9 \pm 0.5$	$1153 \pm 190$	$1548.7 \pm 0.4$	$1186 \pm 253$
NC	$1547.6 \pm 0.6$	2580	$1547.0 \pm 0.44$	2580
C III]				
BC	$1910.7 \pm 0.6$	3400	$1910.7 \pm 1.1$	3400
NC	$1907.3 \pm 0.4$	1545	$1907.5 \pm 0.3$	1545
VBC	1910.9	8548	1910.9	8548

The results of the MCD decomposition are shown and discussed in Sect. 3.1. One could suspect that residual flux from the lensing galaxy is included in our empirical FeII template. We tested this possibility by fitting a lens galaxy template (as retrieved from the deconvolution of the 2D spectrum, see Paper I) as an independent component. However, the quasar lightcurves derived in this case severely disagree with the OGLE lightcurves, indicating that residual contamination of the spectra by the lensing galaxy is not a severe issue. Finally, we emphasize that the decomposition of the carbon lines in multiple Gaussians of different widths is empirical and does not necessarily isolate emission lines coming from physically different regions. The adopted naming of the components is based on their width; it is line dependent and internal to this paper. Because of the different decompositions found for the C IV and C III] lines, comparison between gaussian components should be performed with care, keeping in mind the actual line widths.

## 2.2. Macro-micro decomposition (MmD)

This technique, introduced in Sluse et al. (2007) and Hutsemékers et al. (2010) and previously nicknamed  $F_M - F_{M\mu}$  decomposition, allows us to determine the fraction of the quasar emission which is not affected by microlensing independently of any modeling of the quasar spectra. Because spectral differences between lensed images are only due to microlensing (here we correct for differential extinction prior to the decomposition), we can assume that the observed spectrum  $F_i$  of image *i* is the superposition of an emission  $F_M$  which is only macro-lensed and of another component  $F_{M\mu}$  both macro and micro-lensed. Using pairs of spectra of different lensed images, it is then easy to extract both components  $F_M$  and  $F_{M\mu}$ . Defining  $M = M_A/M_D (> 0)$  as the macro-magnification ratio between image A and image D, and  $\mu = \mu_A/\mu_D$  as the microlensing ratio between image A and image D, we have:

$$\begin{aligned} F_A &= M \times F_M + M \times \mu \times F_{M\mu} \\ F_D &= F_M + F_{M\mu} \end{aligned} \quad (1)$$

The latter equations can be rewritten:

$$\begin{aligned} F_M &= \frac{F_A/M - \mu \times F_D}{1 - \mu} \\ F_{M\mu} &= \frac{F_D - F_A/M}{1 - \mu}, \end{aligned} \quad (2)$$

where  $\mu$  must be chosen to satisfy the positivity constraint  $F_M > 0$  and  $F_{M\mu} > 0$ . For the macro-magnification factor  $M$ , we adopt the ratio  $M = M_A/M_D = 1.0$  provided by the lens model of the Einstein cross performed by Schmidt et al. (1998). The microlensing ratio  $\mu(t, \lambda)$  is wavelength and time dependent and can be estimated for each epoch based on the ratio of the continuum emission between A & D (Sect. 2.1). Our choice of the continuum flux ratio to derive  $\mu(t, \lambda)$  means that we implicitly assume that the emission line and pseudo-continuum emission arise from regions equally or less microlensed than the continuum, in agreement with theoretical expectations. This assumption is also verified a posteriori as our analysis confirms that the continuum emission is also the most microlensed.

### 2.3. Narrow band technique (NBD)

A simple method to identify differential microlensing within an emission line is to divide the line into various wavelength bands and calculate the flux ratio  $F_A/F_D$  for each of these bands. For the C IV and C III] emission, we chose three different wavelength ranges corresponding respectively to the blue wing (BE;  $-8000 < v < -1500 \text{ km s}^{-1}$ ), the line core (CE;  $-1500 < v < 1500 \text{ km s}^{-1}$ ) and the red wing (RE;  $1500 < v < 8000 \text{ km s}^{-1}$ ) of the line. Before integrating the flux in these ranges, we first remove the continuum and the Fe II+III emission from the spectra based on the results of the MCD fit (Sect. 2.1). For the C III] emission we also remove the Al III and Si III] emission. The central wavelength corresponding to zero velocity is  $1549 \text{ \AA}$  for C IV and  $1907.5 \text{ \AA}$  for C III].

## 3. Microlensing variability in the emission lines

In Paper I, we reported evidence that the broad emission lines in QSO 2237+0305 are significantly microlensed in image A during the monitoring period. In this section, we refine and extend these results (including 2007 data which were absent from Paper I) using the techniques outlined in Sect. 2. This will allow us to derive lightcurves for the C IV and C III] emission lines and identify signatures of microlensing.


### 3.1. C IV and C III] emission lines

First, we investigate how microlensing affects the emission line shape. For this purpose we use the MmD technique (Sect. 2.2). The results of this decomposition for the C IV and C III] emission are shown in Fig. 3 for four representative epochs. On this figure, we also overplot as a dashed line the continuum model derived from the MCD decomposition and the spectrum of image D (dotted line). As indicated by Eq. 1, the spectrum of image D should match  $F_{M\mu}$  in the region of the spectra which are microlensed like the continuum ( $F_{M\mu} = F_D$  when  $F_M = 0$ ). The match between  $F_D$  and  $F_{M\mu}$  in the wings of the emission lines reveals that the corresponding emitting region is microlensed as


much as the continuum (Figure 3). It is also apparent that the microlensed fraction of the lines has a broad Gaussian shape with no flux depletion or flat core in the center (the depletion observed for the C IV emission is caused by the intrinsic absorber). On the other hand, in  $F_M$ , only a narrow emission line is visible for C IV and C III]. If we remove the continuum power law model and Fe II model prior to the decomposition, and plot the C III] and C IV lines on a velocity scale (Figure 4), we find that the decomposed line profiles are rather similar. The remainder of the difference could be caused by Fe II blended with the C III] emission, and therefore not included in our Fe II template, but may also be real. It is also apparent that the line profile is asymmetric with a stronger blue component, and that the C IV line is slightly blueshifted ( $\sim 500 \text{ km s}^{-1}$ ) compared to C III]. Although comparison between epochs is difficult due to intrinsic variability of the emission lines, we observe that the decomposition remains roughly the same during the period P1 (not shown). During periods P2 and P3, both fractions  $F_M$  and  $F_{M\mu}$  of the carbon lines follow the intrinsic brightening of the continuum. We also tentatively see that, during period P3, the line wings appear in  $F_M$ , indicative that the broadest component is differently microlensed than the continuum at these epochs. We note that the use of a macro-magnification ratio  $M \sim 0.85$ , closer to the radio and mid-infrared flux ratios (Falco et al. 1996, Agol et al. 2009, Minezaki et al. 2009), does not change the general features observed in Figure 3.

The slicing of the emission lines in velocity also allows us to study differential microlensing in the emission lines independently of a model of the line profile. Figure 5 displays the flux measured in the blue wing (BE), red wing (RE) and line core (CE) of the carbon emission lines, using the NBD (Sect. 2.3). Standard error bars are calculated based on the photon noise. Although the error bars are large, we clearly see a difference of about 0.3 mag between the flux ratios measured in the core and the wings of the emission lines. Table 3 shows the average flux ratios for periods P1, P2, P3. Error bars for a period are standard errors calculated based on the dispersion of the data points. The systematic difference observed between the blue and red wings of the C IV profile seems to be mostly caused by an asymmetry of the core of the line. Indeed, the offset between blue and red wing decreases as we increase the width of the central velocity range (not shown). The asymmetry of the  $F_M$  fraction of the line visible in Figure 4 supports this finding. When looking to the intrinsic variability of the different bands (i.e. individual lightcurves for images A & D), we observe that the wings vary more significantly on short time scales than the core of the line.

The results of the MCD decomposition are presented in Figure 6. They broadly agree with the velocity slicing technique (Fig. 5, Table 3). Figure 6 shows that the flux ratio in the full C III] profile (diamonds) varies weakly over the monitoring interval. A similar flux ratio is measured in C IV and C III] with evidence for a small increase ( $\Delta m_{AD}$  closer to 0) in 2007. Figure 6, also confirms that the flux ratio measured in the narrower component of the emission lines is closer to 1 (i.e. the macro-lensed ratio  $M_A/M_D$ ) while the flux ratio amounts to nearly 2.5 in the broadest component of the lines (i.e. BC of C IV and VBC of C III]).



**Fig. 2.** Example of multi component decomposition of the CIV (*left*) and C III] (*right*) line profiles (Sect. 2.1) after continuum and Fe II subtraction. The narrowest component of the line (NC) is displayed with a dotted-red line, the broad component (BC) with a dashed-blue line, the very broad component (VBC) of C III] and the absorber in front of CIV are shown in green dot-dashed. The sum of the individual components, corresponding to the line model, is shown with a solid gray line. Characteristics of the components are provided in Table 2.




**Fig. 3.** Macro-micro decomposition (MmD) of the CIV (*left*) and C III]+Si III] (*right*) emission lines obtained from the spectra of images A & D at 4 different epochs. In each panel, the black solid line shows the fraction of the spectrum  $F_{M\mu}$  affected by microlensing and the red solid line shows the emission  $F_M$  which is too large to be microlensed. For comparison, we also display the observed spectrum of image D with a dotted blue line and the power-law continuum used to calculate  $\mu(\lambda, t)$  with a dashed blue line.

### 3.2. Other emission lines


An accurate study of the microlensing occurring in the other emission lines is very difficult because of the pseudo-continuum Fe II+III in emission blended with these lines. The MmD technique gives qualitative information about microlensing for the other emission lines. Figure 7 shows the  $F_M$  and  $F_{M\mu}$

spectra averaged over period P1. We clearly see that not only CIV and C III] have their broadest component microlensed, but also Mg II  $\lambda 2798$ , Al II  $\lambda 1671$ , He II  $\lambda 1640$ , Al III  $\lambda 1857$ . The narrow component of these lines is clearly visible in  $F_M$ . Interestingly, the Si III]  $\lambda 1892$  is not microlensed. The absence of microlensed Si III] emission is also clearly visible in Fig. 3. The bottom panel shows the Fe<sub>UV</sub> template of Vestergaard &





**Fig. 4.** Macro-micro decomposition (MmD) technique applied to the C IV (black) and C III] (red) emission lines observed on 13-10-2006. *Left:* Fraction of the flux  $F_M$  non affected by microlensing, *Center:* Fraction of the line  $F_{M\mu}$  affected by microlensing, *Right:* Full emission line profile. The intensity of C III] are multiplied by 2.2 to ease visual comparison between C IV and C III] profiles. The vertical dashed line indicates the velocity zeropoint corresponding to the systemic redshift.




**Fig. 6.** Time variation of the magnitude difference between images A & D,  $\Delta m_{AD}$ , as measured in the continuum (*center*) and in the different components (Fig. 2) of the C IV (*left*), C III] (*right*) emission lines. The shaded area indicates the range of  $\Delta m_{AD}$  estimated from the macro-model and from MIR measurements. The acronyms NC, BC and VBC refer to the different components (of increasing width) of the C IV & C III] lines as shown in Fig. 2.

**Table 3.** Average values of magnitude difference  $\Delta m_{AD}$  as measured in the blue wing (BE), line core (CE), and red wing (RE) of the C IV and C III] emission lines.

Line	Period	$\Delta m_{AD}$ (mag)		
		BE	CE	RE
C IV	P1	$-0.91 \pm 0.02$	$-0.66 \pm 0.02$	$-1.27 \pm 0.02$
	P2	$-0.80 \pm 0.03$	$-0.66 \pm 0.02$	$-1.21 \pm 0.04$
	P3	$-0.66 \pm 0.02$	$-0.58 \pm 0.02$	$-1.10 \pm 0.02$
C III]	P1	$-1.08 \pm 0.02$	$-0.59 \pm 0.01$	$-1.07 \pm 0.02$
	P2	$-1.18 \pm 0.03$	$-0.62 \pm 0.01$	$-1.14 \pm 0.03$
	P3	$-1.32 \pm 0.05$	$-0.61 \pm 0.02$	$-1.13 \pm 0.02$

$\text{km s}^{-1}$ . Although the iron emission in QSO 2237+0305 differs slightly from the Vestergaard & Wilkes (2001) template (Paper I, Sect. 2.1), we clearly see that only the microlensed pseudo-continuum (i.e. emission above the continuum in  $F_{M\mu}$ ) resembles the  $\text{Fe}_{UV}$  emission but there is no sign of UV iron emission in  $F_M$ . The pseudo-continuum emission visible in  $F_M$  at  $\lambda > 1920 \text{ \AA}$  is unlikely to be produced by  $\text{Fe}_{UV}$  because its shape deviates significantly from expectations. Most likely origin(s) of this emission are UV emission from the host galaxy, Balmer continuum emission and scattered continuum light. Although this feature peaks at about  $2200 \text{ \AA}$  in the quasar rest frame, it is unlikely to be caused by differential extinction in the quasar host galaxy because the light-rays separation between A & D in the host is too small (Falco et al. 1999).






**Fig. 5.** Time dependence of the magnitude difference between images A & D,  $\Delta m_{AD}$ , as measured in the blue wing (BE; blue diamond), line core (CE; green square) and red wing (RE; red hexagons) of the C IV (*Top*) and C III] (*bottom*) emission lines. The shaded area indicates the range of  $\Delta m_{AD}$  estimated from the macro-model and from MIR measurements.

#### 4. Microlensing simulations

In order to derive the size of the regions emitting the carbon lines, we compare the observed microlensing signal with simulated microlensing lightcurves generated for different source sizes. The strategy we adopt is similar to the one explained in Paper II. Instead of searching for the best simulated lightcurve reproducing the data, we follow a Bayesian scheme for which a probability is associated to each simulated lightcurve. A probability distribution is then derived for each parameter that describes the lightcurve. We provide a description of the crucial steps in the next sub-sections.

##### 4.1. Creating the microlensing pattern

We use the state-of-the-art inverse ray-shooting code developed by Wambsganss (1990, 1999, 2001) to construct micro-magnification maps for images A & D. To create these patterns, the code follows a large number of light-rays (of order  $10^{10}$ ), from the observer to the source plane, through a field of



**Fig. 7.** *Top:* Macro-micro decomposition (MmD) applied to the spectra of images A & D averaged over period P1. The black solid line shows the fraction of the spectrum  $F_{M\mu}$  affected by microlensing and the red solid line shows the emission  $F_M$  which is too large to be microlensed. For comparison, we also display the observed spectrum of image D with a dotted-blue line and the power-law continuum of D with a dashed black line. *Bottom:* Fe II+III template from Vestergaard & Wilkes 2001 convolved with a Gaussian of FWHM=2000  $\text{km s}^{-1}$ .

randomly distributed stars. The surface density of stars and the shear  $\gamma$  at the location of the lensed images are those provided by a macro model of the system. Instead of the surface density, the dimensionless quantity  $\kappa$  (the convergence), defined as the ratio between the surface density and the critical density, is used. We use  $(\kappa, \gamma)_A = (0.394, 0.395)$  and  $(\kappa, \gamma)_D = (0.635, 0.623)$  (Kochanek 2004). Due to the location of the lensed images behind the bulge of the lens, we assume that 100% of the matter is in the form of compact objects. This assumption is motivated by the results of Kochanek (2004) who demonstrate that the most likely  $\kappa_{\text{smooth}} = 0$  in QSO 2237+0305. The mass function of microlenses has little effect on the simulations (Wambsganss 1992, Lewis & Irwin 1995, 1996, Congdon et al. 2007) and we therefore assume identical masses in the simulations. The mass of the microlenses sets the Einstein radius  $r_E$ . For QSO 2237+0305, the Einstein radius projects onto the source plane as:

$$r_E = \sqrt{\frac{4GM}{c^2} \frac{D_{os}D_{ls}}{D_{ol}}} = 9.84 \times 10^{16} \sqrt{\frac{M}{0.3M_{\odot}}} \text{cm}, \quad (3)$$


where the  $D$  are angular diameter distances, the indices  $o, l, s$  refer to observer, lens and source and  $M$  is the mass of microlenses. We create magnification patterns with sidelength of  $100 r_E$  and pixel sizes of  $0.01 r_E$ . Tracks drawn in such a pattern would provide simulated microlensing lightcurves for 1 pixel-size sources. In order to simulate lightcurves for other source sizes, we convolve the magnification pattern (after conversion on a linear flux scale) by the intensity profile of the source. As

shown by Mortonson et al. (2005), the exact shape of the source intensity profile has little influence on the lightcurve. It rather depends on the characteristic size of the source. For simplicity, we assume a Gaussian profile. A uniform-disc profile has also been tested for comparison. We construct magnification maps for 60 different source sizes having characteristic scales (Full Width at Half Maximum for the Gaussian and radius for the disc profile) in the range 2-1400 pixels ( $0.02 r_E$  -  $14 r_E$ ).

#### 4.2. Comparison with the data

The patterns created in Sect. 4.1 are used to extract simulated microlensing lightcurves to be compared to the data. Two data sets are used: the OGLE data of QSO 2237+0305 (Udalski et al. 2006), which provides well sampled V-band lightcurves of the continuum variations, and our spectrophotometric lightcurves presented in Sect. 2. In order to study the microlensing signal, we have to get rid of intrinsic variations. This is done by calculating the difference lightcurves between images A & D. Similarly, the simulated microlensing signal is obtained by taking the differences between the simulated tracks for images A & D. The parameters which characterise a simulated microlensing track are the starting coordinates in the magnification patterns ( $\mathbf{x}_{0,A}$ ,  $\mathbf{x}_{0,D}$ ), the track position angle ( $\theta_A$ ,  $\theta_D$ ), and the track length. As the extracted track is compared to a lightcurve obtained over a given time range, the track length corresponds to the transverse velocity of the source expressed in  $r_E$  per Julian day. As in Paper II, we assume that the velocity is the same for the trajectories in A & D and the track orientation  $\theta_A = \theta_D$ . To account for the fact that the shears in A & D are roughly orthogonal (Witt & Mao 1994), we rotate the magnification map of D by  $90^\circ$  prior to the track extraction. The difference of the magnitude between track A and track D should, on average, be equal to the one of the macro-model. Like other authors, we account for the uncertainty on the amount of differential extinction between A & D and on the macro-model flux ratio by allowing for a magnitude offset  $m_0$  between the simulated lightcurves extracted for A and D. The other parameters characterizing the lightcurves are the convergence and shear ( $\kappa, \gamma$ ) fixed in Sec. 4.1. Physical quantities (source size and velocity) are proportional to the mass of microlenses as  $M^{1/2}$ .

In order to build a representative ensemble of lightcurves which is compatible with the data, we follow a four-step strategy similar to the one described in Paper II, in Anguita et al. (2008) and in Kochanek (2004): 1) we pick a set of starting values for the parameters defining the tracks in A & D and vary them to minimize the  $\chi^2$  between the simulated lightcurve and the OGLE lightcurve; 2) we repeat step 1 in order to get a representative ensemble of 10000 tracks and a good coverage of the parameter space; 3) each track estimated in step 2 is computed for other source sizes and the agreement with the BLR lightcurve is quantified using a  $\chi^2$ -type merit function; 4) the  $\chi^2$  estimated in steps (2) and (3) are summed up and used to calculate a likelihood distribution for any track parameter. A summary of the technical details regarding microlensing simulations is given in Appendix B. An example of a good simulated lightcurve fitting the data is shown in Figure 8. In the next sub-



**Fig. 8.** *Top:* Example of a good track fitting the differential lightcurve  $\Delta m_{AD}$  observed for the CIV broad line and for the continuum. *Center:* Corresponding tracks (solid line) in the microlensing pattern of image A (left) and D (right). The large circle represents the CIV source size and the small one the continuum. *Bottom:* Track corresponding to the continuum emission (black solid line) and to the CIV emitting region (dashed gray line) for image A (left) and D (right).

section, we explain how we derive probability distributions for the quantities of interest.

#### 4.3. Probability distribution

The probability of a trajectory  $j$ , defined by the set of parameters  $p = (\kappa, \gamma, M, R_s^c, R_s^l, V, m_0, \mathbf{x}_{0,A}, \mathbf{x}_{0,D})$  (where we drop the index  $j$  for legibility,  $R_s^c$  and  $R_s^l$  refer to the source size of the continuum and of the line) given the data  $D$  is, according to Bayes theorem:

$$P(p_j|D) = \frac{L(D|p_j)P(p_j)}{\sum_{j=1}^n L(D|p_j)P(p_j)} = \frac{L(D|p_j)P(R_{s,j}^c)P(V_j)}{\sum_{j=1}^n L(D|p_j)P(R_{s,j}^c)P(V_j)} \quad (4)$$

where  $n$  is the total number of trajectories in the library,  $L(D|p_j)$  is a likelihood estimator of the data  $D$  given the parameters  $p_j$

and  $P(R_{s,j}^c)$  and  $P(V_j)$  are the priors on the continuum source size and velocity respectively. Only the prior on  $R_s^c$  and on  $V$  appear in Eq. 4 because we use an uniform prior on the other (nuisance) parameters. The prior  $P(R_{s,j}^c)$  is required because we have a non uniform sampling of  $R_s^c$ . The prior is related to the density of trajectories as a function of  $R_{s,j}^c$  given our source sampling in 60 different bins. If  $R_{s,j}^c$  falls in the bin  $b$  of width  $l_b$ , then we have  $P(R_{s,j}^c) \propto l_b$ . The velocity prior  $P(V_j)$  is identical to the one used in Anguita et al. (2008). It is based on previous studies by Gil-Merino et al. (2005) which suggest that the net transverse velocity is smaller than  $0.001 r_E/\text{jd}$  (at 90% confidence level). This corresponds to  $v < 660 \text{ km s}^{-1}$  in the lens plane ( $\sim 6000 \text{ km s}^{-1}$  in the source plane) for  $M \sim 0.1 M_\odot$ , in agreement with the velocity prior used by Kochanek (2004) and Poindexter & Kochanek (2010a).

As explained in Kochanek (2004), the use of the standard likelihood estimator  $L(D|p_j) = \exp(-\chi^2(p_j)/2)$  can lead to misleading results because it is very sensitive to the accuracy of the error bars and to change of the  $\chi^2$  induced by possible outliers. To circumvent this problem, we follow Kochanek (2004) and use the following likelihood estimator:

$$L(D|p_j) = \Gamma \left[ \frac{n_{\text{dof}} - 2}{2}, \frac{\chi^2(p_j)}{2} \right], \quad (5)$$

where  $\Gamma$  is the incomplete gamma function, and  $\chi^2 = \chi_{\text{OGLE}}^2 + \chi_{\text{line}}^2$  is the sum of the  $\chi^2$  obtained when fitting the OGLE data and emission line lightcurves (steps 2 & 3 in Sect. 4.2).

## 5. Inferences of the quasar structure

In this section, we present the results of our microlensing simulations and discuss them in the light of the phenomenological study of the emission lines in A & D. First, we discuss the domain of validity of our simulations and the meaning of the BLR size estimated with our method. Second, we derive the size of the continuum and of the BLR for C IV and C III]. Third, we compare our results to the Radius(BLR)-Luminosity relation derived from reverberation mapping. Fourth, we derive the black hole mass of QSO 2237+0305 using the virial theorem and compare it to black hole mass derived from the accretion disc size. Finally, we discuss the consequences of our analysis for the BLR structure of QSO 2237+0305.

### 5.1. Caveat

In microlensing simulations, we represent the projected BLR with a face-on disc having a Gaussian or a uniform intensity. The exact intensity profile should not affect our estimate of the size (Mortonson et al. 2005) but the sizes retrieved are robust only if the BLR has actually a morphology which is disc-like when projected on the plane of the sky. This corresponds to the case of a spherically symmetric BLR geometry or of an axisymmetric geometry with the axis pointing roughly towards us. Since the microlensing signal remains nearly unchanged for sources with ellipticity  $e = 1 - q < 0.3$  (Fluke & Webster 1999, Congdon et al. 2007), our simulations should also be valid for disc-like sources inclined by up to  $\phi \sim 45^\circ$  w.r.t. the

line of sight. This means that inclination may affect our size estimates by a factor  $\cos \phi > 1/\sqrt{2}$ . Although the exact geometry of the BLR is unknown, there are several line of evidence that QSO 2237+0305 is seen close to face-on. Firstly, QSO 2237+0305 is a Type 1 AGN with a flat radio spectrum ( $\alpha = -0.18$ , Falco et al. 1996), indicative that the source is seen nearly face-on. Secondly, the analysis of the microlens-induced continuum flux variations by Poindexter & Kochanek (2010b) show that the inclination of the accretion disc is  $< 45^\circ$  at 68.3% confidence level. Thirdly, the C IV line shows only a small blueshift which can be interpreted as an evidence that the BLR is seen face-on (Richards et al. 2002).

If the BLR has a disc-like morphology, the projected emission should take place in a ring-like<sup>1</sup> region (with an inner radius  $R_{\text{in}}$  and outer radius  $R_{\text{out}}$ ), rather than in a uniform disc as assumed here. The effect of such a 'hole' in the BLR emission has been studied by Fluke & Webster (1999) who showed that the microlensing lightcurve would be significantly affected only for  $R_{\text{in}} > 0.5 R_{\text{out}}$ . Since the BLR is commonly assumed to have  $R_{\text{out}} > 10 R_{\text{in}}$  (e.g. Murray & Chiang 1995, Collin & Húr 2001, Borguet & Hutsemékers 2010), we shall not consider this geometrical issue.

### 5.2. Absolute and relative size of the continuum and the BLR

The V-band OGLE lightcurve is associated with emission arising mostly from the quasar continuum. The probability distribution of the continuum source size  $R_s^c$  derived from our microlensing study of that lightcurve (Sect. 4) is displayed in Fig. 9. On this figure, we also show, on a color-coded scale, the average velocity of the tracks in each source size bin. We see that the source size and the velocity are strongly correlated as expected from the source size/velocity degeneracy (e.g. Kochanek 2004, Paper II). When we consider a uniform prior on the velocity (not shown), we find that imposing  $m_0$ , the average difference of magnitude between A & D lightcurves, to be equal to the macro-model expectation allows us to exclude very high velocities ( $v > 0.005 r_E/\text{jd}$ ). For the velocity range allowed by the Gil-Merino et al. (2005) velocity prior, the same distributions of  $m_0$  is found whatever the velocity bin considered. The median half light radius of the continuum we derive from our simulations is  $R_s^c = 8.8 \times 10^{15} \text{ cm}$  ( $2.5 \times 10^{15} \text{ cm} < R_s^c < 25.4 \times 10^{15} \text{ cm}$  at 68.3% confidence level,  $M = 0.3 M_\odot$ ) for a Gaussian profile, and  $R_s^c = 12.3 \times 10^{15} \text{ cm}$  ( $3.2 \times 10^{15} \text{ cm} < R_s^c < 36.8 \times 10^{15} \text{ cm}$  at 68.3% confidence level,  $M = 0.3 M_\odot$ ) for a uniform disc. Table 4 compares our measurements with the results of Kochanek (2004, KOC04), Anguita et al. (2008), Paper II and Poindexter & Kochanek (2010b). The half-light radius ( $R_{1/2}$ ) allows the comparison of sizes derived with various intensity profiles. We see that our results are compatible with all other works.

The probability distributions of the half light radius measured for the C IV and C III] emission lines (using a Gaussian

<sup>1</sup> In the case of emission coming from a spherical shell, the projected profile may also look ring-like but emission should also arise from the central part of the disc.


**Table 4.** Comparison of V-band continuum source size of QSO 2237+0305 (Sect. 5.2). The reference is presented in Col. #1, the source profile used in Col. #2, and the published size estimate in Col. #4 (where  $R$ = radius of a uniform disc,  $R_V$ = scale radius of a Shakura-Sunyaev accretion disc,  $\sigma_V$  and FWHM are resp. the standard deviation and the full width at half maximum of a Gaussian). These quantities have been converted to half light radius  $R_{1/2}$  to allow comparison between profiles. Col. #6 & #7 provide the average mass of microlenses ( $\langle M \rangle$ ) used in the microlensing simulations and information regarding the velocity prior.

Reference	Profile	Estimator	Size ( $10^{15}$ cm)	$R_{1/2}$ ( $10^{15}$ cm)	$\langle M \rangle$	Velocity prior
This paper	Gauss	Median	$17.6^{+33.3}_{-12.5}$ (FWHM)	$8.8^{+16.6}_{-6.3}$	$\sqrt{M/0.3M_\odot}$	Gil-Merino 2005
This paper	Uniform disc	Median	$17.4^{+34.6}_{-12.9}$ ( $R$ )	$12.3^{+24.5}_{-9.1}$	$\sqrt{M/0.3M_\odot}$	Gil-Merino 2005
Poindexter et al. (2010b)	Shakura-Sunyaev	Median	$5.8^{+3.8}_{-2.3}$ ( $R_V$ )	$14.1^{9.2}_{-5.6}$	$0.3 M_\odot$	Poindexter et al. (2010a)
Anguita et al. (2008)	Gauss	Most likely	$4.6^{+3.4}_{-3.4}$ ( $\sigma_V$ )	$5.4^{+4.0}_{-4.0}$	$\sqrt{M/0.1M_\odot}$	Uniform
Anguita et al. (2008)	Gauss	Most likely	$1.3^{+0.2}_{-0.7}$ ( $\sigma_V$ )	$1.5^{+0.3}_{-0.8}$	$\sqrt{M/0.1M_\odot}$	Gil-Merino 2005
Paper II	Gauss	Most likely	$40^{+75}_{-35}$ (FWHM)	$20^{+37.5}_{-17.5}$	$\sqrt{M/0.1M_\odot}$	Uniform
Paper II	Gauss	Most likely	$9.2^{+6.9}_{-5.8}$ (FWHM)	$4.6^{+3.4}_{-2.9}$	$0.1 M_\odot$	Similar to Kochanek (2004)
Kochanek (2004)	Shakura-Sunyaev	Median	$4.1^{+6.7}_{-1.8}$ ( $R_V$ )	$10^{+16.3}_{-4.4}$	$[0.1, 1]M_\odot$	Sect. 2.3 of Kochanek (2004)
Kochanek (2004)	Gauss	Median	$5.1^{+4.7}_{-2.8}$ ( $\sigma_V$ )	$5.9^{+5.5}_{-3.2}$	$[0.1, 1]M_\odot$	Sect. 2.3 of Kochanek (2004)

source light profile) are displayed in Figure 10. Reasonably good fits are obtained for all the line components. However the quality of the best fits tends to be poorer for the broadest line component (typical reduced  $\chi^2 \sim 2$ ), the poorer fits being obtained for the very broad component of the C III] line which has a reduced  $\chi^2 \sim 4$ . This suggests that our error bars on the flux ratio are underestimated (by a factor  $\sim \sqrt{2}$ ) when measured in the broadest line components. This is not unexpected as the flux in these components of the emission lines are more prone to measurement errors in the MCD decomposition. Our results are however robust in the sense that they remain unchanged when the error bars are increased by this factor. Median half light radius and 68.3% confidence intervals are provided in Table 5 for both Gaussian and uniform disc intensity profiles of the BLR. These sizes are independent of the intensity profile used for the continuum emission. These results confirm that the most compact region of the quasar is also the one emitting the broadest components of the lines.

While the absolute source sizes derived above depend on the mass of the microlenses, the relative size of the continuum and the BLR are independent of the mass of microlenses. Table 6 displays the ratio between the emission line and the continuum half-light radius size  $R_s^l/R_s^c$  (assuming the same source profile for the continuum and BLR), at 68.3% confidence, for the different components of the BLR identified in Sect. 2.1. We can see that the region emitting the broadest components of the C IV and of the C III] lines (i.e. C IV<sub>BC</sub> and C III]<sub>VC</sub>) are about 4 times larger than the region emitting the V-band continuum. In contrast, the narrowest component of the emission lines arises from a region typically 25 times larger than the continuum.

We shall notice that the radius of the Extreme-UV continuum ionizing the carbon lines (typically  $\lambda < 500 \text{ \AA}$ ) is expected to be much smaller than the V-band continuum, with a dependence of the size with wavelength as  $R \propto \lambda^\xi$  (Paper II and ref. therein). For a standard accretion disc model,  $\xi = 4/3$ , implying that the UV ionizing continuum is more than 7 times smaller than the OGLE-V band continuum.




**Fig. 9.** Probability distribution for the half-light radius of continuum emission in QSO 2237+0305, color-coded as a function of the average (lens plane) velocity of the tracks in a given bin.

Finally, we note that the results of Table 5 & 6 are not sensitive to possible outliers in our BLR lightcurves. We retrieve essentially the same results when the points inconsistent with the OGLE lightcurve (i.e. those which are marked with a 'x' in Fig. 1) are excluded. Similar results, but affected by larger error bars, are also obtained if we do not consider the 2007 data.

### 5.3. C IV Radius-Luminosity relationship

In this subsection, we compare our microlensing-based estimate of the C IV emission line (Sect. 5.2) with the Radius(BLR)-Luminosity relationship (hereafter  $R_{BLR} - L$ ) derived with the reverberation mapping technique (Sect. 1.2). The  $R_{BLR} - L$  relationship for C IV has been obtained only for six objects (Peterson et al. 2005, Kaspi et al. 2007). The luminosity used in this relationship is commonly  $L_\lambda(1350 \text{ \AA})$ . Unfortunately, the latter is out of our spectral range. Instead



**Fig. 10.** Probability distribution for the size of the CIV (*top*) and CIII] (*bottom*) broad emission lines. Measurements for the region emitting the CIV & CIII] full emission line profiles (thick black lines, TOT) and for their components NC, BC and VBC (in order of increasing FWHM, cf. Fig. 2), are presented.

we use  $\lambda L_{\lambda}(1450\text{\AA}) = 10^{45.53} \text{ erg/s}$  (Assef et al. 2010) which has been shown to be equivalent to  $L_{\lambda}(1350\text{\AA})$  (Fig. 4 of Vestergaard & Peterson 2006). This luminosity estimate, directly measured on our spectra, is corrected for macro-magnification and line-of-sight Galactic extinction. The macro-magnification is correct within a factor 2 (Witt & Mao 1994). Because the lensed images are located behind the bulge of the galaxy, it is reasonable to assume that the absolute extinction due to the lens is  $< 1.0$  mag. The finite slit width of  $0.6''$  and seeing  $\sim 0.6''$  may lead to an underestimate of the total flux by a factor up to 2. This leads to an uncertainty factor of 4 on  $\lambda L_{\lambda}(1450\text{\AA})$ . The CIV size has been derived to be  $R_s^{\text{CIV}} = 1.7^{+2.9}_{-1.2} \times 10^{17} \sqrt{M/0.3M_{\odot}} \text{ cm} = 65.6^{+110}_{-46} \sqrt{M/0.3M_{\odot}} \text{ light-days}$  (Tab. 5). Figure 11 shows the  $R_{\text{BLR}}-L$  diagram including our measurement for QSO 2237+0305 and the analytical relation derived by Kaspi et al. (2007) for their so called 'FITEXY' fit. Although the reverberation-mapping size may differ from the half-light radius by a factor up to a few depending on the

**Table 5.** Half-light radius of the line emitting region  $R_s^l$  (in  $10^{15} \sqrt{M/0.3M_{\odot}} \text{ cm}$ ) for a Gaussian (G) and a uniform disc (D) intensity profile. Measurements for the CIV & CIII] full emission line profiles (TOT) and for their individual components NC, BC and VBC (in order of increasing FWHM, cf Fig. 2), are presented.

Line	$R_s^l$ (G)	68.3% CI	$R_s^l$ (D)	68.3% CI
CIV BC	42.9	[14.1,124.1]	56.6	[17.6,168.6]
CIV NC	$> 82.7$	-	$> 103.5$	-
CIV TOT	170.7	[47.7,457.6]	276.6	[72.0,676.0]
CIII] BC	150.1	[51.7,430.5]	209.6	[69.0,627.7]
CIII] NC	$> 97.9$	-	$> 112.2$	-
CIII] VBC	53.6	[21.2,126.9]	41.7	[17.4,103.0]
CIII] TOT	126.4	[36.9,399.5]	163.5	[43.8,570.6]

**Table 6.** Ratio of size  $R_s^l/R_s^c$  of the BLR to continuum region for Gaussian (G) and uniform disc (D) source profiles. Measurements for the region emitting the CIV & CIII] full line profiles and for their components NC, BC and VBC (in order of increasing FWHM), as shown in Fig. 2, are presented.

Line	$R_s^l/R_s^c$ (G)	68.3% CI	$R_s^l/R_s^c$ (D)	68.3% CI
CIV BC	3.9	[1.6,13.4]	3.0	[1.1,12.0]
CIV NC	27.7	[9.0,89.3]	22.2	[7.1,73.8]
CIV TOT	18.9	[4.6,72.3]	17.5	[4.5,61.2]
CIII] BC	16.9	[4.9,63.1]	15.2	[4.0,58.9]
CIII] NC	29.4	[10.5,103.6]	25.6	[8.1,82.5]
CIII] VBC	5.0	[2.4,14.3]	2.3	[0.9,7.9]
CIII] TOT	13.8	[0.7,53.8]	12.6	[3.4,49.4]

geometry and density of the BLR gas, we find a remarkably good agreement with the  $R-L^{\alpha}$  relationship.

#### 5.4. Black hole mass estimates


The BLR size we derive from microlensing can be used to estimate a black hole mass using the virial theorem:

$$M_{\text{BH}} = \frac{f}{G} R_{\text{BLR}} \text{FWHM}_{\text{BLR}}^2, \quad (6)$$

where  $f=1$  is a geometric factor, which encodes assumptions regarding the BLR geometry (e.g. Vestergaard & Peterson 2006, Collin et al. 2006), calibrated from the results of Onken<sup>2</sup> et al. (2004). Using  $\text{FWHM}_{\text{CIV}} = 3960 \pm 180 \text{ km s}^{-1}$  (Assef et al. 2010), we derive  $M_{\text{BH}} = 2.0^{3.4}_{-1.5} \times 10^8 M_{\odot}$ . Consistent measurements, but generally affected by larger error bars, are derived using the other broad components of the emission lines. Hereafter we refer to this black hole mass estimate as  $M_{\text{BH}}^{\text{BLR}}$ . As we argued earlier, inclination w.r.t. the line of sight may reduce  $R_{\text{BLR}}$  by  $\cos \phi > 1/\sqrt{2}$ .

<sup>2</sup> Onken et al. derive  $f = 5.5$  but using  $\sigma^2$  instead of  $\text{FWHM}^2$  in the virial relation. The corresponding value of  $f$  when using the FWHM is  $f = 1$ .





**Fig. 11.**  $R_{\text{CIV}}-L$  diagram combining reverberation mapping measurement of  $R_{\text{CIV}}$  ('x' symbol, Peterson et al. 2005, Kaspi et al. 2007) and our microlensing based size for QSO 2237+0305. The solid line is the analytical fit to the reverberation mapping data from Kaspi et al. (2007), which has a slope  $\alpha = 0.52$ .

Our estimate of  $M_{\text{BH}}^{\text{BLR}}$  may be uncertain by a factor of a few due the variation of  $f$  from object to object. A value of  $f$  larger than our fiducial value  $f = 1.0$  may be expected for axi-symmetric geometry of the BLR seen nearly face-on. The study of several plausible theoretical models of the BLR allowed Collin et al. (2006) to show that inclination of axi-symmetric BLR models may noticeably increase  $f$ . Observationally, Decarli<sup>3</sup> et al. (2008) found  $f$  in the range  $[0.5, 10]$ , with larger values ( $f > 4$ ) only for objects with  $\text{FWHM} < 4000 \text{ km s}^{-1}$ .

The use of the virial theorem to derive  $M_{\text{BH}}$  based on the CIV line is still debated. For this line, non gravitational effects, such as obscuration and radiation pressure, are probably affecting the CIV profile (Baskin & Laor 2005, Marziani et al. 2006). Obscuration, if present, should lead to an underestimate of  $\text{FWHM}(\text{CIV})$  and therefore to an overestimate of the  $M_{\text{BH}}^{\text{BLR}}$ . The effect of radiation pressure might be more subtle. While Marconi et al. (2008) have argued that radiation pressure implies an underestimate of  $M_{\text{BH}}$ , recent calculations of Netzer & Marziani (2010) show that orbits of BLR clouds are affected in such a way by radiation pressure that the product  $R_{\text{BLR}} \text{FWHM}^2$  (and therefore  $M_{\text{BH}}$ ) will vary by at most 25% due to radiation pressure. Our finding of roughly similar  $M_{\text{BH}}$  for different components of the line, emitted in regions differently affected by radiation pressure, further supports the idea that  $M_{\text{BH}}$  in QSO 2237+0305 is not strongly biased by non gravitational effects.

A black hole mass estimate can also be derived from the size of the accretion disc. In the accretion disc theory, the radius  $R_\lambda$  at which the disc temperature equals the photon energy,

$kT = hc/\lambda_{\text{rest}}$  is given by the following expression (Poindexter & Kochanek 2010b):

$$R_\lambda = \left( \frac{45 G \lambda_{\text{rest}}^4 M_{\text{BH}} \dot{M}}{16 \pi^6 h c^2} \right)^{1/3} \quad (7)$$

where  $M_{\text{BH}}$  is the black hole mass and  $\dot{M}$  is the accretion rate. Rewriting this expression for the black hole mass, we get:

$$M_{\text{BH}}^{\text{disc}} = 0.68 \times \left( \frac{R_{1/2}}{10^{15} \text{ cm}} \right)^{3/2} \left( \frac{\lambda_{\text{rest}}}{0.2 \mu\text{m}} \right)^{-2} \eta_{0.1}^{1/2} \left( \frac{L}{L_E} \right)^{-1/2} 10^8 M_\odot \quad (8)$$

where  $0.1 \eta_{0.1} = \eta = L/(\dot{M} c^2)$  is the radiative efficiency of the accretion disc,  $R_{1/2} = 2.44 R_\lambda$  is the half light radius of a Shakura-Sunyaev accretion disc model and  $L/L_E$  is the luminosity in units of Eddington luminosity. Using the accretion disc half light radius  $R_s = 8.8 \times 10^{15} \text{ cm}$  (Sect. 5.2), we get  $M_{\text{BH}}^{\text{disc}} = 17.7 \times 10^8 M_\odot$  ( $2.7 \times 10^8 M_\odot < M_{\text{BH}}^{\text{disc}} < 8.7 \times 10^9 M_\odot$  at 68.3% confidence) if we assume that  $L = L_E$  and  $\eta = 0.1$ . Unfortunately,  $M_{\text{BH}}^{\text{disc}}$  is also prone to systematic errors and depends strongly on the exact temperature profile of the disc. The black hole mass derived with this method assumes a temperature profile  $T \propto R^{-\xi'}$  with  $\xi' = 3/4$  (Shakura & Sunyaev 1973). A different value of  $\xi'$  in the range  $[0.5, 1]$  may lead to  $M_{\text{BH}}^{\text{disc}}$  larger/smaller by a factor as large as  $\sim 5$  (e.g. Poindexter & Kochanek 2010b). The  $M_{\text{BH}}^{\text{disc}}$  is thus more uncertain and less accurate than  $M_{\text{BH}}^{\text{BLR}}$ .

By comparing the isotropic luminosity  $L = 4.0 \times 10^{46} \text{ erg/s}$  of QSO 2237+0305 (Agol et al. 2009) to the Eddington Luminosity  $L_E = 1.38 \times 10^{38} M_{\text{BH}}/M_\odot \text{ erg/s}$ , we derive a black hole mass  $M_{\text{BH}}^E > 2.9 \times 10^8 M_\odot$ , consistent with  $M_{\text{BH}}^{\text{BLR}}$  and  $M_{\text{BH}}^{\text{disc}}$ .

### 5.5. The structure of the BLR

Based on the MCD decomposition of the spectra, we have shown that the CIV and CIII] lines can be decomposed into 2 (resp. 3) components of increasing width (Fig. 2). The possibility that these line components correspond to physically distinct regions in the BLR has been discussed by several authors (e.g. Brotherton et al. 1994b, Marziani et al. 2010) but remains debated. The variable amount of microlensing in these components allows us to shed new light on this scenario.

The differential microlensing lightcurves observed for the Gaussian components of the emission line (Fig. 6) are essentially flat, even when a substantial microlensing signal is observed in the continuum. They also show increasing microlensing effect in the broadest line components. Our simulations show that it is possible to reproduce both the continuum and BLR lightcurves by changing only the source size. We clearly find that the most compact regions have the broadest emission. This excludes BLR models for which the width of the emission line is proportional to the size of the emitting region. Our results are compatible with a variation of the BLR size with  $\text{FWHM}^{-2}$  as expected if the emitting regions are virialized. This arise in various geometries, i.e., for a spherically isotropic BLR, a Keplerian disc, or a disc+wind model similar to one proposed by Murray & Chiang (1995, 1997).

<sup>3</sup> Decarli et al. use a different definition of  $f$ :  $f_{\text{thispaper}} = f_{\text{Decarli}}^2$ .

The other techniques we use to analyse the spectra provide additional evidence that the BLR of C IV and C III] is composed of several regions. The "narrow-band" technique (Sect. 2.3) reveals that  $\Delta m_{AD}$  is closer to the macro model value (i.e.,  $\Delta m_{AD} \sim -0.1 \pm 0.1$ ) in the core of the emission lines than in the wings. But even in the central  $200 \text{ km s}^{-1}$  part of the line, we find  $\Delta m_{AD} \sim -0.6 \text{ mag}$ , indicating that there is microlensed flux in the core of the line. The MmD technique (Sect. 2.2) provides further clues on the origin of this flux. The shape of  $F_{M\mu}$  for the carbon lines, smoothly increases from the base to the line center, and is compatible with a single-peak profile. The lack of "accretion disc-like" profile with a flatter or depleted core (see e.g. Bon et al. 2009) does not exclude a disc like geometry of the BLR because disc emission can also produce single peaked profiles (Robinson 1995, Murray & Chiang 1997, Down & al. 2010). The MmD supports the use of a Gaussian to empirically decompose the emission lines with the MCD technique and a composite model of the BLR formed by at least 2 different regions.

Clues to the physical conditions occurring in these regions should come from the flux ratios measured between lines. A detailed analysis, however, needs very high signal to noise measurements (based e.g. on an average spectrum of QSO 2237+0305), proper account of Galactic reddening and detailed photo-ionisation models. This is beyond the scope of this paper and is devoted to a future publication. We shall however mention several features already visible in our spectrum and which should provide interesting constraints on photo-ionisation models. First, there is no broad Si III] emission. The ratio Si III]/C III] is very sensitive to the electronic density (Keenan et al. 1987, Aoki & Yoshida 1999). The absence of Si III] in the broad component of the emission indicates that the electronic density  $n_e$  in the region emitting the broad component of C III] is smaller than in the region emitting the narrow portion of the lines. Second, there is no clear narrow Fe II+III emission, suggesting that the latter arises from the inner part of the BLR or from a very compact region. This contrasts with microlensing observed in the lensed quasar RXJ 1131-1231 (Sluse et al. 2007) where narrow and broad emission were found for the 'optical' Fe II+III emission ( $\lambda \in [3000, 5000] \text{ \AA}$ ), the UV Fe II being poorly covered by the spectra in that system. Third, the ratio C III]/C IV is found to be nearly the same in the broad and narrow component of the line (Fig. 4). This advocates for a similar ionisation parameter  $U$  in these two regions (Mushotzky & Ferland 1984), which is *a priori* surprising since  $U$  should vary with the distance. In their decomposition of the emission lines based on the principal component analysis technique, Brotherton et al. (1994b) faced a similar problem, and found that C III]/C IV measured for the narrow part of the line (their so called intermediate line region) was too large by an order of magnitude compared to photo-ionisation models. We speculate that metallicity and optically thick/thin nature of the BLR gas are likely to play an important role in reproducing these ratios (Snedden & Gaskell 1999).

Small differences between the C IV and C III] line profile are visible in the total line profile (Fig. 4): the C III] profile peaks nearly at the systemic redshift and is rather symmetric while the C IV line shows a small blueshift and an excess of

emission in the blue (i.e. blue asymmetry), in agreement with previous studies (Brotherton et al. 1994a). Figure 4 shows that these differences are visible in the narrow component of the line. On the other hand, the very broad component of these lines is slightly asymmetric for both lines, showing again a blue excess but a more significant one for C IV. Because the intrinsic absorber is superimposed on the C IV profile, it is hard to assess whether the broadest component of the profile is in fact blueshifted like the narrow one. Finally, we mention that the core of the carbon lines varies less than the base of the line, in agreement with the finding of Wilhite (2006) based on a large sample of SDSS quasars. This finding is consistent with the above arguments showing that the broadest part of the line arises from a region closer to the continuum.

## 6. Conclusions

For the first time we have derived the size of the broad line region in QSO 2237+0305 based on spectrophotometric monitoring data, consisting of 39 different epochs obtained between Oct. 2004 and Dec. 2007. To reach this goal, we have measured differential lightcurves between images A & D for the C IV and C III] broad emission lines and compared them to microlensing simulations. This leads to the determination of the half-light radius of the C IV emitting region:  $R_{CIV} \sim 66 \text{ lt-days}$  ( $19 \text{ lt-days} < R_{CIV} < 176 \text{ lt-days}$  at 68.3% confidence), in very good agreement with the  $R_{BLR} - L$  relation derived from reverberation mapping (Kaspi et al. 2007). The size we derive for C III] is  $R_{CIII] \sim 49 \text{ lt-days}$  ( $14 \text{ lt-days} < R_{CIII] < 154 \text{ lt-days}$  at 68.3% confidence), compatible with the size estimated for the region emitting C IV.

Thanks to the variable amount of microlensing observed within a given emission line, we can also derive information on the structure of the broad line region. Differential lightcurves obtained for various velocity slices of the C IV and C III] lines show that the wings of the lines are more microlensed than the core, indicating that the former arise in a more compact region. This finding is confirmed by two other techniques. The first one demonstrates that a broad and single-peaked fraction of the emission line is microlensed, while a narrower fraction is unaffected by microlensing. This technique also suggests a slightly different structure for the C IV and C III] emission regions, with the narrow C IV emission being blueshifted w.r.t. the systemic redshift. The second technique assumes that the emission lines can be decomposed into a sum of Gaussian components. The C IV is decomposed into a narrow (FWHM  $\sim 2600 \text{ km s}^{-1}$ ) and a broad (FWHM  $\sim 6300 \text{ km s}^{-1}$ ) component. In order to reproduce the small differences between the C IV and C III] lines, we need three Gaussian profiles (FWHM  $\sim 1550, 3400, 8550 \text{ km s}^{-1}$ ) to reproduce the latter. The lightcurves derived for these components are essentially flat and show that microlensing is more important when the FWHM increases. Although the individual Gaussian line components do not necessarily isolate individual emission regions, we find that these lightcurves are compatible with the continuum lightcurves, provided only the size of the emission region is modified. This allows us to derive a half-light radius for the regions emitting these components as well as their size relative to the continuum. The radii are



consistent with the virial hypothesis and a radius which varies as  $\text{FWHM}^{-2}$ . Using the virial theorem, we derive a black hole mass  $M_{BH} \sim 2.0 \times 10^8 M_\odot$  ( $0.5 \times 10^8 M_\odot < M_{BH} < 5.4 \times 10^8 M_\odot$  at 68.3% confidence).

Our analysis supports the finding by other authors (Brotherton et al. 1994a, Marziani et al. 2010) that the regions emitting the C IV and C III] lines are composed of at least two spatially distinct components, one emitting the narrow core of the line and another, more compact, emitting a broadest component. The broad (resp. narrow) component of the C IV and C III] lines do not have exactly the same profiles. The flux ratio C III]/C IV is very similar when measured in the broad and in the narrow components of the lines. This suggests that the ionisation parameter  $U$  is nearly the same in the two regions, a surprising result since the narrow components are found to arise in regions several times larger than the broad components. Other lines observed in our spectra seem to arise from a least two components: Mg II  $\lambda 2798$ , Al II  $\lambda 1671$ , He II  $\lambda 1640$ , Al III  $\lambda 1857$ . The situation is different for Si III]  $\lambda 1892$  which does not show emission from a broad component, indicative of a smaller electronic density  $n_e$  in the region emitting the broadest part of the emission lines. On the other hand, we detect broad microlensed Fe II+III but no “extended” emission. This suggests that  $\text{Fe}_{UV}$  is produced in the inner part of the BLR or in a very compact region. Obtaining spectro-photometric monitoring data in the near-infrared where Balmer lines are detectable would be very useful to constrain photo-ionisation models and compare the microlensing signal in Balmer (e.g. H $\beta$ ), high ionisation (e.g. C IV) and  $\text{Fe}_{opt}$  lines.

We have demonstrated that the spectrophotometric monitoring of microlensing in a lensed quasar is a powerful technique to probe the inner regions of quasars, to measure the size of the broad line region and infer its structure. More work is still needed to take full advantage of the method, but our results are very promising. Several improvements are possible to increase the accuracy on our size measurements and better characterize the BLR structure. First, microlensing simulations reproducing the signal in more than 2 lensed images should allow one to narrow the final probability distribution on the source size. Second, the implementation of spectral fitting using Markov-Chain Monte-Carlo should allow a more appropriate estimate of the error bars and more advanced modeling of the individual spectral components. Third, a fully coherent scheme should be developed in order to consistently model the emission line shape and the corresponding source intensity profile used in the simulation. These improvement will be the subject of a future work.

*Acknowledgements.* We thank B. Borguet for useful discussions, and the referee, T. Boroson, for valuable suggestions. DS acknowledges support from the Humboldt Foundation. This project is partially supported by the Swiss National Science Foundation (SNSF).

## References

- Abajas, C., Mediavilla, E., Muñoz, J. A. et al. 2002, *ApJ*, 576, 640  
 Agol, E., Gogarten, S. M., Gorjian, V., Kimball, A. 2009, *ApJ* 697, 1010  
 Anguita, T., Schmidt, R. W., Turner, E. L. et al. 2008, *A&A*, 480, 327

- Aoki, K., Yoshida, M. 1999, *ASPC* 162, 385  
 Assef, R.J., Denney, K.D., Kochanek, C.S. et al. 2010, *ApJ*, submitted (arXiv:1009.1145v1)  
 Bachev, R., Marziani, P., Sulentic, J. W. et al. 2004, *ApJ*, 617, 171  
 Baskin, A., Laor, A., 2004, *MNRAS* 350, L31  
 Baskin, A., Laor, A., 2005, *MNRAS* 356, 1029  
 Bentz, M. C., Peterson, B.M., Pogge, R.W., Vestergaard, M., Onken, C.A. 2006, *ApJ*, 644, 133  
 Bentz, M. C., Walsh, J. L., Barth, A. J. 2009, *ApJ*, 705, 199  
 Bentz, M. C., Horne, K., Barth, A. J. et al., 2010, *ApJ*, 720, L46  
 Bon, E., Popović, L. Č., Gavrilović, N., Mura, G. L., Mediavilla, E. 2009 *MNRAS*, 400, 924  
 Borguet, B., Hutsemékers, D. 2010, *A&A*, 515, 22  
 Boroson T. A., Green, R. F. 1992, *ApJS*, 80, 109  
 Brotherton, M. S., Wills, B.J., Steidel, C.C., Sargent, W.L.W. 1994 *ApJ*, 423, 131  
 Brotherton, M. S., Wills, B.J., Francis, P.J., Steidel, C.C. 1994 *ApJ*, 430, 495  
 Cardelli, J. A., Clayton, G. C., Mathis, J. S., 1989, *ApJ*, 345, 245  
 Chen, K., Halpern, J. P. 1989, *ApJ*, 344, 115  
 Clavel J., Reichert, G. A., Alloin, D. et al. 1991, *ApJ*, 366, 64  
 Collin, S., Huré, J.-M. 2001, *A&A*, 372, 50  
 Collin, S., Kawaguchi, T., Peterson, B. M., Vestergaard, M. 2006 *A&A*, 456, 75  
 Courbin, F., Magain, P., Kirkove, M., & Sohy, S. 2000, *ApJ*, 529, 1136  
 Congdon, A.B., Keeton, C.R., Osmer, S.J. 2007, *MNRAS*, 376, 263  
 Corbin, M.R. 1990, *ApJ*, 347, 346  
 Corbin, M.R. 1995, *ApJ*, 447, 496  
 Corbett, E., Robinson, A., Axon, D. J., Young, S., Hough, J. H. 1998, *MNRAS*, 296, 721  
 Decarli, R., Labita, M., Treves, A., Falomo, R. 2008, *MNRAS*, 387, 1237  
 Denney, K. D., Peterson, B. M., Pogge, R. W. et al. 2009, *ApJ*, 704, L80  
 Dietrich, M., Hamann, F., Appenzeller, I., et al. 2003, *ApJ*, 596, 817  
 Down, E. J., Rawlings, S., Sivia, D. S., Baker, J. C., 2010, *MNRAS*, 401, 633  
 Eigenbrod, A., Courbin, F., Sluse, D., et al. 2007, *A&A*, 480, 647 (Paper I)  
 Eigenbrod, A., Courbin, F., Meylan, G., et al. 2008, *A&A*, 490, 933 (Paper II)  
 Elvis, M. 2000, *ApJ*, 545, 63  
 Falco, E. E., Lehar, J., Perley, R. A., et al. 1996, *AJ*, 112, 897  
 Falco, E. E., Impey, C. D., Kochanek, C. S. et al. 1999, *ApJ*, 523, 617  
 Fine, S., Croom, S. M., Bland-Hawthorn, J. et al. 2010 arXiv1005.5287  
 Francis, P. J., Hewett, P.C., Foltz, C.B., Chaffee, F.H. 1992, *ApJ*, 398, 476  
 Fluke, C. J., Webster, R. L. 1999, *MNRAS*, 302, 68  
 Gaskell, C. M. *ApJ*, 263, 79  
 Gaskell, C. M., Klimek, E.S., Nazarova, L.S. arXiv0711.1025  
 Gaskell, C. M. 2009, *NewAR*, 53, 140  
 Gaskell, C. M. 2010a, *IAUS* 267, 203 (arXiv:1003.0036)  
 Gaskell, C. M. 2010b, *ApJ* submitted (arXiv:1008.1057)  
 Gil-Merino, R., Wambsganss, J., Goicoechea, L. J., Lewis, G. F. 2005, *A&A*, 432, 83  
 Gültekin, K., Richstone, D. O., Gebhardt, K. et al. 2009, *ApJ*, 698, 198  
 Horne, K., Peterson, B. M., Collier, S. J., Netzer, H. 2004, *PASP*, 116, 465  
 Hintzen, P., Maran, S.P., Michalitsianos, A.G. et al. 1990, *AJ*, 99, 45  
 Huchra, J., Gorenstein, M., Kent, S., et al. 1985, *AJ*, 90, 691  
 Hutsemékers, D., Surdej, J., van Drom, E. 1994, *Ap&SS*, 216, 361

- Hutsemékers, D., Borguet, B., Sluse, D., Riaud, P., Anguita, T., 2010, *A&A*, 519, 103
- Hu, C., Wang, J., Ho, L. C. et al. 2008, *ApJ*, 683, L115
- Jarvis, M. J., McLure, R. J. 2006, *MNRAS*, 369, 182
- Kollatschny, W., Dietrich, M. 1996, *A&A*, 314, 43
- Kaspi, S., Maoz, D., Netzer, H. 2005, *ApJ*, 629, 61
- Kaspi, S., Brandt, W. N., Maoz, D. et al. 2007, *ApJ*, 659, 997
- Kayser, R., Refsdal, S., Stabell, R. 1986, *A&A*, 166, 36
- Keenan, F. P., Kingston, A. E., Dufton, P. L. 1987, *MNRAS*, 225, 859
- Kochanek, C. S. 2004, *ApJ*, 605, 58
- Krolik, J. H., Horne, K., Kallman, T. R. et al. 1991, *ApJ*, 371, 541
- Laor, A. 2007, *ASPC*, 373, 384
- Lewis, G. F.; Irwin, M. J. 1995, *MNRAS*, 276, 103
- Lewis, G. F.; Irwin, M. J. 1996, *MNRAS*, 283, 225
- Lewis, G. F.; Belle, K. E. 1998, *MNRAS*, 297, 69
- Magain, P., Courbin, F., Sohy, S. 1998, *ApJ*, 494, 452
- Marconi, A., Axon, D. J., Maiolino, R. et al. 2008, *ApJ*, 678, 693
- Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., Moles, M. 1996, *ApJS*, 104, 37
- Marziani, P., Dultzin-Hacyan, D., Sulentic, J. W. 2006 *New Developments in Black Hole Research*, edited by Paul V. Kreidler, p.123
- Marziani, P., Sulentic, J. W., Negrete et al. 2010, *MNRAS*, accepted
- McLure, R. J., Dunlop, J. S. 2002, *MNRAS*, 331, 795
- Metcalfe, R. B., Moustakas, L. A., Bunker, A. J., et al. 2004, *ApJ*, 607, 43
- Morgan, C. W., Kochanek, C. S., Morgan, N. D., Falco, E. E. 2010, *ApJ*, 712, 1129
- Mortonson, M.J., Schechter, P.L., Wambsganss, J. 2005, *ApJ*, 628, 594
- Motta, V., Mediavilla, E., Munoz, J. A., Falco, E. 2004, *ApJ*, 613, 86
- Minezaki, T., Chiba, M., Kashikawa, N., Inoue, K. T., Kataza, H. 2009, *ApJ*, 697, 610
- Mushotzky, R. & Ferland, G. J. 1984, *ApJ*, 278, 558
- Murray, N., Chiang, J. 1995, *ApJ*, 454, 105
- Murray, N., Chiang, J. 1997, *ApJ*, 474, 91
- Netzer, H. 2003, *ApJ*, 583, L5
- Netzer, H., Marziani, P. *ApJ*, 2010, Accepted (arXiv:1006.3553)
- Onken, C. A., Ferrarese, L., Merritt, D. et al. 2004, *ApJ*, 615, 645
- Peterson B. M., Balonek, T. J., Barker, E. S. et al. 1991, *ApJ*, 368, 119
- Peterson, B. M., Bentz, M. C., Desroches, L.-B. et al. 2005, *ApJ*, 632, 799 (+Erratum)
- Poindexter, S., Kochanek, C.S. 2010a, *ApJ*, 712, 658
- Poindexter, S., Kochanek, C.S. 2010b, *ApJ*, 712, 668
- Popović, L. Č., Mediavilla, E. G., Jovanović, P.; Muñoz, J. A. 2003, *A&A*, 398, 975
- Popović, L. Č., Mediavilla, E., Bon, E., Ilić, D. 2004 *A&A*, 423, 909
- Popović, L. Č. 2006, *SerAJ*, 173, 1
- Rice, M. S., Martini, P., Greene, J. E. 2006, *ApJ*, 636, 654
- Richards, G. T., Vanden Berk, D. E., Reichard, T. A. et al. 2002, *AJ*, 124, 1
- Risaliti, G., Salvati, M., Marconi, A. 2010, arXiv1010.2037
- Rix, H. W., Schneider, D. P.; Bahcall, J. N. 1992, *AJ*, 104, 959
- Robinson, A. 1995, *MNRAS*, 272, 647
- Schmidt, R., Webster, R. L., Lewis, G. F. 1998, *MNRAS*, 295, 488
- Schneider, P. & Wambsganss, J. 1990, *AAP*, 237, 42
- Shakura, N.I., Sunyaev, R.A. 1973, *A&A*, 24, 337
- Sluse, D., Claeskens, J.-F., Hutsemékers, D., et al. 2007, *A&A*, 468, 885
- Sluse, D., Claeskens, J.-F., Hutsemékers, D., et al. 2008, *RMxAC*, 32, 83
- Snedden, S.A., Gaskell, M.C. 1999, *ApJ*, 521, 91
- Smith, J. E., Robinson, A., Young, S., Axon, D. J., Corbett, E. A. 2005, *MNRAS*, 359, 846
- Sulentic, J. W., Marziani, P., Zwitter, T., Calvani, M. 1995, *ApJ*, 438, L1
- Sulentic, J.W., Marziani, P. 1999, *ApJ*, 518, L9
- Sulentic, J. W., Marziani, P., Dultzin-Hacyan, D. 2000, *ARA&A*, 38, 521
- Udalski, A., Szymański, M., Kubiak, M., et al. 2006, *AcA*, 56, 293
- Vanden Berk, D. E., Richards, G. T., Bauer, A. et al. 2001, *AJ*, 122, 549
- Vestergaard, M., Wilkes, B. J., Barthel, P. D. 2000, *ApJ*, 538, L103
- Vestergaard, M., Wilkes, B. J., 2001, *ApJS*, 134, 1
- Vestergaard, M., Peterson, B. M. 2006, *ApJ*, 641, 689
- Wambsganss, J. 1990 PhD Thesis (Munich University)
- Wambsganss, J., Paczyński, B. 1991, *AJ*, 102, 864
- Wambsganss, J. 1992, *ApJ*, 386, 19
- Wambsganss, J., Paczyński, B. 1994, *AJ*, 108, 1156
- Wambsganss, J. 1999 *JCoAM*, 109, 353
- Wambsganss, J. 2001, *PASA*, 18, 207
- Wayth, R. B., O'Dowd, M., Webster, R. L. 2005, *MNRAS*, 359, 561
- Wills, B. J., Brotherton, M. S., Fang, D., Steidel, C. C., Sargent, W. L. W. 1993, *ApJ*, 415, 563
- Wilhite, B. C., Vanden Berk, D. E., Brunner, R. J., Brinkmann, J. V. 2006, *ApJ*, 641, 78
- Witt, H. J., Mao, S. 1994, *ApJ*, 429, 66
- Whittle, M. 1985, *MNRAS*, 213, 33
- Young, S. Axon, D. J., Robinson, A., Hough, J. H., Smith, J. E. 2007, *Nature*, 450, 74
- Yee, H.K.C., De Robertis, M.M. 1991, *ApJ*, 381, 386
- Zamfir, S., Sulentic, J. W., Marziani, P. 2008, *MNRAS*, 387, 856
- Zhu, L., Zhang, S. N., Tang, S. 2009, *ApJ*, 700, 1173

## Appendix A: Averaged emission line profile

We explain here how we construct an averaged emission line profile and how we use it to choose a reasonable multi-Gaussian decomposition of the broad emission lines. Following the prescriptions of the multi-component decomposition described in Sec. 2.1, we are able to subtract the quasar pseudo-continuum emission (including the Fe II+III emission) from the quasar spectra. This procedure leads to a spectrum containing only the broad emission lines. Then, we normalize their flux and compare the line profiles averaged over periods P1, P2 and P3. This procedure reveals that the profile of the C III] emission line remains basically unchanged over the whole monitoring period. In contrast, there is evidence for a small change of the C IV line profile between 2005 and 2007. The high signal to noise emission line spectra of C III] and C IV are then fitted with a sum of 3 Gaussian-line profiles. For C III], we find that the best fit is obtained with a narrow emission component with  $\text{FWHM} = 1540 \text{ km s}^{-1}$ , an intermediate emission component  $\text{FWHM}=3400 \text{ km s}^{-1}$  and a broad component with  $\text{FWHM}=8600 \text{ km s}^{-1}$ . For C IV, the emission line is well fitted with only two emission components with  $\text{FWHM} \sim 2570 \text{ km s}^{-1}$  and  $\text{FWHM} \sim 5150 \text{ km s}^{-1}$ . The absorption component is also fitted with a  $\text{FWHM} \sim 1250 \text{ km s}^{-1}$  for P1 and P2 and  $\text{FWHM} \sim 900 \text{ km s}^{-1}$  for P3. The difference of width of the absorption comes from the increased resolution of the spectra obtained during period P3. Higher resolution spectra (Hintzen et al. 1990, Yee & De Robertis 1991) show that the C IV absorption system is composed of at least 2 different absorption clouds associated with the quasar. Narrow absorption systems are also present in front of C III] emission (Motta et al. 2004) but they do not appear as significant features in our spectra and are therefore not part of our modelling.

If we now compare the decomposition of the line profile in images A & D, we find that the width of the individual Gaussian profiles are similar but not their relative intensities, as already suggested by the analysis performed in Paper I. During our fitting procedure, we assume that the FWHM of the Gaussian components are identical for A & D, except for C IV(BC) and C IV(AC). The widths are also fixed to the fiducial values derived here.

## Appendix B: Microlensing fit

We provide here the technical details regarding the comparison of the simulated tracks with data. The following source sizes are considered: 2, 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 240, 280, 320, 360, 400, 440, 480, 520, 560, 600, 640, 680, 720, 760, 800, 840, 880, 920, 960, 1000, 1040, 1080, 1120, 1160, 1200, 1240, 1280, 1320, 1360 and 1400 pixels. For each of the 60 source sizes  $R_s$ , we simulate 10000 pairs of lightcurves for both images A & D by tracing source trajectories across the magnification patterns. Bilinear interpolation is used for the track extraction. Each track is characterised by the set of parameters described in Sect. 4.2:  $p = (R_s, V, m_0, \theta, \mathbf{x}_{0,A}, \mathbf{x}_{0,D}, M)$ . Each simulated dif-

ference lightcurve  $\Delta m'_k(p)$  is then compared to the data  $D = \Delta m_k$  by measuring the goodness of fit with a  $\chi^2$  statistic:

$$\chi^2(p) = \sum_{k=1}^{n_{\text{obs}}} \left( \frac{\Delta m_k - \Delta m'_k(p)}{\sigma_k} \right)^2 \quad (\text{B.1})$$

where  $\sigma_k$  are the uncertainties of the data and  $n_{\text{obs}}$  is the number of data points. For OGLE data, we use those obtained between modified julian day (i.e. jd-2450000) 3100 and 4500, and we bin the data points obtained the same night, which leads to  $n_{\text{obs}} = 181$ . The error  $\sigma_k$  is the quadratic sum of the photometric error  $\sigma_{\text{OGLE}}$  provided by OGLE, of the standard deviation  $\sigma_{\text{bin}}$  between binned points, and of the systematic underestimation of the error  $\sigma_{\text{sys}}$  of the OGLE data. The error  $\sigma_{\text{sys}}$  is estimated using a polynomial fitting of the difference lightcurves (see Paper II) and is estimated to be  $\sigma_{\text{sys}} = 0.05 \text{ mag}$ . The data points corresponding to the BLR lightcurve have been described in Sect. 3.1.

The brute force approach which consists in randomly picking track parameters and estimate a  $\chi^2$  is very inefficient in getting even a few tracks reproducing the data unless a large amount of computing time is available. To overcome this problem, we follow the procedure explained in Paper II and, for each set of random parameters we have generated, we optimize the 6 parameters  $(m_0, \mathbf{x}_A, \mathbf{x}_D, \theta)$  with a  $\chi^2$  based minimization algorithm using a Levenberg-Marquardt least-square routine. Contrary to what has been done in Paper II, we do not optimize the track velocity to avoid possible bias on the velocity distribution induced by the minimization routine. In addition, we have windowed the magnification map in such a way that the starting point of a track is always more than 1 track length away from the border of the map. Our final library of tracks fitting OGLE data contains  $6 \times 10^5$  tracks (10000 track per source size) with a  $\chi^2/n_{\text{obs}} < 8.5$ .

The only parameter which is varied to compare the tracks of our library to the BLR lightcurves is the source size  $R_s$ . As explained in Sect. 4.2, we extract, for each track of our library (step 2), the same track in the 59 other magnification maps corresponding to the 59 source sizes. We then search for the best source size using a minimization routine and calculate tracks for arbitrary source sizes (in the range 0.01-14  $r_E$ ) by means of a quadratic spline interpolation between different source sizes.