

## Comparison of the Sequences of Class A $\beta$ -Lactamases and of the Secondary Structure Elements of Penicillin-Recognizing Proteins

B. JORIS,<sup>1</sup> P. LEDENT,<sup>1</sup> O. DIDEBERG,<sup>1</sup> E. FONZÉ,<sup>1</sup> J. LAMOTTE-BRASSEUR,<sup>1</sup> J. A. KELLY,<sup>2</sup>  
J. M. GHUYSEN,<sup>1</sup> AND J. M. FRÈRE<sup>1\*</sup>

*Laboratoire d'Enzymologie et Laboratoire de Cristallographie, Centre d'Ingénierie des Protéines, Institut de Chimie, B6, Université de Liège, B-4000 Sart Tilman (Liège 1), Belgium,<sup>1</sup> and Biochemistry and Biophysics, University of Connecticut, Storrs, Connecticut 06268<sup>2</sup>*

Received 15 April 1991/Accepted 26 August 1991

**The sequences of class A  $\beta$ -lactamases were compared. Four main groups of enzymes were distinguished: those from the gram-negative organisms and bacilli and two distinct groups of *Streptomyces* spp. The *Staphylococcus aureus* PC1 enzyme, although somewhat closer to the enzyme from the *Bacillus* group, did not belong to any of the groups of  $\beta$ -lactamases. The similarities between the secondary structure elements of these enzymes and those of the class C  $\beta$ -lactamases and of the *Streptomyces* sp. strain R61 DD-peptidase were also analyzed and tentatively extended to the class D  $\beta$ -lactamases. A unified nomenclature of secondary structure elements is proposed for all the penicillin-recognizing enzymes.**

In the last few years, many different  $\beta$ -lactamase sequences have become available. Most of them are active serine enzymes, and they have been divided into three major classes (classes A, C, and D) on the basis of their sequences. Crystallographic studies have revealed details of the three-dimensional structures of four class A and one class C  $\beta$ -lactamases and of a related penicillin-recognizing enzyme, the *Streptomyces* sp. strain R61 DD-peptidase. These data indicate, as expected, that most of the major structural features of the four class A enzymes are nearly superimposable and, in addition, that the peptidase and the class C enzyme appear to be built according to very similar patterns. Class A  $\beta$ -lactamases, for which the most detailed structural information is available, have been used as models for the whole group of active serine penicillin-recognizing enzymes. In this study we compared the sequences of class A  $\beta$ -lactamases and searched for corresponding areas in the other serine  $\beta$ -lactamases and the *Streptomyces* sp. strain R61 DD-peptidase.

**Origins of compared sequences.** (i) Class A  $\beta$ -lactamases. Class A  $\beta$ -lactamase sequences were those from *Bacillus licheniformis* 749/C, *Bacillus cereus* 569/H (type I), TEM-1, *Staphylococcus aureus* PC-1 (1), PSE-3 encoded by plasmid Rms149, and *Rhodospseudomonas capsulata* (8); SHV-1 (also called PIT-2) (5); *Bacillus cereus* 5/B (50); *Bacillus cereus* 569/H (type III) (24); LEN-1 (*Klebsiella pneumoniae*) (3); *Klebsiella oxytoca* E23004 (4); PSE-4 (7); *Actinomadura* sp. strain R39 (22); *Streptomyces albus* G (11); *Streptomyces lavendulae*, *Streptomyces badius*, and *Streptomyces fradiae* (15); and *Streptomyces aureofaciens* (2).

Two different enzymes secreted by *Streptomyces cacaoi* have been described. In this report, to distinguish the two sequences, we added the abbreviation of the institution where the gene was sequenced to the name of the strain, as follows: ULg and NDRE for Lenzini et al. (33) and Forsman et al. (15), respectively. A large number of extended-spectrum  $\beta$ -lactamases (SHV-2 and all the TEM enzymes) have been described recently. Sequencing of the corresponding genes often showed that they were derived from the parent

enzymes by a very small number of amino acid substitutions (see, for instance, references 6 and 36). Similarly, the OHIO-1 and the *Citrobacter diversus* ULA-27  $\beta$ -lactamases are closely related to the SHV-1 (47) and *K. oxytoca* (43) enzymes, respectively. Thus, only the TEM-1, SHV-1, and *K. oxytoca* enzymes were included in the comparison.

Unless otherwise stated, the ABL consensus numbering scheme of class A  $\beta$ -lactamases (2) is used throughout this report.

(ii) **Class C  $\beta$ -lactamases.** Class C  $\beta$ -lactamase sequences were those from *Escherichia coli* K-12 (26); *Citrobacter freundii* OS60 (34); *Enterobacter cloacae* P99, Q908R, and MHN1 (16); *Serratia marcescens* SR50 (38); and *Pseudomonas aeruginosa* (35).

(iii) **Class D  $\beta$ -lactamases.** The aligned sequences of class D  $\beta$ -lactamases were those from OXA-1 (41), OXA-2 (9), and PSE-2 (23) were used. A consensus numbering scheme for class D  $\beta$ -lactamases is proposed in this report. The putative active Ser residue was arbitrarily for class D  $\beta$ -lactamases assigned the number 70 (as in the ABL consensus numbering scheme).

(iv) **Penicillin receptor.** The penicillin-binding, C-terminal domain of the *blaR* gene has been expressed as a 26,000- $M_r$  soluble protein in *E. coli* (28). It is referred to as the BLAR-CTD protein, and its numbering is according to the class D consensus scheme.

(v) **DD-Peptidase.** The DD-peptidase was that from *Streptomyces* sp. strain R61 (14). The numbering system used is that of the mature protein.

### MATERIALS AND METHODS

The comparison matrix (Table 1) was constructed by comparing pairwise the sequences of the  $\beta$ -lactamases of class A with the help of two algorithms. The distances algorithm requires aligned sequences and compares them residue by residue (12). The alignments were those of Ambler et al. (2), but the C- and N-terminal portions were deleted so that all compared sequences extended from residues 30 to 285 (ABL consensus numbering scheme). The final score represents the number of matches divided by the length of the shorter sequence, excluding the gaps. In the

\* Corresponding author.

TABLE 1. Comparison of 20 class A  $\beta$ -lactamases

Enzyme	Comparison scores <sup>a</sup>																			
	LEN-1	Kl.o.	SHV-1	Rh.c.	PSE-3	PSE-4	TEM-1	A.R39	B.c.I	B.c.I5	B.c.III	Ba.l.	St.a.	S.au.	S.al.	S.c.Lg	S.c.N	S.ba.	S.fr.	S.la.
LEN-1		43	91	46	51	49	68	41	42	43	41	42	33	41	40	42	44	41	40	40
Kl.o.	-45		42	41	43	43	44	46	47	47	45	45	35	44	43	44	43	47	42	44
SHV-1	-111	-45		46	52	51	70	41	42	42	41	41	34	40	41	42	44	40	41	41
Rh.c.	-54	-45	-54		46	45	47	40	36	36	36	39	29	37	40	36	38	39	37	37
PSE-3	-57	-46	-57	-55		50	49	48	45	47	42	44	36	41	42	46	43	45	44	42
PSE-4	-58	-50	-60	-54	-64		47	40	42	42	39	38	40	37	37	37	41	40	40	38
TEM-1	-92	-48	-92	-55	-58	-56		42	44	43	42	43	36	42	42	45	46	43	42	40
A.R39	-42	-47	-42	-39	-50	-41	-40		66	66	63	64	44	46	47	54	61	65	49	49
B.c.I	-46	-49	-46	-38	-48	-48	-46	-74		90	68	67	47	47	46	56	55	61	45	44
B.c.I5	-46	-52	-46	-38	-50	-50	-47	-76	-103		68	66	47	47	47	56	55	61	47	44
B.c.III	-45	-53	-45	-39	-49	-47	-46	-74	-81	-84		76	49	45	48	53	53	57	49	47
Ba.l.	-47	-52	-47	-41	-46	-44	-48	-75	-81	-83	-94		52	41	47	53	55	62	47	45
St.a.	-33	-36	-35	-24	-33	-36	-36	-45	-50	-52	-54	-57		35	35	44	43	45	35	33
S.au.	-43	-44	-43	-34	-41	-38	-46	-41	-44	-48	-47	-43	-32		59	44	45	46	65	65
S.al.	-44	-49	-45	-34	-42	-42	-44	-44	-51	-52	-53	-50	-33	-74		45	46	49	68	68
S.c.Lg	-40	-38	-41	-32	-39	-37	-43	-52	-57	-60	-55	-54	-45	-40	-46		58	60	46	44
S.c.N	-48	-48	-47	-42	-46	-48	-52	-68	-65	-70	-66	-68	-46	-48	-50	-60		68	44	44
S.ba.	-45	-51	-45	-40	-47	-43	-46	-70	-69	-72	-69	-72	-47	-46	-50	-62	-86		48	46
S.fr.	-47	-46	-48	-33	-46	-44	-50	-48	-49	-51	-54	-52	-34	-81	-88	-47	-52	-50		76
S.la.	-46	-50	-46	-34	-44	-43	-46	-49	-49	-50	-52	-50	-30	-78	-88	-44	-48	-49	-92	

<sup>a</sup> The positive values above the diagonal were calculated on the basis of the distances algorithm. They represent the percentage of very similar or identical residues (see text). The negative values below the diagonal were obtained with the help of the SEQDP algorithm. The gross values were divided by 10. Kl.o., *Klebsiella oxytoca*; Rh.c., *Rhodospseudomonas capsulata*; A.R39, *Actinomadura* sp. strain R39; B.c.I, B.c.I.5, and B.c.III:  $\beta$ -lactamases I, 15b, and III, respectively, of *Bacillus cereus*; Ba.l., *Bacillus licheniformis*; St.a., *Staphylococcus aureus*; S.au., *Streptomyces aureofaciens*; S.al., *Streptomyces albus* G; S.c.Lg, *Streptomyces cacaoi* ULg; S.c.N, *Streptomyces cacaoi* NDRE; S.ba., *Streptomyces badius*; S.fr., *Streptomyces fradiae*; S.la., *Streptomyces lavendulae*.

present analysis, a match was scored when the substitution had a value greater than 1 in the table of Gribskov and Burgess (19), which is a normalized form of the table of Dayhoff (10). The chosen value corresponds to a high degree of structural conservation, since identical residues yield a score of 1.5, while the mean value is zero.

The SEQDP algorithm gives a score for the best alignment of two sequences, as described by Goad and Kanehisa (18) and Needleman and Wunsch (39). The final score is the sum of the individual scores obtained for each pair of residues minus a gap penalty of 8. The original table of Dayhoff (10) was used, and the scores varied from 0 to -17, with the lower value representing identical residues. The sequences that were compared were the same as those above, from residues 30 to 285 (ABL consensus numbering scheme), but the gaps were introduced by the program itself, in contrast to the distances algorithm, in which the gaps were imposed a priori.

Secondary structure predictions were performed as described by Robson and Garnier (44).

## RESULTS AND DISCUSSION

**Amino acid sequence comparisons. (i) Class A  $\beta$ -lactamases.** The alignment of the class A  $\beta$ -lactamase sequences has been described by Ambler et al. (2). Table 1 presents the comparison matrices which were obtained with residues 30 to 285 of the ABL numbering scheme. It should be stressed that the deletion of the N- and C-terminal residues increased the similarities between the various enzymes, since the alignments show that these areas exhibit higher degrees of variation (2).

Both comparison methods yielded very similar results (Table 1). The SEQDP algorithm generally yields higher scores, since it uses a replacement matrix, while the other procedure counts matches only above a certain level. Two

pairs of enzymes, SHV-1-LEN-1 and *B. cereus* I-*B. cereus* 5b, exhibited particularly high scores (greater than 90). They were not considered to be different proteins in the following analysis. The enzymes of *S. aureus* and *R. capsulata* consistently exhibited poor scores. Moreover, they were extremely different from each other; indeed, the score obtained by comparing them with each other was the lowest score of all. To position each enzyme in the complete population, we computed an average score as the mean of all the scores (the distances program was used) for a given enzyme, with the standard deviation representing the heterogeneity of its behavior. Those averages ranged from  $39.5 \pm 5$  for the *R. capsulata* and *S. aureus* enzymes to  $51 \pm 10$  for the *B. licheniformis* enzyme and were generally higher for the *Streptomyces* and *Bacillus*  $\beta$ -lactamases than for those originating from gram-negative species. Those scores were probably somewhat biased because of the presence of seven *Streptomyces* and four *Bacillus* enzymes in the sample. As a consequence, we tried to group the various enzymes into a limited number of enzyme families. The basic idea was that a given enzyme should exhibit a significantly higher average score against its own enzyme family than against the others. This resulted in the classification shown in Table 2 comprising four enzyme families and three "loners," in which each enzyme nicely fulfills that condition. The major surprise was that the *Streptomyces* enzymes fell into two distinct enzyme families and that one of them was much closer to the enzyme family of the bacilli than to that of the other *Streptomyces* spp. Moreover, the *Streptomyces* I enzyme family could not even be considered as an intermediate between the bacilli and the *Streptomyces* II enzyme families since the average score of the latter enzyme family was higher with the bacilli than with the other *Streptomyces* spp. enzyme families (46.5 versus 45.6, although the difference is probably not meaningful). The three loners were the *S. aureus*, the *K. oxytoca*, and the *R. capsulata* enzymes. Although the *S. aureus* and

TABLE 2. Classification of the enzymes and score for each family

Enzyme	Score for enzyme family:			
	1 (gram-negative bacteria)	2 (bacilli)	3 ( <i>Streptomyces</i> I)	4 ( <i>Streptomyces</i> II)
LEN-1 (SHV-1)	56.8	42.2	41.5	40.5
TEM-1	58.5	42.8	44.7	41.5
PSE-3	50.5	45	44.7	42.3
PSE-4	49.8	40	39.3	37.8
Mean	54 ± 4.4	42.5 ± 2	42.6 ± 2.6	40.5 ± 2
<i>Actinomadura</i> sp. strain R39	42.4	64	60	47.8
Bcl (Bcl 5b)	43.2	72.6	57.3	45.9
BcIII	41	69	54.3	47.3
<i>B. licheniformis</i>	41.6	68	56.7	45
Mean	42.1 ± 1	68.4 ± 3.5	57 ± 2.3	46.5 ± 1.3
<i>S. cacaoi</i> ULG	42.4	54.4	59	44.8
<i>S. cacaoi</i> NDRE	42.8	55.8	63	44.8
<i>S. badius</i>	41.8	61.2	64	47.3
Mean	42.3 ± 0.5	57 ± 3.6	62 ± 2.6	45.6 ± 1.4
<i>S. aureofaciens</i>	40.6	45.2	45	63
<i>S. albus</i> G	40.2	47	46.7	65
<i>S. fradiae</i>	41.4	47.4	46	69.7
<i>S. lavendulae</i>	40.2	45.8	44.7	69.7
Mean	40.6 ± 0.6	46.4 ± 1.0	45.6 ± 1.0	66.9 ± 3.4
<i>K. oxytoca</i>	43 ± 1	46 ± 1	44.7 ± 2.1	43.3 ± 1
<i>S. aureus</i>	35.8 ± 2.7	47.8 ± 3	44 ± 1	34.5 ± 1
<i>R. capsulata</i>	46 ± 1	37.4 ± 2	37.7 ± 1.5	37.8 ± 1.5
ROB-1	42.2 ± 2.5	53.8 ± 1.5	48.7 ± 1.5	44.2 ± 0.5
LIPEN	41 ± 2.3	69 ± 8	55 ± 1.7	44 ± 1

the *K. oxytoca* enzymes appeared to be closer to those of the bacilli and the *R. capsulata* enzyme appeared to be closer to those of the gram-negative bacteria, because of their low scores, they could not be included in those families without destroying the homogeneity of the bacilli and gram-negative bacteria enzyme families. As noted above, the three loners were also extremely different from each other. It should be noted, however, that the *K. oxytoca* enzyme was not a real loner, since the enzyme of *Citrobacter diversus* certainly belongs to the same enzyme family. The *Actinomadura* sp. strain R39 enzyme was much closer to the bacilli than to the *Streptomyces* enzymes, which is a somewhat surprising behavior for a member of the *Actinomycetales* taxonomic family. Finally, in the gram-negative bacteria, the enzymes LEN-1, SHV-1, and TEM-1 formed a rather homogeneous group. On the contrary, the enzymes PSE-3 and PSE-4 could not be considered to be more related to each other than to the enzymes from other gram-negative bacteria.

We further tested our classification with two other enzymes which were inadvertently omitted during the first analysis. Data for those two enzymes, ROB-1 (29) and LIPEN (30), appear at the bottom of Table 2, where they are compared with the various enzyme families. Clearly, LIPEN presents all the characteristics of a member of the bacilli enzyme family, including a good similarity with the *Streptomyces* I enzyme family. Although the ROB-1 enzyme was more similar to the bacilli enzyme family, it cannot be included in that family without drastically decreasing its

average internal score. It could be considered a distant relative, but it also exhibits the same preference for the *Streptomyces* I enzyme family as do the true members of the bacilli enzyme family.

At present, it is not possible to build a phylogenetic tree with the proteins compared here. Pastor et al. (42) have recently proposed a tree, some aspects of which are at variance with our classification. The reasons for those differences are not clear, but it remains difficult to understand how the *K. oxytoca* and *S. albus* G enzymes, which, when compared by our criteria, exhibited a very low score of 43, could be considered to be closely related by those investigators.

(ii) **Class C  $\beta$ -lactamases and the *Streptomyces* sp. strain R61 DD-peptidase.** Alignment of the five known distinct sequences of class C  $\beta$ -lactamases (16, 35, 38) reveals 93 invariant residues (26%), and this does not allow an easy pinpointing of functionally important conserved residues. However, by introducing the *Streptomyces* sp. strain R61 DD-peptidase into the comparison, a more significant identification of such conserved areas could be performed (27). Recent three-dimensional data show that Y159 in the DD-peptidase superimposes on Y150 in the  $\beta$ -lactamases, which suggests that the YSN sequence of the peptidase corresponds to the YAN sequence of the class C  $\beta$ -lactamase enzymes. Moreover, the hydroxyl group of the Y residue superimposes on that of S130 in class A  $\beta$ -lactamases.

(iii) **Class D  $\beta$ -lactamases and BLAR-CTD.** A surprisingly

					40
OXA1	....MKNTIH	INFAIF.LII	ANIIYSSASA	STDISTVASP	LFEGTE...G
OXA2	.....MAIR	IFAILFSIFS	LATFAHAQEG	TLERSDARK.	FFSEFQA.KG
PSE2	.....M	KTFAAYVIA	CLSTALAGS	ITENTSANK	.FSA.EAVNG
BLAR-CTD	.....	.....	.....PGT	NVEYDYST.	FFDKFSAS.G
Consensus	-----	-----	-----	-E-----	F--A--G
			70		90
OXA1	CFLLYD.AST	NAEIAQFNKA	KCATQMAPDS	TFKIALSLMA	FDAEII.DOK
OXA2	TIVVADERQA	DRAMLVFDPV	RSKKRYSPAS	TFKIPHTLFA	LDAGAVRDEF
PSE2	VFVLCKSSSK	SCATN..DLA	RASKEYLPAS	TFKIPNAIIG	LETGVIKNEH
BLAR-CTD	GFVLFN.SNR	KKYT.IYNRK	ESTSRFAPAS	TYKVSALLA	LESGIITKND
Consensus	-FVL-----	-----	-----PAS	TFKI--L-L-A	L--G-I--G
				140	
OXA1	TIFKDKKTPK	GMEIHNNSHT	PKTHMDFSVV	WVSQEIQKI	RLNKIKNYLK
OXA2	QIFRWDGVR	GFAGHNDQDD	LRSAHRNSTV	WYELFAKEI	GDDKARRYLK
PSE2	QVFKHDGKPR	AMKQIERDLT	LRGAIQVSAV	PVFQIAREV	GEVRMQKYLK
BLAR-CTD	SHMTDGTQY	PYKEWINDQDD	LFSAMSSSTT	WYFQKLDRI	GEDHLRHYLK
Consensus	--F--LDG--	----WN-D--	L--AM--S-V	WV-Q-----	I-G-----YLK
				190	
OXA1	DFDYGNQDFS	GDKERNNGLT	EAWLESSLKI	SPEEQIQFLR	KIINHLPVK
OXA2	KIDYGNADPS	TSNG.....	DYWIEGSLAI	SAEQEIAFLR	KLYRNELPFR
PSE2	KFSYGNQNIS	GG.....ID	KFWLEGQLRI	SAVNDQVLE	SLYLNKLSAS
BLAR-CTD	SIHYGNEDFS	VP.....A	DYWLDGSLQI	SPLQGVNLIK	KFYDNEFDK
Consensus	---YGN-D-S	-----	---LEGLS-I	S--EQ--FL-	K-Y-N-L---
				240	
OXA1	NSAIENTIEN	MYLQDLNST	KLYGKTGAGF	TANRT....	LONGWFEFGI
OXA2	.VEHORLVKD	LMIVEAGRNI	ILRAKTGHEG	RMG.....	HWVGAVE...
PSE2	K.ENQLIVKE	ALVTEAAPEY	LVHSGTGFSG	VGTESNPGVA	HWVGAVE...
BLAR-CTD	QSNIE.TVKD	SIRLEESNGR	VLSGKTGTSV	INGELHAG..	WFIGYVE...
Consensus	-----VK-	---E---	-L--KTG--	-----	W--GWVE---
				290	
OXA1	ISKSGHKYVF	VSALTGNLGS	NLTSSIKAKK	NAITILNTLN	L.....
OXA2	..WPTGVSFF	ALNIDTPNRM	DDLFKREAIIV	RAILRSIEAL	PPNPVNSDA AR
PSE2	..KETEVYFF	AFNMDIONES	KLPLRKS IPT	KIMESGIIIG	G.....
BLAR-CTD	..TADNTFFF	AVHIQGEKRA	AGSSAAEIAL	SILDKGIYP	SVSR.....
Consensus	-----FF	A-----	-----	-----	-----

FIG. 1. Aligned sequences of BLAR-CTD and class D  $\beta$ -lactamases. Strictly invariant residues are underlined. A residue was considered as a consensus when it was present in at least three of the proteins.

high degree of similarity has been found between the OXA-2  $\beta$ -lactamase and the C-terminal domain of the product of the *blaR* gene (BLAR-CTD) (51), although BLAR-CTD exhibits only penicillin-binding properties and an exceedingly low hydrolytic activity. Figure 1 shows the aligned sequences, and Table 3 provides results of the comparisons that were performed as described above for the class A enzymes. It was remarkable to observe that the BLAR-CTD fragment exhibited higher scores with OXA-2 and PSE-2 than OXA-1. In fact, those scores characterize by far the highest degree of similarity ever found between a  $\beta$ -lactamase and a penicillin-binding protein. By contrast, by using the SEQDP algorithm and the same scale, the comparison between the *Streptomyces* sp. strain R61 DD-peptidase and a class C  $\beta$ -lactamase

TABLE 3. Comparison matrix for the class D enzymes

Enzyme	Comparison scores <sup>a</sup>				
	$\beta$ -Lactamase			BLAR-CTD	
	OXA-1	OXA-2	PSE-2	<i>B. licheniformis</i>	Tn552
OXA-1		30	30	33	31
OXA-2	-25		43	44	33
PSE-2	-28	-43		39	35
<i>B. licheniformis</i>	-32	-42	-40		47
Tn552	-31	-30	-31	-57	

<sup>a</sup> For details regarding the scores, see footnote a of Table 1.




FIG. 2. General arrangement of the secondary structure (A) and the four conserved structural elements (B) in class A  $\beta$ -lactamases. (A) A ribbon representation of the polypeptide chain of the *S. albus*  $\beta$ -lactamase. The dashed lines indicate possible direct connections (see text) in class D enzymes. (B) The circles indicate the positions of the  $\alpha$  carbon of the conserved residues composing the four structural elements described in the text. 1a is Ser-89 and ABL-70 and 1b is Lys-92 and ABL-73 of structural element 1. 2a is Ser-153 and ABL-130, 2b is Asp-154 and ABL-131, and 2c is Asn-155 and ABL-132 of structural element 2. 3 is Glu-191 and ABL-166 of structural element 3. 4a is Lys-259 and ABL-234, 4b is Thr-260 and ABL-235, and 4c is Gly-261 and ABL-236 of structural element 4. The numbering of the *S. albus* G residues includes the signal peptide. It should be noted that two errors were found in the sequence published by Dehottay et al. (11): residue Gly-128 should be eliminated and the tripeptide Ala-Gln-Leu at residues 140 to 142 should be replaced by the dipeptide Gly-Met. The complete sequence is thus two residues shorter (312 residues). Taking into account these corrections, the secondary structure assignments are as follows (numbers are residue numbers):  $\alpha$ 1, 48 to 60 (ABL 28 to 40);  $\beta$ 1, 63 to 70 (ABL 43 to 50);  $\beta$ 2, 76 to 79 (ABL 56 to 60);  $\alpha$ 2, 89 to 102 (ABL 70 to 83);  $\alpha$ 2a, 118 to 124 (ABL 99-103);  $\alpha$ 3, 133 to 138 (ABL 111a to 115);  $\alpha$ 4, 142 to 151 (ABL 119 to 128);  $\alpha$ 5, 155 to 165 (ABL 132 to 142);  $\alpha$ 6, 168 to 177 (ABL 145 to 154);  $\alpha$ 8, 206 to 217 (ABL 185 to 194);  $\alpha$ 9, 224 to 235 (ABL 201 to 212);  $\alpha$ 10, 244 to 247 (ABL 221 to 224);  $\beta$ 3, 253 to 260 (ABL 230 to 237);  $\beta$ 4, 266 to 273 (ABL 244 to 251);  $\beta$ 5, 280 to 287 (ABL 259 to 269);  $\alpha$ 11, 297 to 310 (ABL 276 to 289).

yielded a score of -12.4 (27) which, in that previous analysis, was the highest inter-enzyme family score.

A gene homologous to the *blaR* gene has been found in Tn552, a transposable element from *S. aureus* (45). Its sequence has been determined, and its C-terminal portion is included in the comparison presented on Table 3.

**General comparison.** The presently available X-ray structures show that four class A  $\beta$ -lactamases (13, 21, 37, 46) and one class C  $\beta$ -lactamase (40) and the R61 DD-peptidase (31) are composed of two domains, one  $\alpha/\beta$  and one all  $\alpha$  (Fig. 2).

TABLE 4. Four structural elements that limit the active site<sup>a</sup>

Enzyme	Sequence and residue no.			
	Element 1	Element 2	Element 3	Element 4
<i>S. albus</i> G	S-V-F-K 70	S-D-N 130	E-P-E-L-N 166	D-K-T-G 233
<i>S. aureus</i>	S-T-S-K	S-D-N	E-I-E-L-N	D-K-S-G
<i>B. licheniformis</i>	S-T-I-K	S-D-N	E-P-E-L-N	D-K-T-G
<i>C. freundii</i>	S-V-S-K 64	Y-A-N 150	D-A-E-A-Y <sup>b</sup> 217	H-K-T-G 314
<i>Streptomyces</i> sp. strain R61	S-V-T-K 62	Y-S-N 159	D-S-T-E-Q 225	G-H-T-G 297
Class D <sup>c</sup>	S-T-F-K 70	Y-G-N 144	E-X-X-L-X 175	X-K-T-G 214
BLAR-CTD <sup>c</sup>	S-T-Y-K 70	Y-G-N 144	D-G-S-L-Q 175	G-K-T-G 214

<sup>a</sup> The numbering of the various proteins is as follows: class A  $\beta$ -lactamases, ABL (2), *C. freundii*, consensus class C (16); *Streptomyces* sp. strain R61: mature protein sequence (14); class D and BLAR-CTD, see Fig. 1.

<sup>b</sup> The identification of element 3 in the class C  $\beta$ -lactamase of *C. freundii* is tentative since the C<sub>α</sub> coordinates of the enzyme are not available.

<sup>c</sup> For the class D  $\beta$ -lactamases and the BLAR-CTD, the identification of elements 1 and 4 cannot be easily challenged, but that of elements 2 and 3 is based only on sequence alignments.

The penicillin-binding site is in a crevice at the interface between the two domains. In the six enzymes, the active site is limited by four structural elements which occupy similar positions (Table 4).

(i) **Element 1.** Element 1 is the sequence -S\*-X-X-K-, where S\* is the active serine, at the N terminus of the  $\alpha$ 2 helix.

(ii) **Element 2.** Element 2 is the SDN loop of class A  $\beta$ -lactamases, corresponding, respectively, to YAN and YSN sequences in class C enzymes and the DD-peptidase. This loop is between helices  $\alpha$ 4 and  $\alpha$ 5 in class A  $\beta$ -lactamases. However, in class A  $\beta$ -lactamases, that serine appears to be involved in maintaining the functional positioning of the two domains (25) and in the protonation of the substrate's leaving group (32), roles which might be very different from that of the tyrosine in class C  $\beta$ -lactamases, which could act as a general base in the catalytic phenomenon (40).

(iii) **Element 3.** Element 3 is an acidic residue situated just before or at the N terminus of a one-turn helix or of a loop whose first three residues present helix-like characteristics. The residue is E in class A  $\beta$ -lactamases and D in class C  $\beta$ -lactamases and the peptidase. In class A  $\beta$ -lactamases, site-directed mutagenesis indicates that E166 (ABL consensus numbering scheme) plays an important role in catalysis (17). Accordingly, its side chain points into the substrate-binding cavity. In class C  $\beta$ -lactamases and the *Streptomyces* sp. strain R61 DD-peptidase, the side chain does not point into the active site, and modifications of that residue do not appear to alter the enzyme activity (20, 49). Even if, in those cases, it plays no functional role in catalysis, the reproducible presence of that residue might still represent another element that is common to all the structures.

(iv) **Element 4.** Element 4 is the KT(S)G sequence on the  $\beta$ 3 strand of class A  $\beta$ -lactamases facing the SDN loop on the other side of the active serine. The same KTG sequence is found on the  $\beta$ 7 strand of the class C enzyme and HTG on the  $\beta$ 3 strand of the DD-peptidase, in similar positions.

Finally, it is interesting to remember that the SXXX and KTG elements are also found in all penicillin-binding pro-

teins that have been described so far. Moreover, a SXN sequence, which might correspond to element 2 is also found in all penicillin-binding protein sequences (48).

Those four elements can serve as markers to compare the folding of the four classes of enzymes discussed here. By using the numbering of the helices and  $\beta$  strands as they occur in the sequence in class A  $\beta$ -lactamases (Fig. 3), the following conclusions can be made.

(i) The residues preceding the active Ser contribute one helix and two  $\beta$  strands to the  $\alpha/\beta$  domain.

(ii) The active serine is at the N terminus of the hydrophobic helix  $\alpha$ 2, which is followed by a loop containing one short helix ( $\alpha$ 3) in the *B. licheniformis* and *S. aureus* class A  $\beta$ -lactamases and in the *Streptomyces* sp. strain R61 DD-peptidase. The *S. albus* G  $\beta$ -lactamase contains one additional helix ( $\alpha$ 2a), and the class C enzyme contains two additional short helices. This is also a hydrophilic part of the molecule where a short (12-residue) insertion occurs in class C  $\beta$ -lactamases and the DD-peptidase.

(iii) Thereafter, six helices occur successively in the  $\beta$ -lactamases ( $\alpha$ 4 to  $\alpha$ 9). When it is present, the fourth helix ( $\alpha$ 7) is very short (one turn) and sometimes reduces to a loop where two or three residues present helix-like angle values. Active-site element 2 is between  $\alpha$ 4 and  $\alpha$ 5 in all the enzymes, and element 3 is just at the beginning of the short helix  $\alpha$ 7. Helix  $\alpha$ 8 is at the border between the all  $\alpha$  and the  $\alpha/\beta$  domains. It is in contact with the  $\beta$  sheet and might play an important role in the stability of the whole structure. In this all- $\alpha$  part of the molecules, two insertions appear to occur in the class C enzyme and the DD-peptidase, one between  $\alpha$ 4 and  $\alpha$ 5 and the other between  $\alpha$ 6 and  $\alpha$ 7. This might explain why the orientation of the  $\alpha$ 4 helix is somewhat different in the class A enzyme and the DD-peptidase.

(iv) The C-terminal part of the polypeptide chain belongs to the  $\alpha/\beta$  domain. Structures common to all the enzymes are, in the class A nomenclature, helices  $\alpha$ 10 and  $\alpha$ 11;  $\beta$  strands  $\beta$ 3,  $\beta$ 4, and  $\beta$ 5; and loops between  $\alpha$ 9 and  $\alpha$ 10 and between  $\alpha$ 10 and  $\beta$ 3. In addition to these structures, the class C enzymes contain three short  $\beta$  strands between  $\alpha$ 9 and  $\alpha$ 10 and another one between  $\alpha$ 10 and  $\beta$ 3. This could




FIG. 3. Analysis of the secondary structures of the *Streptomyces* sp. strain R61 DD-peptidase and the class A and C  $\beta$ -lactamases. The distances (in residues) between each secondary structure element are given. The circles represent  $\alpha$  helices, and the jagged lines represent the  $\beta$  strands. Heavy bars indicate the areas where important insertions are observed in the *Streptomyces* sp. strain R61 (S. R61) and class C enzymes. The consensus elements bordering the active site are highlighted in the lower part of the figure. The proposed ABL numbering of the secondary structure elements, corresponding to that of the *S. aureus* enzyme, is used for the four enzymes. When different, the original numbering is also shown under the corresponding value of the ABL consensus numbering scheme. In the proposed unified nomenclature, a secondary structure which is absent in the reference protein is assigned the number of the preceding helix or strand with an additional letter. To illustrate this point, two class A  $\beta$ -lactamases are included. In the *S. albus* G enzyme, an additional helix is found between *S. aureus* helices 2 and 3. It is referred to as helix  $\alpha_{2a}$ . Conversely, in the *S. albus* G enzyme, helix  $\alpha_7$  reduces to a loop and cannot be considered to be a helix. The additional  $\beta$  strands in the class C  $\beta$ -lactamase are accordingly numbered  $\beta_{2a}$ ,  $\beta_{2b}$ ,  $\beta_{2c}$ , and  $\beta_{2d}$ , since they are inserted between  $\beta_2$  and  $\beta_3$ . In the upper part of the figure, the numberings of the S.R61 DD-peptidase and of the *S. albus*  $\beta$ -lactamase include the signal peptides.

explain one major difference between the class C  $\beta$ -lactamases and the enzymes of the other classes. In class C  $\beta$ -lactamases, helix  $\alpha_{10}$  appears to run parallel to  $\alpha_1$  and antiparallel to  $\alpha_{11}$ , while it has the opposite orientation in the DD-peptidase and in class A (Fig. 4). The  $\alpha/\beta$  domain contains only one active-site element (KTG), which is on  $\beta_3$  in all cases.

At this point, we wish to stress again that, on the basis of the available knowledge, the role of structural elements 2 and 3 might be quite different in class A and C  $\beta$ -lactamases. Starting with the rather safe assumption that a general base is necessary for activating the hydroxyl of the active Ser residue, that general base would be E166 (element 3) in class A  $\beta$ -lactamases and the anion of Y150 (element 2) in class C  $\beta$ -lactamases.

One can also attempt to fit the sequence of class D  $\beta$ -lactamases to the general model, on the basis of the conserved active-site elements and secondary structure pre-




FIG. 4. Organization of the  $\alpha/\beta$  domains in the class A and *Streptomyces* sp. strain R61 enzymes (A) and in class C enzymes (B).

dictions. These predictions show that the active serine is at the end of an  $\alpha/\beta$  area and at the beginning of a hydrophobic helix. There are 74 residues between the active serine and the YGN element; thus, between those elements there are 30 more residues in the class D  $\beta$ -lactamases than there are in class A  $\beta$ -lactamases, but this is also an insertion area in both the DD-peptidase and the class C  $\beta$ -lactamases. By contrast, the distance between the YGN sequence and the third element (E140) is only 15 residues, and no secondary structure is predicted in that area. The class A structure shows that a direct passage from  $\alpha 5$  to  $\alpha 7$  involving the deletion of  $\alpha 6$  might be possible. Similarly, the distance between E140 and the KTG sequence is only 49 residues, i.e., 19 residues shorter than in class A  $\beta$ -lactamases. Assuming that helix  $\alpha 8$  is necessary for the stability of the molecule, one might predict, again on the basis of the class A structure, a direct passage from the C-terminal end of  $\alpha 9$  to the N terminus of  $\beta 3$ , thus involving the deletion of  $\alpha 10$ . Between the active Ser and residue 187, just before KTG, only helices are predicted. In the C-terminal portion (residues 187 to 255), predictions indicate an  $\alpha/\beta$  domain. The proposed shortcuts in class D enzymes are shown in Fig. 2A.

In conclusion, the class A  $\beta$ -lactamase model can be used as a reference for all serine  $\beta$ -lactamases and probably for many DD-peptidases. We propose that, in addition to the ABL consensus numbering scheme (2), the numbering of class A secondary structure elements also be adopted for all active serine penicillin-recognizing enzymes, as we have done in Fig. 3.

#### ACKNOWLEDGMENTS

This work was supported, in part, by the Belgian Programme on Interuniversity Poles of Attraction initiated by the Science Policy Programming Prime Minister's Office, Belgian State (PAI 19); an action concertée with the Belgian government (convention 86/91-90); the fonds de la Recherche Scientifique Médicale (contract 3.4537.88); and a convention tripartite between the Région Wallonne, SmithKline Beecham, United Kingdom, and the University of Liège. B.J. is chercheur qualifié of the Fonds National de la Recherche Scientifique (Brussels). P.L. is research fellow of the Institut pour l'Encouragement de la Recherche Scientifique dans l'Industrie et l'Agriculture (Brussels).

#### REFERENCES

- Ambler, R. P. 1980. The structure of  $\beta$ -lactamases. *Philos. Trans. R. Soc. London B Biol. Sci.* **289**:321-331.
- Ambler, R. P., A. F. Coulson, J. M. Frère, J. M. Ghuysen, M. Forsman, B. Joris, R. Levesque, G. Tiraby, and S. G. Waley. 1991. A standard numbering scheme for the class A  $\beta$ -lactamases. *Biochem. J.* **276**:269-270.
- Arakawa, Y., M. Ohta, N. Kido, Y. Fujii, T. Komatsu, and N. Kato. 1986. Close evolutionary relationship between the chromosomally encoded  $\beta$ -lactamase gene of *K. pneumoniae* and the Tem  $\beta$ -lactamase gene mediated by R plasmid. *FEBS Lett.* **207**:69-74.
- Arakawa, Y., M. Ohta, N. Kido, M. Mori, H. Ito, T. Komatsu, Y. Fujii, and N. Kato. 1989. Chromosomal  $\beta$ -lactamase of *Klebsiella oxytoca*, a new class A enzyme that hydrolyzes broad-spectrum  $\beta$ -lactam antibiotics. *Antimicrob. Agents Chemother.* **33**:63-70.
- Barthelemy, M., J. Peduzzi, and R. Labia. 1988. Complete amino acid sequence of p453-plasmid-mediated PIT-2  $\beta$ -lactamase. *Biochem. J.* **251**:73-79.
- Barthelemy, M., J. Peduzzi, H. B. Yaghlane, and R. Labia. 1988. Single amino-acid substitution between SHV-1  $\beta$ -lactamase and cefotaxime-hydrolysing SHV-2 enzyme. *FEBS Lett.* **231**:217-220.
- Boissinot, M., and R. C. Levesque. 1990. Nucleotide sequence of the PSE-4 carbenicillinase gene and correlations with the *Staphylococcus aureus* PC1  $\beta$ -lactamase crystal structure. *J. Biol. Chem.* **265**:1225-1230.
- Campbell, J. I. A., S. Scahill, T. Gibson, and R. P. Ambler. 1989. The phototrophic bacterium *Rhodospseudomonas capsulata* sp108 encodes an indigenous class A  $\beta$ -lactamase. *Biochem. J.* **260**:803-812.
- Dale, J. W., D. Godwin, D. Massakowska, P. Stephenson, and S. Wall. 1985. Sequence of the OXA2  $\beta$ -lactamase: comparison with other penicillin-reactive enzymes. *FEBS Lett.* **191**:39-44.
- Dayhoff, M. O. 1979. Atlas of protein sequence and structure, vol 5, suppl. 3. National Biochemical Foundation, Silver Spring, Md.
- Dehottay, P., J. Dusart, F. De Meester, B. Joris, J. Van Beeumen, T. Erpicum, J. M. Frère, and J. M. Ghuysen. 1987. Nucleotide sequence of the gene encoding the *Streptomyces albus* G  $\beta$ -lactamase precursor. *Eur. J. Biochem.* **166**:345-350.
- Devereux, J., P. Haeblerli, and O. Smithies. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.* **12**:387-395.
- Dideberg, O., P. Charlier, J. P. Wery, P. Dehottay, J. Dusart, T. Erpicum, J. M. Frère, and J. M. Ghuysen. 1987. The crystal structure of the  $\beta$ -lactamase *Streptomyces albus* G at 0.3 nm resolution. *Biochem. J.* **245**:911-913.
- Duez, C., C. Piron-Fraipont, B. Joris, J. Dusart, M. Urdea, J. Martial, J. M. Frère, and J. M. Ghuysen. 1987. Primary structure of the *Streptomyces* R61 extracellular DD-peptidase. *Eur. J. Biochem.* **162**:509-518.
- Forsman, M., B. Häggström, L. Lindgren, and B. Jaurin. 1990. Molecular analysis of  $\beta$ -lactamases from four species of *Streptomyces*: comparison of amino acid sequences with those of other  $\beta$ -lactamases. *J. Gen. Microbiol.* **136**:589-598.
- Galleni, M., F. Lindberg, S. Normark, S. Cole, N. Honoré, B. Joris, and J. M. Frère. 1988. Sequence and comparative analysis of three *Enterobacter cloacae ampC*  $\beta$ -lactamase genes and their products. *Biochem. J.* **250**:753-760.
- Gibson, R. M., H. Christensen, and S. G. Waley. 1990. Site-directed mutagenesis of  $\beta$ -lactamase I. Single and double mutants of Glu-166 and Lys-73. *Biochem. J.* **272**:613-619.
- Goad, W. B., and M. I. Kanehisa. 1982. Pattern recognition in nucleic acid sequences. I. A general method for finding local homologies and symmetries. *Nucleic Acids Res.* **10**:247-263.
- Gribskov, M., and R. R. Burgess. 1986. Sigma factors from *E. coli*, *B. subtilis*, phage T4 are homologous proteins. *Nucleic Acids Res.* **14**:6745-6763.
- Hadonou, A. M., and B. Joris. Unpublished data.
- Herzberg, O. 1991. Refined structure of  $\beta$ -lactamase from *Staphylococcus aureus* PC1 at 2.0 Å resolution. *J. Mol. Biol.* **217**:701-719.
- Houba, S., S. Willem, C. Duez, C. Molitor, J. Dusart, J. M. Frère, and J. M. Ghuysen. 1989. Nucleotide sequence of the gene encoding the active-site serine  $\beta$ -lactamase from *Actinomyces* R39. *FEMS Microbiol. Lett.* **65**:241-246.
- Huovinen, P., S. Huovinen, and G. Jacoby. 1988. Sequence of PSE-2  $\beta$ -lactamase. *Antimicrob. Agents Chemother.* **32**:134-136.
- Hussain, M., F. J. Pastor, and J. O. Lampen. 1987. Cloning and sequencing of the *blaZ* gene encoding  $\beta$ -lactamase III, a lipoprotein of *B. cereus* 569/H. *J. Bacteriol.* **169**:579-586.
- Jacob, F., B. Joris, O. Dideberg, J. Dusart, J. M. Ghuysen, and J. M. Frère. 1990. Engineering a novel  $\beta$ -lactamase by a single point mutation. *Protein Eng.* **4**:79-86.
- Jaurin, B., and T. Grundström. 1981. AmpC cephalosporinase of *Escherichia coli* K12 has a different evolutionary origin from that of  $\beta$ -lactamases of penicillinase type. *Proc. Natl. Acad. Sci. USA* **78**:4897-4901.
- Joris, B., J. M. Ghuysen, G. Dive, A. Renard, O. Dideberg, P. Charlier, J. M. Frère, J. Kelly, J. Boyington, P. Moews, and J. Knox. 1988. The active-site-serine penicillin-recognizing enzymes as members of the *Streptomyces* R61 DD-peptidase family. *Biochem. J.* **250**:313-324.
- Joris, B., P. Ledent, T. Kobayashi, J. O. Lampen, and J. M. Ghuysen. 1990. Expression in *Escherichia coli* of the carboxy terminal domain of BLAR sensory-transducer protein of *Bacil-*

- lus licheniformis* as a water-soluble Mr-26000 penicillin-binding protein. FEMS Microbiol. Lett. 70:107-113.
29. Juteau, J.-M., and R. C. Levesque. 1990. Sequence analysis and evolutionary perspectives of ROB-1  $\beta$ -lactamase. Antimicrob. Agents Chemother. 34:1354-1359.
  30. Kato, C., Y. Nakano, and K. Horikoshi. 1989. The nucleotide sequence of the lipo-penicillinase gene of alkalophilic *Bacillus* sp. strain 170. Arch. Microbiol. 151:91-94.
  31. Kelly, J. A., J. R. Knox, H. Zhao, J. M. Frère, and J. M. Ghuysen. 1989. Crystallographic mapping of  $\beta$ -lactams bound to a D-alanyl-D-alanine peptidase target enzyme. J. Mol. Biol. 209:281-295.
  32. Lamotte-Brasseur, J., G. Dive, O. Dideberg, P. Charlier, J. M. Frère, and J. M. Ghuysen. 1991. Mechanism of acyl transfer by the class A serine  $\beta$ -lactamase of *Streptomyces albus* G. Biochem. J. 279:213-221.
  33. Lenzini, M. V., H. Ishihara, J. Dusart, H. Ogawara, B. Joris, J. Van Beeumen, J. M. Frère, and J. M. Ghuysen. 1988. Nucleotide sequence of the gene encoding the active-site serine  $\beta$ -lactamase from *Streptomyces cacaoi*. FEMS Microbiol. Lett. 49:371-376.
  34. Lindberg, F., and S. Normark. 1986. Sequence of the *Citrobacter freundii* chromosomal *ampC*  $\beta$ -lactamase gene. Eur. J. Biochem. 156:441-445.
  35. Lodge, J. M., S. D. Michin, L. J. V. Piddock, and S. J. W. Busby. 1990. Cloning and analysis of the structural gene and regulatory region of the *Pseudomonas aeruginosa* chromosomal *ampC*  $\beta$ -lactamase. Biochem. J. 272:627-631.
  36. Mabilat, C., S. Goussard, W. Sougakoff, R. C. Spencer, and P. Courvalin. 1990. Direct sequencing of the amplified structural gene and promoter for the extended-broad-spectrum  $\beta$ -lactamase TEM-9 (RHH-1) of *Klebsiella pneumoniae*. Plasmid 23:27-34.
  37. Moews, P. C., J. R. Knox, O. Dideberg, P. Charlier, and J. M. Frère. 1990.  $\beta$ -lactamase of *Bacillus licheniformis* 749/C at 2 Å resolution. Proteins Struct. Funct. Genet. 7:156-171.
  38. Momura, K., and T. Yoshida. 1990. Nucleotide sequence of the *Serratia marcescens* SR50 chromosomal *ampC* gene. FEMS Microbiol. Lett. 70:295-300.
  39. Needleman, S. B., and C. D. Wunsch. 1970. A general method applicable to the search for similarities in the amino acid sequence of two proteins. J. Mol. Biol. 48:443-453.
  40. Oefner, C., A. Darcy, J. J. Daly, K. Gubernator, R. L. Charnas, I. Heinze, C. Hubschwerlen, and F. K. Winkler. 1990. Refined crystal structure of beta-lactamase from *Citrobacter freundii* indicates a mechanism for beta-lactam hydrolysis. Nature (London) 343:284-288.
  41. Ouellette, M., L. Bissonnette, and P. Roy. 1987. Precise insertion of antibiotic resistance determinants into Tn21 like transposons: nucleotide sequence of the OXA-1  $\beta$ -lactamase gene. Proc. Natl. Acad. Sci. USA 84:7378-7382.
  42. Pastor, N., D. Piñero, A. M. Valdés, and X. Soberón. 1990. Molecular evolution of class A  $\beta$ -lactamases: phylogeny and patterns of sequence conservation. Mol. Microbiol. 4:1957-1965.
  43. Perilli, M., N. Franceschini, B. Segatore, G. Amicosante, A. Oratore, C. Duez, B. Joris, and J. M. Frère. FEMS Microbiol. Lett., in press.
  44. Robson, B., and J. Garnier. 1986. Introduction to proteins and protein engineering. Elsevier, Amsterdam.
  45. Rowland, S.-J., and K. G. H. Dyke. 1990. Tn552, a novel transposable element from *Staphylococcus aureus*. Mol. Microbiol. 4:961-975.
  46. Samraoui, B., B. Sutton, R. Todd, P. Artimyuk, S. G. Waley, and D. Phillips. 1986. Tertiary structure similarity between a class A  $\beta$ -lactamase and a penicillin-sensitive D-alanyl-D-alanine carboxypeptidase-transpeptidase. Nature (London) 320:378-380.
  47. Shlaes, D. M., C. Currie-McCumber, A. Hull, I. Behlau, and M. Kron. 1990. OHIO-1  $\beta$ -lactamase is a part of the SHV-1 family. Antimicrob. Agents Chemother. 34:1570-1576.
  48. Spratt, B. G., and K. D. Cromie. 1988. Penicillin binding proteins of gram negative bacteria. Rev. Infect. Dis. 10:699-711.
  49. Tsukamoto, K., R. Kikura, R. Ohno, and T. Sawai. 1990. Substitution of aspartic acid-217 of *Citrobacter freundii* cephalosporinase and properties of the mutant enzymes. FEBS Lett. 264:211-214.
  50. Wang, W., P. S. F. Mezes, Y. Q. Yang, R. W. Blacher, and J. O. Lampen. 1985. Cloning and sequencing of the  $\beta$ -lactamase I gene of *Bacillus cereus* 5/B and its expression in *Bacillus subtilis*. J. Bacteriol. 163:487-492.
  51. Zhu, Y. F., I. H. A. Curran, B. Joris, J. M. Ghuysen, and J. O. Lampen. 1990. Identification of BlaR, the signal transducer for  $\beta$ -lactamase production in *Bacillus licheniformis*, as a penicillin-binding protein with strong homology to the OXA-2  $\beta$ -lactamase (class C) of *Salmonella typhimurium*. J. Bacteriol. 172:1137-1141.