



## COUPLING OF PRIMARY PRODUCTION AND CALCIFICATION AT THE CONTINENTAL MARGIN

Jérôme Harlay<sup>1</sup>, Lei Chou<sup>1</sup>, Nathalie Roevros<sup>1</sup>, Roland Wollast<sup>1</sup>, Bruno Delille<sup>2</sup>, Katrien Aerts<sup>3</sup>, Pascale-Emmanuelle Lapernat<sup>4</sup>

During the *Belgica* BG02/11 cruise (22 April - 11 May 2002) in the northern Bay of Biscay, a slope survey covering transects across the continental margin from the La Chapelle Bank to the Goban Spur area were conducted. The sampling campaign was assisted in addition by remote sensing data from SeaWiFS to locate the coccolithophore blooms, as characterised by white, highly reflective regions as shown by ocean colour imagery. Experiments of  $^{14}\text{C}$  incorporation were carried out along the shelf break to determine the production rate of both organic and inorganic particulate carbon in the photic zone.




SeaWiFS data received at the NERC Dundee University Receiving Station and processed by the Remote Sensing Group of the Plymouth Marine Laboratory.



Coccolithophores, among which *Emiliania huxleyi* is the most widespread species in the world's oceans, are the dominant calcifying phytoplankton in the area of investigation.

Their importance in biogeochemical cycling is supported not only by their capacity for organic matter production *via* **photosynthesis**, the **biological pump**, but also by their ability to produce calcified **coccoliths**, the **carbonate pump**, which represents a major contribution to the particulate carbon flux to the deep ocean.

In the photic zone, the **biological pump** removes inorganic carbon from the surface ocean, while the **carbonate pump**, according to the equation  $\text{Ca}^{2+} + 2\text{HCO}_3^- \rightleftharpoons \text{CaCO}_3 + \text{H}_2\text{O} + \text{CO}_2\uparrow$  releases  $\text{CO}_2$  and consumes Total Alkalinity to produce biogenic calcium carbonate. It is generally accepted that the overall effect of photosynthesis and calcification constitutes a net sink of carbon from the atmosphere. There remains still large uncertainties in the production and fate of biogenic calcium carbonate in the oceanic carbon cycle.




The primary production observed during our field survey was mainly associated to the coccolithophores. The total transfer of dissolved inorganic carbon to the particulate phase in the studied area is estimated to be  $1.44 \pm 0.40 \text{ gC m}^{-2} \text{ d}^{-1}$  (mean  $\pm$  std dev) for the shelf break and  $0.64 \pm 0.44 \text{ gC m}^{-2} \text{ d}^{-1}$  for the slope region. The elevated standard deviation associated to those numbers is due to the various stages encountered within the Coccolithophore bloom. The organic carbon produced is rapidly remineralised in the water column during settling, while the calcium carbonate resists much better to dissolution. Consequently, the carbon deposited and preserved in the sediments is mainly present as calcium carbonate and inorganic carbon flux becomes therefore significant at a global marine scale.

Recent investigations have shown that calcium carbonate could dissolve even in shallow and intermediate waters where this mineral phase is over-saturated. It is therefore important to understand the mechanisms associated with this process. This aspect constitutes the follow-up of the present research.

**Chlorophyll-a** concentrations increase in surface waters from less than 1 to more than 4  $\mu\text{g}\cdot\text{l}^{-1}$  as one moves from offshore (i.e. Station 7) towards the continental shelf (i.e. Station 5). Furthermore, there is a constant increase of Chl-a concentration from north (Station 16) to south (Station 2).


These concentrations are typical of those commonly observed during the spring phytoplankton bloom in this area.



The profiles of **total alkalinity** normalized to a salinity of 35 indicate a consumption by calcifying phytoplankton in surface waters. In the case of the offshore stations, the pattern of the vertical distribution is almost linear, denoting a minor effect of primary production on this parameter.


The large decrease observed in surface waters of station 16 indicate a late stage of the bloom, when all the coccoliths are spread, giving milky waters, whereas southern stations are in an early stage.

This is rather due to a temporal effect of the sampling than a physiological delay due to low latitudes.



<sup>14</sup>C uptake vs. light intensity experiments have allowed the characterisation of the photosynthetic properties of the phytoplankton. Organic carbon uptake is measured by acidification of the filters with Acetic Acid (0.01 M). The inorganic carbon uptake is obtained by differential scintillation counting between acidified filters and non-treated ones.



The parallelism of the two curves indicates that photosynthesis and calcification are intimately coupled. Carbon fluxes in  $\text{g C m}^{-2} \text{d}^{-1}$  have been calculated by integrating the optimal primary production (with the simulated light intensity of a sunny day and its related photoperiod) and the associated calcification rate.



Primary production and calcification rate are presented for the stations located on the shelf and represent the different stages of the bloom that we encountered during the cruise:

Station 5 was sampled at the beginning of the bloom. It's characterized by low levels of primary production, calcification as well as integrated Chl-a.

Station 8, 11, 14 and 16 were sampled during the middle stage and the end of the bloom, while the characteristic milky waters were observed, associated to a reduced Chl-a content (stations 14 and 16).



## AKNOWLEDGMENTS

This study is funded by the Belgian Federal Office for Scientific, Technical and Cultural Affairs (OSTC) under contract n° EV/11/5A. We would like to thank the captain, officers and crewmembers of the R/V Belgica for their logistic support. J. Backers and J.-P. De Blauwe from the Management Unit of the North Sea and Scheldt Estuary Mathematical Model (M.U.M.M.) in Ostend are greatly acknowledged for their assistance in the operation of the CTD rosette and in data acquisition. Data analysis and some representations are performed with ODV software, for which we are very grateful to Dr R. Schlitzer (<http://www.awi-bremerhaven.de/GEO/ODV/>)

Project Internet Site: <http://www.ulb.ac.be/sciences/dste/ocean/carbonate/frame.html>