

Carbon dioxide in European coastal waters

A.V. Borges(1), L.-S. Schiettecatte(1), G. Abril (2), B. Delille(1), F. Gazeau (3)

(1) University of Liège, (2) Université de Bordeaux 1, (3) Netherlands Institute of Ecology (alberto.borges@ulg.ac.be)

We compiled from literature annually integrated air-water fluxes of carbon dioxide (CO₂) computed from field measurements, in 19 coastal European environments that were gathered into 3 main ecosystems: inner estuaries, upwelling continental shelves and non-upwelling continental shelves. Air-water CO₂ fluxes were scaled at European regional level and compared to fluxes of atmospheric CO₂ in other aquatic and terrestrial compartments. Continental shelves are significant sinks for atmospheric CO₂ at an average rate of -1.9 molC/m²/yr that scaled at European level corresponds to an absorption of atmospheric CO₂ of -68.1 TgC/yr. This sink is equivalent to the one reported for the terrestrial biosphere of -66.1 TgC/yr, based on carbon-stock change models. Estuaries are significant sources of CO₂ to the atmosphere at an average rate of 49.9 molC/m²/yr that is higher than the CO₂ emission to the atmosphere from rivers and streams (26.9 molC/m²/yr) and lakes (7.6 molC/m²/yr). The scaled emission of CO₂ to the atmosphere from inner estuaries of about 67.0 TgC/yr would almost fully balance the sink of atmospheric CO₂ computed for continental shelves, and is higher than the emission of CO₂ to the atmosphere from continental aquatic systems of 36.5 TgC/yr. However, the scaled emission of CO₂ from estuaries to the atmosphere is inconsistent with the potential emission of CO₂ based on the fate of river organic carbon during estuarine transit. This discrepancy is most probably due to the poorly constrained surface area estimate of inner estuaries.