

# Multimodular Penicillin-Binding Proteins: An Enigmatic Family of Orthologs and Paralogs

COLETTE GOFFIN AND JEAN-MARIE GHUYSEN\*

Centre d'Ingénierie des Protéines, Université de Liège, Institut de Chimie, B-4000 Sart Tilman (Liège), Belgium

INTRODUCTION .....	1079
THE PENICILLOYL SERINE TRANSFERASES: A SUPERFAMILY OF MULTIPLE PERSONALITIES .....	1079
HIERARCHICAL ANALYSIS OF MULTIMODULAR PBPs .....	1082
CORE-BASED CLUSTERING OF MULTIMODULAR PBPs .....	1083
EVOLUTION OF MULTIMODULAR PBPs .....	1086
FUNCTIONS OF MULTIMODULAR CLASS A PBPs .....	1087
FUNCTIONS OF MULTIMODULAR CLASS B PBPs .....	1088
Class B PBPs of Gram-Negative Bacteria .....	1088
Class B PBPs of Gram-Positive Bacteria .....	1090
Class B PBPs in Cell Morphogenesis .....	1090
AN AMAZING PANOPLY OF MULTIMODULAR PBPs .....	1091
PENICILLIN-ORIENTED EVOLUTION .....	1091
TUNING UP THE CATALYTIC TWO-STROKE ENGINE .....	1091
CONCLUSIONS AND FUTURE DIRECTIONS .....	1092
ACKNOWLEDGMENTS .....	1092
REFERENCES .....	1092

## INTRODUCTION

Trypsin, chymotrypsin, and other acyl serine transferases catalyze the transfer of the electrophilic group  $R_1-CO$  of ester, thioester, and amide (peptide) carbonyl donors  $R_1-CO-X-R_2$  to an acceptor,  $HY$ , via the formation of a serine-ester-linked acyl ( $R_1-CO-$ ) enzyme intermediate.  $X$  denotes an oxygen atom, a sulfur atom, or an NH group.  $R_1$  and  $R_2$  denote the substituents of the scissile  $CO-X$  bond. When  $HY$  is  $H_2O$ , the carbonyl donor is hydrolyzed and the product of enzyme deacylation is  $R_1-COOH$ . When  $HY$  is an amino compound,  $NH_2-R_3$ , the carbonyl donor is transpeptidated and the product of enzyme deacylation is  $R_1-CO-NH-R_3$ .

Specialized acyl serine transferases are involved in the assembly and metabolism of the bacterial cell wall peptidoglycan. They have in common the ability to catalyze the rupture of the  $\beta$ -lactam amide bond of penicillin and the formation of a serine ester-linked penicilloyl enzyme. However, this intermediate is almost completely inert, the enzyme catalytic center turns over very slowly, once or less per hour, and the inactivated acyl serine transferases are easily detectable as penicillin-binding proteins (PBPs).

The production of  $\beta$ -lactamases of classes A, C, and D is a remarkable defensive mechanism that bacteria have developed to protect their wall peptidoglycan-synthesizing machinery against the toxic effect of penicillin. The serine  $\beta$ -lactamases hydrolyze penicillin into penicilloate via the formation of a serine ester-linked penicilloyl enzyme that is hydrolytically labile. On good  $\beta$ -lactam substrates,  $\beta$ -lactamases can turn over 1,000 times or more per second.

The PBPs and serine  $\beta$ -lactamases have been discussed in recent reviews (22, 24–28, 38, 39, 48, 49). This article focuses

on questions that biochemists still strive to answer concerning the multimodular PBPs, which, globally, are the lethal targets of penicillin in susceptible bacteria. To apprehend the problem, we shall first position the multimodular PBPs within the penicilloyl serine transferases superfamily (24).

## THE PENICILLOYL SERINE TRANSFERASES: A SUPERFAMILY OF MULTIPLE PERSONALITIES

The bacterial cell wall peptidoglycan (23) is a covalently closed, net-like polymer in which glycan strands are cross-linked by peptides (Fig. 1A). The glycan portion is made up of alternating  $\beta$ -1,4-linked units of *N*-acetylglucosamine and *N*-acetylmuramic acid arranged in linear chains. The carboxyl groups of the *N*-acetylmuramic acid residues are involved in amide linkages to terminal L-alanine residues of the peptide units L-alanyl- $\gamma$ -D-glutamyl-L-diaminoacyl-D-alanine. Also, neighboring peptide-substituted glycan strands are cross-linked by peptide bridges which extend from the carboxyl group of the terminal D-alanine of one peptide unit to the side chain amino group of the diamino acid residue of another peptide unit.

The nature of the diamino acid residue of the peptide units, the composition and length of the peptide bridges, and the extent of peptidoglycan cross-linking vary according to the bacterial species. In *Escherichia coli*, the diamino acid residue is *meso*-diaminopimelic acid, the glycan chains are substituted by peptide monomers and cross-linked peptide dimers, most of the interpeptide bridges are direct D-alanyl-(D)-*meso*-diaminopimelic acid bonds, and peptide oligomers larger than dimers have not been identified.

The immediate precursor of the wall peptidoglycan is lipid II (Fig. 1B). A disaccharide peptide is linked to a  $C_{55}H_{89}$  undecaprenyl lipid carrier via a pyrophosphate bridge involving  $C_1$  of *N*-acetylmuramic acid, and the peptide borne by *N*-acetylmuramic acid is a pentapeptide which terminates with the sequence D-alanyl-D-alanine. From this precursor, the assembly of lipid-transported disaccharide pentapeptide units into

\* Corresponding author. Centre d'Ingénierie des Protéines, Université de Liège, Institut de Chimie, B6, B-4000 Sart Tilman (Liège), Belgium. Phone: 32-4-366.33.95. Fax: 32-4-366.33.64. E-mail: jmghuysen@ulg.ac.be.




FIG. 1. (A and B) Structure of the wall peptidoglycan of *E. coli* (A) and assembly from the lipid II intermediate (B). G, *N*-acetylglucosamine; M, *N*-acetylmuramic acid; Dpm, *meso*-diaminopimelic acid; transmembrane bar, C<sub>55</sub>H<sub>89</sub> isoprenoid alcohol carrier; Mp, *N*-acetylmuramyl pentapeptide. (C) Reaction 1, glycan chain elongation at the nonreducing end of the chain; reaction 2, glycan chain elongation at the reducing end of the chain. (D) Overall reactions catalyzed by the DD-transpeptidases, DD-carboxypeptidases, and DD-endopeptidases.

polymeric peptidoglycan requires two enzymatic activities, a glycosyl transferase and an acyl serine transferase (66).

The glycosyl transferase-catalyzed transglycosylation proceeds through displacement of the pyrophosphate of lipid II by the 4-hydroxyl group of *N*-acetylglucosamine of the growing glycan chain (reaction 1 in Fig. 1C) or displacement of the pyrophosphate of the growing glycan chain by the 4-hydroxyl group of *N*-acetylglucosamine of lipid II (reaction 2 in Fig. 1C). This latter mechanism has been observed in *Bacillus licheniformis* and *Micrococcus luteus* (68, 69). In *Escherichia coli*, the absence of nascent peptidoglycan has been established, implying that its binding to preexisting peptidoglycan is concomitant with its synthesis.

The acyl serine transferase (transpeptidase)-catalyzed peptidoglycan cross-linking is made at the expense of the *D*-alanyl-*D*-alanine bond of the pentapeptide units (Fig. 1D). The reaction proceeds via the formation of a serine ester-linked peptidyl (-*L*-alanyl- $\gamma$ -*D*-glutamyl-*L*-diaminoacyl-*D*-alanyl) enzyme with the concomitant release of the carboxy-terminal *D*-alanine of the pentapeptide, and it is achieved by the transfer of the peptidyl moiety to the side chain amino group of the diamino acid residue of another peptide. Because the reaction involves breaking a *D*-alanyl-*D*-alanine bond, the transferase is classified as a DD-transpeptidase.

PBP1a and PBP1b of *E. coli* each catalyze the conversion of lipid II into peptidoglycan in *in vitro* assays (50). These two PBPs are the prototypes of bifunctional PBPs which combine

in a single polypeptide chain the required transglycosylase and DD-transpeptidase activities. Essentially, a noncleavable signal peptide which functions as a transmembrane anchor is fused to the amino end of a transglycosylase non-penicillin-binding (n-PB) module, which itself is fused to the amino end of an acyl serine transferase (DD-transpeptidase) penicillin-binding (PB) module. The two catalytic modules form a single polypeptide chain that folds on the exterior of the plasma membrane.

To allow the bacterial cell to grow and divide, morphogenetic networks channel peptidoglycan assembly into wall expansion and septum formation in a cell-cycle-dependent fashion. Central to these networks are PBPs which are similar in their modular design to the bifunctional (transglycosylase-acyl serine transferase) PBPs. However, the n-PB module is not a transglycosylase (1). *E. coli* PBP2 and PBP3 are involved in cell shape maintenance and cell division, respectively (53). They are the prototypes of bifunctional PBPs which combine in a single polypeptide chain a morphogenetic determinant n-PB module and an acyl serine transferase PB module.

Throughout the bacterial cell cycle, the wall peptidoglycan undergoes constant chemical changes that do not impair the tensile strength of the polymer (34). Monofunctional serine DD-carboxypeptidases/PBPs hydrolyze *D*-alanyl-*D*-alanine bonds (Fig. 1D). They control the extent of peptidoglycan cross-linking by limiting the number of pentapeptide units available for transpeptidation. Monofunctional serine DD-carboxypeptidases/PBPs also hydrolyze peptidoglycan interpeptide bonds,




FIG. 2. Acyl serine transferase-catalyzed reaction on  $R_1\text{—CO—X—}R_2$  carbonyl donors. (A) Overall reaction. (B) Role of a water molecule as proton transmitter. X is O, S, or NH (in penicillin, the CO—N bond is endocyclic). HY is an acceptor, i.e., water or an amino compound. The carbonyl of the donor and that of the acyl enzyme are polarized by NH groups of the enzyme polypeptide backbone. TS, transition state; \*, scissile bond of the carbonyl donor (step 1) and scissile bond of the acyl enzyme (step 2). The shaded area symbolizes the active-site environment.

which in some bacteria extend between two D-centers in the  $\alpha$ -position to a free carboxylate, for example a D-alanyl-(D)-*meso*-diaminopimelic acid bond (Fig. 1D). *E. coli* PBP4 through PBP7 are DD-carboxy/endopeptidases. Loss of these PBPs is tolerated (73). Their functions and those of many other peptidoglycan hydrolases presumably include a role in the recycling of old peptidoglycan, a role as zipper during cell division, and a role as space maker for the insertion of new peptidoglycan material.

The serine  $\beta$ -lactamases, the monofunctional PBPs, and the PB modules of the multimodular PBPs fulfill different functions, and similarity in their amino acid sequences is, globally, almost nonexistent. However, they operate on  $R_1\text{—CO—X—}R_2$  carbonyl donors by the same proton abstraction-donation mechanism, and their catalytic centers have a remarkably well-conserved topology.

In step 1 of the catalyzed reaction (Fig. 2A), the proton of the serine  $\gamma\text{OH}$  is abstracted, the activated  $\text{O}\gamma$  attacks the carbonyl carbon atom of the CO—X bond, and the abstracted proton is back-donated to the adjacent X atom, resulting in the formation of the serine-ester-linked acyl enzyme. In step 2 (for the reaction to reach completion), the proton of the acceptor HY (an amino group or water) is abstracted, the activated  $\text{Y}$  attacks the carbonyl carbon atom of the CO—O ester bond of the acyl enzyme and the abstracted proton is back-donated to the  $\text{O}\gamma$  atom of the serine residue.

The catalytic centers that perform this double proton shuttle

are defined by three amino acid groupings, referred to as motifs. These motifs occur in the same order and with roughly the same spacing along the polypeptide chains, defining a common amino acid sequence signature (Fig. 3). Polypeptide folding brings the three motifs close to each other, forming a cavity at the boundary between an all- $\alpha$  domain and an  $\alpha/\beta$  domain, which itself consists of a five-stranded  $\beta$ -sheet protected by  $\alpha$ -helices (38).

As shown in Fig. 4, motif 1,  $\text{S}_x\text{K}$  (where S is the essential serine residue and x is a variable amino acid residue), is at the amino end of helix  $\alpha_2$  of the all- $\alpha$  domain and occupies a central position in the catalytic center. Motif 2,  $[\text{S}/\text{Y}]_x[\text{N}/\text{C}]$ , is on a loop connecting two  $\alpha$ -helices of the all- $\alpha$  domain and defines one side of the catalytic center. Motif 3,  $[\text{K},\text{H}][\text{T},\text{S}]\text{G}$ , is on strand  $\beta_3$  of the  $\beta$ -sheet and defines the other side of the catalytic center. Although secondary structures vary in number, size, and orientation, the spanning distances between most of the heavy atoms of the side chains of the active-site-defining motifs 1, 2, and 3 differ by less than 1 Å (Fig. 4). Variations are observed at the top and bottom of the cavities. At this latter position, the side chain of the glutamic acid residue of motif  $\text{Ex}_2\text{LN}$ , found only in the class A  $\beta$ -lactamases, points toward the inside of the catalytic center.

This background of structural similarity and catalytic diversity illustrates the concept that evolution often obscures the function. In the absence of direct biochemical data, the function of a penicilloyl serine transferase cannot be identified on

Monofunctional PBPs		1		2		3			
DD-transpeptidase	S. K15*	NH <sub>2</sub>	34 <sup>35</sup> — Sx <sub>2</sub> K	57 <sup>96</sup> — SxC	114 <sup>213</sup> — KTG	46 <sup>213</sup> — KTG	COOH	262	
DD-carboxypeptidase	S. R61*	NH <sub>2</sub>	61 <sup>62</sup> — Sx <sub>2</sub> K	94 <sup>159</sup> — YxN	136 <sup>298</sup> — HTG	48 <sup>298</sup> — HTG	COOH	349	
	Eco 5	NH <sub>2</sub>	43 <sup>44</sup> — Sx <sub>2</sub> K	62 <sup>110</sup> — SxN	100 <sup>213</sup> — KTG	158 <sup>213</sup> — KTG	COOH	374	
	Bsu 5	NH <sub>2</sub>	35 <sup>36</sup> — Sx <sub>2</sub> K	60 <sup>100</sup> — SxN	124 <sup>227</sup> — KTG	183 <sup>227</sup> — KTG	COOH	413	
DD-endopeptidase	Eco 4	NH <sub>2</sub>	41 <sup>42</sup> — Sx <sub>2</sub> K	240 <sup>286</sup> — SxN	108 <sup>397</sup> — KTG	57 <sup>397</sup> — KTG	COOH	457	
<b>Multimodular PBPs</b>									
Tgase/Tpase	Eco 1a	NH <sub>2</sub>	464 <sup>465</sup> — Sx <sub>2</sub> K	55 <sup>524</sup> — SxN	189 <sup>716</sup> — KTG	131 <sup>716</sup> — KTG	COOH	850	
	Eco 1b	NH <sub>2</sub>	509 <sup>510</sup> — Sx <sub>2</sub> K	58 <sup>572</sup> — SxN	123 <sup>698</sup> — KTG	143 <sup>698</sup> — KTG	COOH	844	
Cell cycle	Eco 2	NH <sub>2</sub>	329 <sup>330</sup> — Sx <sub>2</sub> K	53 <sup>387</sup> — SxN	154 <sup>544</sup> — KSG	86 <sup>544</sup> — KSG	COOH	633	
	Eco 3	NH <sub>2</sub>	306 <sup>307</sup> — Sx <sub>2</sub> K	48 <sup>359</sup> — SxN	132 <sup>494</sup> — KTG	91 <sup>494</sup> — KTG	COOH	588	
	Spn 2x*	NH <sub>2</sub>	336 <sup>337</sup> — Sx <sub>2</sub> K	54 <sup>395</sup> — SxN	149 <sup>547</sup> — KSG	200 <sup>547</sup> — KSG	COOH	750	
<b>β-lactamases</b>									
Class A (Eco Tem)*		NH <sub>2</sub>	44 <sup>45</sup> — Sx <sub>2</sub> K	56 <sup>165</sup> — SxN	101 <sup>269</sup> — KTG	51 <sup>269</sup> — KTG	COOH	263	
Class C (Eclo P99)*		NH <sub>2</sub>	64 <sup>65</sup> — Sx <sub>2</sub> K	81 <sup>150</sup> — YxN	162 <sup>315</sup> — KTG	45 <sup>315</sup> — KTG	COOH	362	
Class D (Oxa2)		NH <sub>2</sub>	50 <sup>51</sup> — Sx <sub>2</sub> K	70 <sup>125</sup> — YxN	61 <sup>189</sup> — KTG	62 <sup>189</sup> — KTG	COOH	254	

α2                      α4 - α5                      β3

FIG. 3. Amino acid sequence signature of the penicilloyl serine transferases superfamily. S. K15, *Streptomyces* strain K15; S. R61, *Streptomyces* strain R61; Eco, *E. coli*; Bsu, *B. subtilis*; Spn, *S. pneumoniae*; Eclo, *E. cloacae*; Tgase, transglycosylase; Tpase, transpeptidase. PBPs and β-lactamases marked by an asterisk are of known three-dimensional structure (at 3.5-Å resolution only for Spn PBP2x). Motifs 1, 2, and 3 are at the top, and secondary structures are at the bottom. The monofunctional PBPs and β-lactamases are synthesized with a cleavable signal peptide. The R61 PBP and the β-lactamases are secreted; *E. coli* PBP4 is, somehow, loosely attached to the membrane; the *Streptomyces* strain K15 PBP is membrane associated via an internal hydrophobic segment; *E. coli* PBP5 and probably *B. subtilis* PBP5 are membrane associated via a carboxy-terminal amphiphilic helix. The multimodular PBPs are synthesized with a noncleavable signal peptide that serves as membrane anchor.

the basis of amino acid sequence or even three-dimensional structure comparisons.

#### HIERARCHICAL ANALYSIS OF MULTIMODULAR PBPs

By limiting our scrutiny to the small number of available sequences, it was proposed in 1991 (24) that, depending on the motifs borne by the n-PB modules, the multimodular PBPs fall into class A (whose prototypes are the *E. coli* bienzymatic PBP1a and PBP1b) and class B (whose prototypes are the *E. coli* cell cycle PBP2 and PBP3). In recent years, the databases have expanded considerably and several bacterial genomes have been sequenced (5, 12, 20, 21, 36, 41, 65). These advances allow for better standards in screening similarity and for improvement of our ability to apprehend the structure-function relationships and, perhaps, to manipulate the functions of the PBPs in wall peptidoglycan assembly and cell morphogenesis.

Sixty-three multimodular PBPs have been analyzed. They are listed in Table 1, with their identifier codes and accession numbers. Some PBPs have been characterized biochemically, and others have been identified only on the basis of genome sequence data. This applies, in particular, to the PBPs of the spiral-shaped *Helicobacter pylori* (the causative agent of peptic ulcer disease) and *Borrelia burgdorferi* (the causative agent of Lyme disease), the filamentous *Aquifex aeolicus*, and the cyanobacterium *Synechocystis* sp. strain PCC863. *A. aeolicus* is a member of the most deeply branching family within the bac-

terial domain and is one of the most thermophilic bacteria known, with growth temperature maxima near 95°C. *Synechocystis* sp. strain PCC863 carries genes for oxygenic photosynthesis and is also a very deeply branching family member. Chloroplasts are believed to have evolved from cyanobacterial ancestors which developed an endosymbiotic relationship with a eukaryotic host cell.

Classically, the relationships between genes from different genomes are represented as a system of orthologs and paralogs. Ortholog genes in different organisms have evolved from a common ancestor, and the encoded ortholog proteins normally retain the same function. Paralog genes are related by duplication within a genome, and paralog proteins normally evolve new functions. Often, the picture is more complex because genes may have been acquired by horizontal transfer, so that paralogs may have arisen from orthologs and vice versa.

To shed light on the structure-function and evolutionary relationships of the multimodular PBPs and the differential roles that they play in wall peptidoglycan assembly, the amino acid sequences of the 63 PBPs have been analyzed by hierarchically combining pairwise comparisons of two sequences, either a sequence and a preexisting alignment or two preexisting alignments (2, 45). By using PileUp from the Wisconsin package, the full-size PBPs generate the dendrograms shown in Fig. 5. The PBP identifier codes are at the bottom, the sequences joined toward the bottom have greater similarity than the sequences joined further to the top, and the vertical




FIG. 4. Schematic representation of the catalytic center of the penicilloyl serine transferases. Average distances in angstroms between heavy atoms (O-O, O-N, S-N, and N-N) of side chains of the active-site-defining motif 1 (on helix  $\alpha_2$ ), motif 2 (on a loop), and motif 3 (on strand  $\beta_3$ ) are given. The values derived from X-ray data apply to the proteins marked by an asterisk in Fig. 3. Motif 1 is, invariably, Sx<sub>2</sub>K. Motif 2, SxN, is replaced by SxC in *Streptomyces* strain K15 PBP and by YxN in *Streptomyces* strain R61 PBP and class C  $\beta$ -lactamases. Motif 3, K[T/S]G, is replaced by HTG in *Streptomyces* strain R61 PBP.

axis is roughly calibrated in similarity scores calculated for selected pairs of sequences.

A score is the standard deviation value above that expected from a run of 100 randomized pairs of sequences with the same amino acid composition as the two sequences under comparison. The score calculated for a pair of identical sequences varies depending on the length and, to some extent, the composition of the polypeptide chain. Thus, the scores calculated for *Bacillus subtilis* PBP1a (914 amino acid residues) and *Pseudomonas aeruginosa* PBP3a (565 amino acid residues) are 173 and 145, respectively. In spite of these limitations, dendrograms calculated for proteins with different molecular masses give a reasonably accurate picture of the hierarchical relationship. Scores larger than 15 express statistically significant similarity, scores smaller than 5 express lack of similarity, and scores between 15 and 5 define a cutoff region.

The results of this analysis (Fig. 5) are consistent with those described recently (13, 49, 55). The 63 full-size PBPs fall into two unbridgeable classes, A and B (score,  $<<5$ ). Class A comprises 29 PBPs. The PBPs of gram-negative bacteria fall into three clusters (whose prototypes are *E. coli* PBP1a, *Synechocystis* PBP1a, and *E. coli* PBP1b), the PBPs of gram-positive bacteria fall into three clusters (whose prototypes are *B. subtilis* PBP4, *Streptococcus pneumoniae* PBP1a, and *S. pneumoniae* PBP1b), and the mycobacterial PBPs form one distinct cluster. Class B comprises 34 PBPs. The PBPs of gram-negative bacteria fall into two clusters (whose prototypes are *E. coli* PBP2 and *E. coli* PBP3), and the PBPs of gram-positive bacteria fall into three clusters (whose prototypes are *Enterococcus faecium* PBP5, *S. pneumoniae* PBP2x, and *S. pneumoniae* PBP2b). Classes A and B contain PBP outliers, which fall outside the clusters.

#### CORE-BASED CLUSTERING OF MULTIMODULAR PBPs

To identify features that may be responsible for the clustering shown in Fig. 5, the amino acid sequences of the 63 full-size PBPs have been aligned in a way that highlights the class-specific motifs. In Fig. 6 and 7, the sequences for class A and class B PBPs, respectively, are presented in the same order from top to bottom as in the dendrogram of Fig. 5 from left to right. The sequences each start at the first dicarboxylic acid, D or E, immediately downstream from the hydrophobic sequence assumed to function as membrane anchor. The conserved motifs that contain identities are numbered at the top of the figures. Equivalent amino acid residues (A and G; K, R, and H; D and E; I, L, and V) are in boldface type. The intermotif distances are given as the number of amino acid residues. Since it is known that motif 4 of *S. pneumoniae* class B PBP2x is the fusion site between strand  $\beta_6$  at the carboxy end of the n-PB module and helix  $\alpha_1$  at the amino end of the PB module (57), motif 4 of all class B PBPs and the equivalent motif 6 of all class A PBPs are assumed to represent the intermodule junctions.

The picture which emerges from these alignments is one of distinctive motifs in the n-PB modules of class A PBPs versus class B PBPs, of conserved penicilloyl serine transferase motifs in the PB modules of both class A and class B PBPs, and of adducts occurring at various places along the polypeptide chains.

Cytosolic tails are present when the sequence between the amino end of the protein and the first dicarboxylic acid, D or E, is more than about 60 to 70 amino acid residues long. Carboxy-terminal extensions are present when the sequence between motif K[T/S]G and the carboxy end of the protein is

TABLE 1. Code, accession number, class, and size for multimodular PBPs

Source <sup>a</sup>	PBP code	Data bank <sup>b</sup>	Accession no.	Subclass <sup>c</sup>	Residue no.
Gram-positive bacteria					
<sup>+</sup> <i>Bacillus subtilis</i>	Bsu1	S	P39793	A3	914
	Bsu2c	S	P38050	A4	714
	Bsu4	S	P40750	A5	624
	Bsu3	S	P42971	B1	668
	BsuVD	S	Q03524	B3	645
	Bsu2b	S	Q07868	B4	716
	Bsu2a	S	P54488	B5	716
<i>Enterococcus faecalis</i>	Efas5	E	X78425	B1	679
<i>Enterococcus faecium</i>	Efam5	E	X84859	B1	673
	EfamC	E	U94707	B4	742
<i>Enterococcus hirae</i>	Ehi3r	P	A36903	B1	678
	Ehi5	E	X62280	B1	678
	Ehi3s	E	Y13922	B4	730
<i>Staphylococcus aureus</i>	Sau2	P	S43693	A3	716
	Sau2a	P	JQ0773	B1	668
	SauB	E	U94706	B4	646
<i>Staphylococcus sciuri</i>	Ssc2	E	Y09223	B1	666
<i>Streptococcus oralis</i>	Sor1a	S	Q00573	A3	637
<i>Streptococcus pneumoniae</i>	Spn1a	S	Q04707	A3	719
	Spn2a	E	AJ002292	A4	731
	Spn1b	E	AJ002291	A5	821
	Spn2x	S	P14677	B4	750
	Spn2b	S	P10524	B5	679
<i>Streptococcus pyogenes</i>	Spy263	O	contig 263	A3	721
	Spy250	O	contig 250	A4	778
	Spy286	O	contig 286	A5	723
	Spy290	O	contig 290	B4	752
<i>Streptococcus thermophilus</i>	Sth2b	E	U58210	B5	704
<i>Actinomycetales</i>					
<i>Mycobacterium leprae</i>	Mle1*	E	L39923	A	686
	Mle1	E	L01263	A	821
<i>Mycobacterium tuberculosis</i>	Mtu1	E	Z80775	A	665
<i>Streptomyces clavuligerus</i>	Sclpcbr	E	U56256	A:B	551
<i>Streptomyces coelicolor</i>	Scoa3	E	Y14206	B:B2	770
Gram-negative bacteria					
<i>Citrobacter freundii</i>	Cfr1b	P	S57580	A2	846
<sup>+</sup> <i>Escherichia coli</i>	Eco1a	S	P02918	A1	850
	Eco1b	S	P02919	A2	844
	Eco1c	E	U88571	A	770
	Eco2	S	P08150	B2	633
	Eco3	S	P04286	B3	588
<sup>+</sup> <i>Haemophilus influenzae</i>	Hin1a	S	P31776	A1	853
	Hin1b	S	P45345	A2	781
	Hin2	P	C64044	B2	651
	Hin3	P	G64184	B3	610
<i>Neisseria gonorrhoeae</i>	Ngo1	E	U72876	A1	798
	Ngo2	P	S49090	B3	582

Continued on following page

TABLE 1—Continued

Source <sup>a</sup>	PBP code	Data bank <sup>b</sup>	Accession no.	Subclass <sup>c</sup>	Residue no.
<i>Neisseria meningitidis</i>	Nme1	E	U80933	A1	798
	Nme2	E	X59624	B3	584
<i>Pseudomonas aeruginosa</i>	Pae1a	E	U73780	A1	822
	Pae3	P	S54872	B3	579
	Pae3a	E	X95517	B3	565
Spiral-shaped and filamentous gram-negative related bacteria					
+ <i>Helicobacter pylori</i>	Hpy1a	E	AE000573	A2	659
	Hpy2	E	AE000654	B:B2	588
	Hpy3	T	HP1556	B:B3	615
+ <i>Borrelia burgdorferi</i>	Bbu3	G	AE001173	A1	932
	Bbu2	G	AE001171	B:B2	599
	Bbu1	G	AE001125	B3	629
+ <i>Aquifex aeolicus</i>	Aaemrca	G	AE000699	A1	726
	AaeA1	G	AE000728	B2	595
	AaeA2	G	AE000695	B3	578
Cyanobacteria					
+ <i>Synechocystis</i> strain PCC6803	Syn1a	K	sll0002	A1	885
	Synmrca	K	sll1434	A1	650
	SynmrCb	K	slr1710	A1	749
	Syn3	K	sll1833	B3	607

<sup>a</sup> + denotes that the genome has been sequenced.

<sup>b</sup> E, EMBL DNA data bank; G, GenBank; K, Kazusa DNA Research Institute data bank at <http://www.kazusa.or.jp>; O, Oklahoma University DNA data bank at <http://www.genome.ou.edu>; P, PIR data bank; S, Swiss-Prot data bank; T, TIGR Microbial data bank at <http://www.tigr.org>.

<sup>c</sup> Letters without suffixes indicate that the PBP does not belong to a defined subclass. A:B indicates that the n-PB and PB modules of the PBP are of classes A and B, respectively. B:B2 and B:B3 indicate that the n-PB module of the PBP is an outlier of class B.

more than about 60 to 70 amino acid residues long. The carboxy-terminal extensions vary widely in size. Residues 780 to 844 in *E. coli* PBP1b are dispensable (37), but residues 762 to 780 at the carboxy end of the PB module are not (42). Internal inserts (boxed in Fig. 6 and 7) are also present. They occur between the membrane anchor and motif 1 in class A and class B PBPs, downstream from the junction site in class A PBPs, and between motifs 8 and 9 of the PB modules of class A PBPs.

As a result of these alignments, the cores of the n-PB modules can be defined as the sequences extending from the amino end of motif 1 to the carboxy end of motif 6 (class A) or motif 4 (class B). Likewise, the cores of the PB modules can be defined as the sequences starting 60 amino acid residues upstream from motif Sx<sub>2</sub>K and terminating 70 amino acid residues downstream from motif KTG or at the carboxy end of PBPs which have no carboxy-terminal extensions. On this basis, the hierarchical analysis of the cores of the n-PB modules (n-PB cores) and the cores of the associated PB modules (PB cores) leads to important observations.

The n-PB cores which bear the distinctive class-specific motifs fall into two unbridgeable groups, class A and class B (scores,  $\leq 5$ ), as expected. Contrary to expectations, the PB cores which all bear the penicilloyl serine transferases motifs also fall in the same unbridgeable classes A and B (scores,  $\leq 5$ ).

As shown in Fig. 8, the n-PB cores of the 29 class A PBPs form a continuum of diverging sequences which, from the *E. coli* PBP1a cluster to the *S. pneumoniae* PBP1b cluster, are related by scores larger than 15. In contrast, the associated PB cores of PBPs from gram-negative bacteria fall into two subclasses, A1 (whose prototype is *E. coli* PBP1a) and A2 (whose prototype is *E. coli* PBP1b), which form a supercluster; and

those of PBPs from gram-positive bacteria fall into three subclasses, A3 (whose prototype is *S. pneumoniae* PBP1a), A4 (whose prototype is *S. pneumoniae* PBP2a), and A5 (whose prototype is *S. pneumoniae* PBP1b), which also form a supercluster. It should also be noted that the PB cores of *A. aeolicus* PBP mrca, *Synechocystis* PBPs 1a, mrca, and mrCb, and *B. burgdorferi* PBP3 belong to subclass A1; the PB core of *H. pylori* PBP1a belongs to subclass A2; the PB cores of *Mycobacterium leprae* PBP1\* and *M. tuberculosis* PBP1 form a distinct cluster; and the PB cores of *E. coli* PBP1c, *M. leprae* PBP1, and *Streptomyces clavuligerus* PBP pCbr are outliers. *M. leprae* PBP1 is thermostable and binds penicillin with a very low affinity (3). In contrast, *M. leprae* PBP1\* is thermolabile and binds penicillin with a high affinity (43).

As shown in Fig. 9, the n-PB and PB cores of class B PBPs from gram-positive bacteria fall into three distinct subclasses, B1 (whose prototype is *Enterococcus faecium* PBP5), B4 (whose prototype is *S. pneumoniae* PBP2x), and B5 (whose prototype is *S. pneumoniae* PBP2b). Remarkably, an n-PB core of subclass B1, B4, or B5 is linked to a PB core of subclass B1, B4, or B5, respectively. As also shown in Fig. 9, the n-PB and PB cores of class B PBPs from gram-negative bacteria fall into two distinct subclasses, B2 (whose prototype is *E. coli* PBP2) and B3 (whose prototype is *E. coli* PBP3). Almost invariably, an n-PB core of subclass B2 or B3 is linked to a PB core of subclass B2 or B3, respectively. It should be noted that the n-PB cores of *H. pylori* PBP2, *B. burgdorferi* PBP2, and *Streptomyces coelicolor* PBP3a are outliers and that the corresponding associated PB cores belong to subclass B2 (thin and broken arrows in Fig. 9). Likewise, the n-PB core of *H. pylori* PBP3 is




FIG. 5. Hierarchical analysis of full-size, multimodular PBPs. PBP codes are given in Table 1. Each cluster (or subclass) is identified by a prototypic PBP. For SD score values and more details, see the text.

an outlier and the associated PB core belongs to subclass B3 (thick arrow in Fig. 9).

*Streptomyces clavuligerus* PBP pcbr is the only wild card which really breaks the pattern. The full-size PBP is of class B, the n-PB core is of class B, and the PB core is of class A. This PBP, whose gene is located downstream from the isopenicillin synthase-encoding gene in the cephamycin cluster, is probably responsible for the penicillin resistance of this organism (56).

The picture which results from the above analysis is almost certainly far short of being complete. As the structure databases continue to expand, existing subclasses will become more populated and new subclasses will be identified. Moreover, PBPs of the spiral-shaped *H. pylori* and *B. burgdorferi* and that of *Streptomyces coelicolor* suggest that mixed subclasses of PBPs may exist. Unrelated n-PB modules may be linked to PB modules belonging to a same subclass.

Adducts (Fig. 6 and 7; Table 2) allow the core-based clustering to be refined. Three PBPs of subclass A2 (Eco1b, Cfr1b, and Hin1b) have structurally related inserts downstream from the membrane anchor (>30% identity [Table 2]). The PBPs of subclass B1 (Efam5, Ehi3r, Ehi5, Efas5, Bsu3, Sau2a, and Ssc2) also have inserts at this position (Table 2). The enterococcal and *Bacillus* inserts are structurally related (>23% identity), the two staphylococcal inserts have 71% identity, and the two groups of inserts are distinctly related to each other (17% identity). Three PBPs of subclass A1 (Eco1a, Hin1a, and Pae1a) have inserts between motif 8 and motif 9 of the PB module which are structurally related (>30% identity [Table 2]). Six PBPs of subclass A1 (Eco1a, Hin1a, Pae1a, Nme1/Ngo1, Aaemrca, and Bbu3) have inserts located downstream from the intermodule junction (Table 2). The *E. coli*, *Haemophilus influenzae*, *P. aeruginosa*, and *Neisseria* inserts are structurally related (>25% identity).

## EVOLUTION OF MULTIMODULAR PBPs

The multimodular PBPs are an exemplary model of molecular evolution that gave rise to two classes of PBPs and to a prolific expansion of subclasses by fusion among several polypeptide chains (membrane anchor, n-PB core, PB core), acquisition of adducts, adaptive radiation, and speciation. Adaptive radiation is a term applied to the spread of species of common ancestry into different niches (70). It is extended in the present context to the spread of PBPs into the gram-negative bacteria, gram-positive bacteria, and other bacterial groups.

These structural changes occurred in combination with conservation of the core-specific amino acid sequence signatures. They resulted in the making of fully integrated polypeptide hybrids. These hybrids combine the main properties of the parental chains and are endowed with new properties because of noncovalent interactions between the constitutive modules.

Polypeptide folding and module swapping illustrate this notion. All the attempts made to produce the PB modules of *E. coli* PBP1b of subclass A2 (unpublished data), *E. coli* PBP3 of subclass B3 (29), *Enterococcus hirae* PBP5 of subclass B1 (51), and *Staphylococcus aureus* PBP2a of the same subclass B1 (71) as independent stable, penicillin-binding entities (using various expression-secretion vectors) failed. In contrast to the monofunctional PBPs, which are autonomous folding entities, the PB modules of multimodular PBPs have lost the ability to fold by themselves. They require the assistance of the associated n-PB modules, a property which implies precise and specific module-module interactions. While correct folding (in terms of penicillin binding) of the associated n-PB and PB modules is independent of the transporting signal peptide sequence, the in vivo activity of *E. coli* PBP3 is membrane anchor module dependent, indicating that the membrane anchor and its cytosolic

CLASS A PBPs	n-PB module						PB module	COOH								
	1	2	3	4	5	6										
<i>E. coli</i>	10 54	9 84	11 117	13 133	12 156	221 234	432 465	534 716								
<i>E. coli</i>	23 D 9	KRF 19	EDSRF <sub>2</sub> EHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 57	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 135	IN <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 55	SN 189	KTG 131	850			
<i>Hin1a</i>	E 10 D 9	QRR 19	EDSRF <sub>2</sub> DEHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 58	G <sub>2</sub> TTX <sub>2</sub> LX <sub>2</sub> Q 121	LN <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> S 17	S <sub>2</sub> X <sub>2</sub> K 54	SN 215	KTG 133	853			
<i>Pae1a</i>	E 16 D 9	MRR 19	EDDNF <sub>2</sub> NHX <sub>2</sub> G 21	GGSTITMQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> IL 57	G <sub>2</sub> TTX <sub>2</sub> RX <sub>2</sub> Q 131	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> SX <sub>2</sub> GG <sub>2</sub> S 17	S <sub>2</sub> X <sub>2</sub> K 55	SRN 172	KTG 124	822			
<i>Mme1</i>	D 16 D 9	QRR 19	EDKRF <sub>2</sub> RHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RQ <sub>2</sub> IL 57	G <sub>2</sub> TTX <sub>2</sub> RX <sub>2</sub> Q 129	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 56	SN 127	KTG 144	798			
<i>Ngol</i>	D 16 D 9	QRR 19	EDKRF <sub>2</sub> RHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RQ <sub>2</sub> IL 57	G <sub>2</sub> TTX <sub>2</sub> RX <sub>2</sub> Q 129	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 56	SN 127	KTG 144	798			
<i>Aaemrca</i>	E 11 D 11	QKR 19	EDRNF <sub>2</sub> HX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> KIMEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 55	R <sub>2</sub> X <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 107	ID <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> S 17	S <sub>2</sub> X <sub>2</sub> K 55	SIN 119	KTG 110	726			
<i>Hpy1a</i>	E 18 D 12	EFR 19	EDTLF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> IL 54	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 41	TD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 61	SN 119	KTG 102	659			
<i>Bsu4</i>	E 31 D 11	ENR 19	EDRHF <sub>2</sub> EHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> LL 83	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 22	IN <sub>2</sub> X <sub>2</sub> H <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> N 17	S <sub>2</sub> X <sub>2</sub> K 51	SYN 124	KTG 51	624			
<i>Bsu2c</i>	D 13 D 11	ENR 19	EDKRF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> IL 60	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 26	IN <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 16	S <sub>2</sub> X <sub>2</sub> K 51	SN 120	KTG 175	714			
<i>Syn1a</i>	E 20 E 5	HDK 19	EDSRF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> QILEY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 59	G <sub>2</sub> TTX <sub>2</sub> NX <sub>2</sub> Q 27	MD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> MX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 50	SN 120	KTG 165	885			
<i>Synmrcb</i>	D 70 D 12	IHR 19	EDTRF <sub>2</sub> EHX <sub>2</sub> G 21	GASTITQQ 16	RK <sub>2</sub> E 12	K <sub>2</sub> DILTY <sub>2</sub> X <sub>2</sub> N 56	LR <sub>2</sub> X <sub>2</sub> VI 60	N <sub>2</sub> X <sub>2</sub> TSX <sub>2</sub> NX <sub>2</sub> Q 27	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> S 18	S <sub>2</sub> X <sub>2</sub> K 50	SEN 132	KTG 53	749			
<i>Bhu3</i>	D 15 D 12	ENR 19	EDIGF <sub>2</sub> SHX <sub>2</sub> G 20	GGSTITQQ 15	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 55	IQ <sub>2</sub> VL 67	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 111	ID <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> X 18	S <sub>2</sub> X <sub>2</sub> K 55	SIN 142	KTG 263	932			
<i>Spn1a</i>	E 12 D 11	ERR 19	EDHRF <sub>2</sub> DX <sub>2</sub> G 20	GGSTITQQ 17	RK <sub>2</sub> E 12	K <sub>2</sub> EILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 61	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 26	VD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GAX <sub>2</sub> H 20	S <sub>2</sub> X <sub>2</sub> K 54	SRN 126	KTG 159	719			
<i>Sor1a</i>	E 12 D 11	ERR 19	EDHRF <sub>2</sub> NHX <sub>2</sub> G 21	GGSTITQQ 17	RK <sub>2</sub> E 12	K <sub>2</sub> EILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 61	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 26	VD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GAX <sub>2</sub> H 20	S <sub>2</sub> X <sub>2</sub> K 54	SRN 126	KTG 76	637			
<i>Spy263</i>	E 12 D 11	EKR 19	EDKRF <sub>2</sub> NHX <sub>2</sub> G 20	GGSTITQQ 17	RK <sub>2</sub> E 12	K <sub>2</sub> EILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 61	G <sub>2</sub> TTX <sub>2</sub> X <sub>2</sub> Q 26	VD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> N 20	S <sub>2</sub> X <sub>2</sub> K 54	SRN 126	KTG 158	721			
<i>Bsu1</i>	D 18 D 11	EKR 19	EDARF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 56	RR <sub>2</sub> VL 63	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 24	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> AG <sub>2</sub> N 17	S <sub>2</sub> X <sub>2</sub> K 52	SRN 124	KTG 338	914			
<i>Sau2</i>	E 12 E 12	QRR 19	EDRNF <sub>2</sub> EHX <sub>2</sub> A 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> DIFQY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 73	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 24	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> LX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 52	SRN 126	KTG 130	716			
<i>Spn2a</i>	D 12 D 10	GQK 22	EDSRF <sub>2</sub> KX <sub>2</sub> G 17	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> QILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 64	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 28	LE <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> X <sub>2</sub> G <sub>2</sub> .N 21	S <sub>2</sub> X <sub>2</sub> K 51	SIN 122	KTG 139	731			
<i>Spy250</i>	D 12 D 14	155 178	EDRTF <sub>2</sub> SNX <sub>2</sub> G 17	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> DILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 64	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 29	LD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GR <sub>2</sub> N 21	S <sub>2</sub> X <sub>2</sub> K 50	SIN 122	KTG 138	778			
<i>E. coli</i>	98 D 10	109 D 12	EQR 19	EDRHF <sub>2</sub> EHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> RILEY <sub>2</sub> X <sub>2</sub> N 59	RR <sub>2</sub> VL 57	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 29	VD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> E 17	S <sub>2</sub> X <sub>2</sub> K 58	SN 123	KTG 143	844		
<i>Cfr1b</i>	98 D 10	109 D 12	EQR 19	EDRHF <sub>2</sub> EHX <sub>2</sub> G 21	GASTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> RILEY <sub>2</sub> X <sub>2</sub> N 59	RR <sub>2</sub> VL 57	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 29	VD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> E 17	S <sub>2</sub> X <sub>2</sub> K 58	SN 123	KTG 143	846		
<i>Hin1b</i>	D 105 E 12	EDR 19	EDRRF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> RILEY <sub>2</sub> X <sub>2</sub> N 59	RR <sub>2</sub> VL 58	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 29	TD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> Q 17	S <sub>2</sub> X <sub>2</sub> K 56	SIN 123	KTG 126	781			
<i>Mle1*</i>	D 15 E 8	GNR 19	EDRNF <sub>2</sub> SNX <sub>2</sub> G 21	GGSTITQQ 18	RK <sub>2</sub> E 12	K <sub>2</sub> DVLOQY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 63	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 24	ID <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 14	S <sub>2</sub> X <sub>2</sub> K 49	SIN 138	KTG 133	686			
<i>Mtu1</i>	D 36 E 8	GNR 19	EDRNF <sub>2</sub> SNX <sub>2</sub> G 20	GGSTITQQ 18	RK <sub>2</sub> E 12	K <sub>2</sub> DVLOQY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 63	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 24	TD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> N 14	S <sub>2</sub> X <sub>2</sub> K 49	SIN 138	KTG 127	665			
<i>Mle1</i>	E 10 D 11	QRR 19	EDKRF <sub>2</sub> DX <sub>2</sub> G 21	GGSTITQQ 26	RK <sub>2</sub> E 12	K <sub>2</sub> EILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> VL 65	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 24	I <sub>2</sub> .X <sub>2</sub> .V <sub>2</sub> AX <sub>2</sub> .D 20	S <sub>2</sub> X <sub>2</sub> K 62	SRN 146	KTG 205	821			
<i>Spn1b</i>	D 26 D 10	LLR 19	EDRHF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> EILEY <sub>2</sub> X <sub>2</sub> N 69	RA <sub>2</sub> VL 87	G <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 27	MD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> IX <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> N 17	S <sub>2</sub> X <sub>2</sub> K 55	SN 132	KTG 167	821			
<i>Spy286</i>	E 10 D 10	LLR 19	EDRHF <sub>2</sub> EHX <sub>2</sub> G 21	GGSTITQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> NILCY <sub>2</sub> X <sub>2</sub> N 69	RQ <sub>2</sub> VL 87	G <sub>2</sub> TTX <sub>2</sub> NX <sub>2</sub> Q 27	TD <sub>2</sub> X <sub>2</sub> G <sub>2</sub> V <sub>2</sub> AX <sub>2</sub> GG <sub>2</sub> D 17	S <sub>2</sub> X <sub>2</sub> K 52	SN 132	KTG 144	723			
<b>Identities</b>	* * * * *															
<i>mTgase</i>	47 E 9	57 NFR 84	EDQKF <sub>2</sub> EHX <sub>2</sub> G 21	GASTISQQ 14	RK <sub>2</sub> E 12	K <sub>2</sub> RILTY <sub>2</sub> X <sub>2</sub> N 55	RR <sub>2</sub> IL 17	242	mTgase NH <sub>2</sub>				34 S <sub>2</sub> X <sub>2</sub> K 57	SO <sub>2</sub> 114	KTG 46	262
<i>E. coli</i>	29 D 19	D 12	IWR 19	EDRWF <sub>2</sub> KHX <sub>2</sub> G 21	GGSTITMQ 13	GK <sub>2</sub> Q 12	K <sub>2</sub> EILTY <sub>2</sub> X <sub>2</sub> N 55	AR <sub>2</sub> VL 45	S <sub>2</sub> TTX <sub>2</sub> D <sub>2</sub> X <sub>2</sub> Q 22	I <sub>2</sub> . . . . .X <sub>2</sub> . . . . .D 20	S <sub>2</sub> X <sub>2</sub> K 50	SIN 112	KTG 256	770		

FIG. 6. Amino acid sequence analysis of full-size, multimodular class A PBPs. Conserved motifs, intermodule junction sites, inserts, and amino- and carboxy-terminal extensions are shown. mTgase, putative monofunctional transglycosylase of *E. coli* (Swiss-Prot accession no. P46022). mTgase, monofunctional dd-transpeptidase/PBP of *Streptomyces* strain K15 (Swiss-Prot accession no. P39042); Mle1\*, high-affinity, thermolabile class A PBP of *M. leprae* (43). Asterisks at the bottom of the figure highlight identities defining the amino acid sequence signatures of the modules. Amino acid residues that do not obey the consensus are in underlined italics. For more details, see the text.

tail have more sophisticated functions than that of a simple anchoring device (29, 31). A plausible interpretation of the core-based clustering of the class A PBPs (Fig. 8) is that the gram-negative PBPs of subclasses A1 and A2 are paralogs (i.e., performing different functions); the gram-positive PBPs of subclasses A3, A4, and A5 are also paralogs; and the gram-negative PBPs of subclass A1 or A2 and the gram-positive PBPs of subclass A3, A4, or A5 may be orthologs (i.e., performing similar functions in these two groups of bacteria). Likewise, the clustering of the class B PBPs (Fig. 9) suggest that the gram-negative PBPs of subclasses B2 and B3 are paralogs; the gram-positive PBPs of subclasses B1, B4, and B5 are also paralogs; and the gram-negative PBPs of subclass B2 or B3 and the gram-positive PBPs of subclass B4 or B5 may be orthologs. As discussed below, the gram-positive PBPs of subclass B1 have no equivalents in the gram-negative bacteria. The ensuing sections confront this scheme with facts. The associated n-PB and PB cores comprise about 500 to 550 amino acid residues, whereas the full-size PBPs are polypeptide chains 600 to 900 amino acid residues long. Because the inserts and extensions are not taken into consideration, an

important proportion of the information may be lost in some cases. Moreover, the biochemistry audit of the multimodular PBPs is far short of reality. As stated above, attributing a function to a protein in the absence of direct biochemical data is a dangerous exercise.

**FUNCTIONS OF MULTIMODULAR CLASS A PBPs**

*E. coli* PBP1a of subclass A1 and *E. coli* PBP1b of subclass A2 are the only class A PBPs which have been identified biochemically as bifunctional transglycosylase (n-PB module)-transpeptidase (PB module) entities. However, the class A PBPs have two remarkable features. The n-PB cores have an extended amino acid sequence signature in the form of six conserved motifs (Fig. 6), and they underwent steady divergence without marked adaptive radiation (Fig. 8), suggesting conserved functionality. In contrast, the PB cores, while retaining the penicilloyl serine transferase signature (Fig. 6), have evolved in two gram-negative PBPs of subclasses A1 and A2 that form a supercluster and three gram-positive PBPs of subclasses A3, A4, and A5 that also form a supercluster (Fig.

CLASS B PBPs	n-PB module							PB module							COOH
	1	2	3	4	5	6	7	1	2	3	4	5	6	7	
	$\beta$ 1	$\beta$ 5	loop	$\beta$ 6	$\alpha$ 1	$\beta$ 2	$\alpha$ 2	Loop	$\beta$ 3						
<i>Spn2x</i>															
<i>Efam5</i>	E 134	RGX <sub>169</sub> DRNG	E1 RXYPXG	30 GX <sub>234</sub> GXE <sub>2</sub> D	31 GXDX <sub>335</sub> TDX <sub>3</sub> Q	20 TGD <sub>346</sub> LAX <sub>347</sub> PS <sub>348</sub> DP	34 SX <sub>417</sub> K	54 SDN	134	KTG	58	673			
<i>Ehi3r</i>	E 134	RGX <sub>173</sub> DRNG	E1 RXYPXG	30 GX <sub>239</sub> GXE <sub>2</sub> D	31 GXDX <sub>340</sub> TDX <sub>3</sub> Q	20 TGD <sub>341</sub> LAX <sub>342</sub> PS <sub>343</sub> DP	34 SX <sub>410</sub> K	54 SDN	134	KTG	58	678			
<i>Ehi5</i>	E 134	RGX <sub>173</sub> DRNG	E1 RXYPXG	30 GX <sub>239</sub> GXE <sub>2</sub> D	31 GXDX <sub>340</sub> TDX <sub>3</sub> Q	20 TGE <sub>373</sub> LXV <sub>374</sub> PS <sub>375</sub> DP	34 SX <sub>422</sub> K	54 SDN	134	KTG	59	678			
<i>Efas5</i>	E 131	RGX <sub>159</sub> DRNG	E1 RXYPXG	29 GX <sub>234</sub> GXE <sub>2</sub> D	31 GX <sub>341</sub> TX <sub>342</sub> TDX <sub>3</sub> Q	20 KGD <sub>343</sub> LAX <sub>344</sub> PS <sub>345</sub> DF	34 SX <sub>423</sub> K	54 SDN	134	KTG	58	679			
<i>Bsu3</i>	D 133	RGX <sub>151</sub> DKNG	E5 RXYPXG	31 GX <sub>241</sub> GXE <sub>2</sub> D	27 GXDX <sub>321</sub> TDX <sub>3</sub> Q	20 TGE <sub>354</sub> LAX <sub>355</sub> PS <sub>356</sub> DP	34 SX <sub>403</sub> K	55 SDN	136	KTG	57	668			
<i>Sau2a</i>	D 123	RGX <sub>150</sub> DRNN	E1 RXYPXG	32 GX <sub>218</sub> GXE <sub>2</sub> D	32 GXDX <sub>319</sub> TDX <sub>3</sub> Q	20 TGE <sub>352</sub> LAX <sub>353</sub> PS <sub>354</sub> DV	34 SX <sub>401</sub> K	55 SDN	132	KSG	68	668			
<i>Ssc2</i>	E 118	RGX <sub>68</sub> DRNN	E1 RXYPXG	32 GX <sub>206</sub> GXE <sub>2</sub> D	31 GXDX <sub>247</sub> TDX <sub>3</sub> Q	20 TGE <sub>289</sub> LAX <sub>290</sub> PS <sub>291</sub> DV	34 SX <sub>337</sub> K	55 SDN	132	KSG	68	666			
<i>Eco2</i>	D 18	RGX <sub>172</sub> DRNG	E7 RXYPXG	36 GX <sub>214</sub> GXE <sub>2</sub> E	31 GXDX <sub>255</sub> TDX <sub>3</sub> Q	20 TGG <sub>288</sub> LAX <sub>289</sub> PS <sub>290</sub> DP	35 SX <sub>338</sub> K	53 SAD	154	KSG	86	633			
<i>Hin2</i>	D 19	RGX <sub>76</sub> DRYG	E7 RXYLXG	36 GX <sub>215</sub> GXE <sub>2</sub> E	31 GX <sub>255</sub> SX <sub>256</sub> TDX <sub>3</sub> Q	20 D <sub>259</sub> X <sub>260</sub> LAX <sub>261</sub> PS <sub>262</sub> DN	35 SX <sub>339</sub> K	53 SSD	154	KSG	96	651			
<i>AaeA1</i>	E 21	RGX <sub>53</sub> DRNG	E3 RXYPXG	45 GX <sub>187</sub> GXE <sub>2</sub> V	17 GX <sub>228</sub> SX <sub>229</sub> TDX <sub>3</sub> Q	23 TGE <sub>311</sub> LAX <sub>312</sub> PN <sub>313</sub> NP	30 SX <sub>366</sub> K	51 SC <sub>367</sub> D	151	KTG	79	595			
<i>Hpy2</i>	E 19	RGX <sub>53</sub> DRND	E9 RXYPXN	30 GX <sub>187</sub> GXE <sub>2</sub> N	31 N <sub>211</sub> X <sub>212</sub> SX <sub>213</sub> DX <sub>3</sub> Q	20 NGE <sub>261</sub> LXV <sub>262</sub> PE <sub>263</sub> NL	35 SX <sub>311</sub> K	51 SVD	144	KTG	62	588			
<i>Ebu2</i>	D 16	RGX <sub>54</sub> DSNS	E9 RXYLXD	33 GX <sub>145</sub> GXE <sub>2</sub> D	32 GX <sub>223</sub> DX <sub>224</sub> PNX <sub>225</sub> Q	20 TGA <sub>259</sub> LAX <sub>260</sub> PYX <sub>261</sub> SM	26 SX <sub>304</sub> K	51 SSN	156	KTG	73	599			
<i>ScoA3</i>	E 18	RGX <sub>69</sub> D-RD	E9 RXYAXF	36 GX <sub>215</sub> GXE <sub>2</sub> D	33 GX <sub>256</sub> DX <sub>257</sub> SX <sub>258</sub> DX <sub>3</sub> Q	39 TGR <sub>303</sub> LXV <sub>304</sub> PD <sub>305</sub> XXX	36 SX <sub>359</sub> K	50 SC <sub>360</sub> D	197	KTG	154	588			
<i>Scipcbr</i>	D 128	RGX <sub>71</sub> DRANG	..... 18	..... GX <sub>167</sub> GX <sub>168</sub> QX <sub>169</sub> D	34 G <sub>219</sub> X <sub>220</sub> TDX <sub>3</sub> Q	22 TGN <sub>269</sub> LAX <sub>270</sub> PS <sub>271</sub> XXX	10 SX <sub>307</sub> K	51 SC <sub>308</sub> N	129	KTG	55	551			
<i>Eco3</i>	D 19	RGX <sub>91</sub> DRSG	E7 RXYPXG	15 GX <sub>208</sub> GXE <sub>2</sub> D	31 AX <sub>249</sub> DX <sub>250</sub> SX <sub>251</sub> DX <sub>3</sub> Q	27 TGE <sub>289</sub> LAX <sub>290</sub> PS <sub>291</sub> XNP	23 SX <sub>327</sub> K	48 SSN	132	KTG	91	588			
<i>Hin3</i>	D 19	RGX <sub>62</sub> DRNG	E7 RXYPXV	15 GX <sub>153</sub> GXE <sub>2</sub> N	31 AX <sub>216</sub> DX <sub>217</sub> SX <sub>218</sub> DX <sub>3</sub> Q	27 TGE <sub>256</sub> LAX <sub>257</sub> PS <sub>258</sub> XNP	23 SX <sub>294</sub> K	48 SSN	130	KTG	95	610			
<i>Pae3</i>	D 21	RGX <sub>55</sub> DRNG	E2 RXYPXG	15 GX <sub>148</sub> GXE <sub>2</sub> D	32 G <sub>211</sub> X <sub>212</sub> SX <sub>213</sub> DX <sub>3</sub> Q	27 TGE <sub>251</sub> LAX <sub>252</sub> PTX <sub>253</sub> NP	23 SX <sub>294</sub> K	51 SSN	132	KSG	92	579			
<i>Pae3a</i>	D 21	RGX <sub>55</sub> DRNG	E2 RXYPXG	15 GX <sub>148</sub> GXE <sub>2</sub> D	32 G <sub>211</sub> X <sub>212</sub> SX <sub>213</sub> DX <sub>3</sub> Q	27 TGE <sub>251</sub> LAX <sub>252</sub> PS <sub>253</sub> XNP	23 SX <sub>294</sub> K	50 SSN	132	KSG	84	565			
<i>Syn3</i>	E 21	RX <sub>75</sub> DSQG	E8 RXYPXG	15 GX <sub>188</sub> GXE <sub>2</sub> R	33 DX <sub>232</sub> QX <sub>233</sub> TDX <sub>3</sub> Q	27 DGA <sub>272</sub> LAX <sub>273</sub> PTX <sub>274</sub> NP	23 SX <sub>310</sub> K	51 SSN	145	KTG	73	607			
<i>Nme2</i>	E 15	RGX <sub>75</sub> DRNG	E3 RXYPXG	15 GX <sub>188</sub> GXE <sub>2</sub> E	34 GX <sub>232</sub> DX <sub>233</sub> SX <sub>234</sub> DX <sub>3</sub> Q	27 TGE <sub>272</sub> LAX <sub>273</sub> PA <sub>274</sub> MEP	23 SX <sub>310</sub> K	49 SSN	132	KTG	83	584			
<i>Ngo2</i>	E 15	RGX <sub>63</sub> DRNG	E3 RXYPXG	15 GX <sub>188</sub> GXE <sub>2</sub> E	34 GX <sub>232</sub> DX <sub>233</sub> SX <sub>234</sub> DX <sub>3</sub> Q	27 TGE <sub>272</sub> LAX <sub>273</sub> PA <sub>274</sub> MDP	23 SX <sub>310</sub> K	49 SSN	132	KTG	81	582			
<i>AaeA2</i>	E 18	RGX <sub>76</sub> TSDG	E10 RXYPXG	102 AX <sub>215</sub> GXE <sub>2</sub> D	32 AX <sub>258</sub> DX <sub>259</sub> SX <sub>260</sub> DX <sub>3</sub> Q	27 TGA <sub>299</sub> LAX <sub>300</sub> PD <sub>301</sub> DP	23 SX <sub>337</sub> K	54 SSN	127	KTG	72	578			
<i>Spn2x</i>	D 18	RGX <sub>57</sub> DRNG	E10 RXYPXG	23 GX <sub>215</sub> GXE <sub>2</sub> N	33 GX <sub>258</sub> DX <sub>259</sub> TX <sub>260</sub> X <sub>3</sub> Q	27 TGE <sub>299</sub> LAX <sub>300</sub> PT <sub>301</sub> MDA	24 SX <sub>337</sub> K	54 SSN	149	KSG	200	750			
<i>Spy290</i>	D 24	RGX <sub>76</sub> DRNG	E10 RXYPXG	24 GX <sub>192</sub> GXE <sub>2</sub> N	33 GX <sub>264</sub> DX <sub>265</sub> TX <sub>266</sub> X <sub>3</sub> Q	27 TGE <sub>304</sub> LAX <sub>305</sub> PT <sub>306</sub> NNA	25 SX <sub>343</sub> K	55 SSN	148	KSG	198	751			
<i>EfamC</i>	E 15	RGX <sub>60</sub> DRYG	E10 RXYPXG	23 GX <sub>221</sub> GXE <sub>2</sub> N	33 GX <sub>264</sub> DX <sub>265</sub> TDX <sub>3</sub> Q	27 TGE <sub>304</sub> LAX <sub>305</sub> PT <sub>306</sub> XNP	24 SX <sub>343</sub> K	54 SSN	143	KTG	193	742			
<i>Ehi3s</i>	E 15	RGX <sub>60</sub> DRNG	E10 RXYPXG	20 GX <sub>218</sub> GXE <sub>2</sub> N	33 GX <sub>264</sub> DX <sub>265</sub> TDX <sub>3</sub> Q	27 TGE <sub>304</sub> LAX <sub>305</sub> PT <sub>306</sub> NP	25 SX <sub>343</sub> K	50 SSN	142	KTG	191	730			
<i>Bsu2b</i>	E 19	RGX <sub>65</sub> DRKG	E9 RXYPXG	20 GX <sub>194</sub> GXE <sub>2</sub> D	33 GX <sub>237</sub> DX <sub>238</sub> TDX <sub>3</sub> Q	27 TGE <sub>277</sub> LAX <sub>278</sub> PS <sub>279</sub> DP	20 SX <sub>314</sub> K	51 SSN	143	KTG	200	716			
<i>SauB</i>	D 17	RGX <sub>59</sub> DRNG	E5 RXYPXG	20 GX <sub>151</sub> GXE <sub>2</sub> D	32 GX <sub>215</sub> DX <sub>216</sub> TDX <sub>3</sub> Q	27 TGE <sub>255</sub> LAX <sub>256</sub> PT <sub>257</sub> XNP	22 SX <sub>294</sub> K	50 SSN	142	KTG	130	646			
<i>BsuVD</i>	D 31	RGX <sub>58</sub> DRNG	E3 RXYPXG	15 GX <sub>167</sub> GXE <sub>2</sub> D	33 GX <sub>215</sub> DX <sub>216</sub> TDX <sub>3</sub> Q	27 NGE <sub>255</sub> LGX <sub>256</sub> PD <sub>257</sub> DP	24 SX <sub>294</sub> K	52 SCN	143	KTG	146	645			
<i>Ebu1</i>	D 28	RGX <sub>90</sub> DRNG	E4 RXYPXG	15 GX <sub>177</sub> GXE <sub>2</sub> N	18 TX <sub>245</sub> DX <sub>246</sub> TDX <sub>3</sub> Q	27 NGE <sub>236</sub> LXV <sub>237</sub> PQ <sub>238</sub> MDA	23 SX <sub>274</sub> K	55 SSN	142	KSG	148	629			
<i>Hpy3</i>	D 12	RFX <sub>61</sub> DKED	E7 RXYPXG	22 GX <sub>205</sub> GXE <sub>2</sub> D	34 GX <sub>249</sub> EX <sub>250</sub> SX <sub>251</sub> DX <sub>3</sub> Q	27 SGE <sub>282</sub> L <sub>283</sub> X <sub>284</sub> NP	23 SX <sub>327</sub> K	51 SSN	142	KTG	85	615			
<i>Spn2b</i>	D 19	RGX <sub>60</sub> DASG	..... 194	..... G <sub>263</sub> X <sub>264</sub> EX <sub>265</sub> E	31 GX <sub>304</sub> DX <sub>305</sub> TDX <sub>3</sub> Q	..... 68	..... SX <sub>365</sub> K	53 SSN	169	KTG	62	679			
<i>Stb2b</i>	E 12	RGX <sub>69</sub> DAKG	..... 195	..... G <sub>273</sub> X <sub>274</sub> EX <sub>275</sub> E	32 GX <sub>314</sub> DX <sub>315</sub> TDX <sub>3</sub> Q	..... 68	..... SX <sub>377</sub> K	52 SSN	168	KTG	65	704			
<i>Bsu2a</i>	D 29	RGX <sub>69</sub> DRNF	..... 195	..... G <sub>273</sub> X <sub>274</sub> EX <sub>275</sub> E	31 GX <sub>314</sub> DX <sub>315</sub> TDX <sub>3</sub> Q	..... 70	..... SX <sub>377</sub> K	50 SSN	178	KTG	97	716			
<b>Identities</b>		** *	** *	** *	** *	** *	** *	** *	** *	** *	** *	** *	** *	** *	

FIG. 7. Amino acid sequence analysis of full-size, multimodular class B PBPs. Conserved motifs, intermodule junction sites, inserts, and carboxy-terminal extensions are shown. Secondary structures of the *S. pneumoniae* PBP2x are shown at the top. Identities (asterisks) define the amino acid sequence signatures of the modules. Amino acid residues that do not obey the consensus are in underlined italics. For more details, see the text.

8), suggesting differential functionality. On the basis of this conjecture, plausible suggestions can be made.

Dicarboxylic amino acid residues E and D of motif 1 and E of motif 3 present in all class A PBPs (Fig. 6) are almost certainly important components of the transglycosylase catalytic center of the n-PB modules. In this respect, *E. coli* PBP1c looks like a class A PBP mutant. The changes R→G and E→Q in motif 3 and R→A in motif 5 suggest that the n-PB module of this PBP may not be a transglycosylase.

Peptidoglycan cross-linking by the acyl serine transferase-PB module of the class A PBPs is more complex than expected. PBP1a of subclass A1 and PBP1b of subclass A2 substitute for each other in *E. coli* (37). Loss of either PBP1a or PBP1b is tolerated, but loss of the two class A PBPs is fatal. Under these extreme conditions, PBP1c cannot rescue *E. coli* from death.

PBP1 of subclass A3, PBP2c of subclass A4, and PBP4 of subclass A5 also substitute for each other in *B. subtilis* (59). Loss of PBP1 is associated with slight growth and cell morphology defects, these defects are exacerbated by the additional loss of PBP4 but not that of PBP2c, and loss of all three PBPs slows growth even further but is not fatal. Under these extreme conditions (and as discussed in the next section), *B. subtilis* PBP3 of subclass B1 might take over the functions of the PBPs of subclasses A3, A4, and A5.

Although performing the same transglycosylase-transpeptidase activities, PBP1a and PBP1b in *E. coli* and, likewise, PBP1, PBP2c, and PBP4 in *B. subtilis* are probably paralogs, prescribing distinct, subtle traits in peptidoglycan cross-linking in these two organisms that are difficult to detect.

**FUNCTIONS OF MULTIMODULAR CLASS B PBPs**

The class B PBPs evolved differently (Fig. 9). The n-PB cores and PB cores underwent adaptive radiation in such a concerted manner that in the gram-negative bacteria an n-PB core of subclass B2 or B3 is linked, almost invariably, to a PB core of subclass B2 or B3, respectively, and in the gram-positive bacteria an n-PB core of subclass B1, B4, or B5 is linked to a PB core of subclass B1, B4, or B5, respectively.

**Class B PBPs of Gram-Negative Bacteria**

*E. coli* PBP2 of subclass B2 and *E. coli* PBP3 of subclass B3 cannot substitute for each other. They are paralogs. *E. coli* PBP3 of subclass B3 and a set of non-PBPs work together to initiate septum formation (53, 67). In recent years, the catalogue of these non-PBPs has grown considerably. The genes encoding PBP3 and six other cell cycle proteins reside in the




FIG. 8. Hierarchical analysis of the n-PB cores and PB cores of the multimodular class A PBPs. PBP codes are given in Table 1.

division and cell wall (*dcw*) cluster at the 2-min region of the chromosome (67). PBP3, FtsQ (10), and FtsL, a protein bearing a putative leucine zipper motif (30), are membrane bound, with the bulk of the polypeptide chains exposed on the outer face of the plasma membrane. FtsW is an integral membrane protein with loops exposed on both faces of the membrane (6). FtsA, an isolog of the DnaK-actin family of ATPases, is cytosolic when phosphorylated and is membrane associated when

unphosphorylated (63). MraW, a protein bearing a putative *S*-adenosylmethionine-binding motif (9), and FtsZ, a GTPase similar to tubulin (47), are cytosolic. FtsZ is of known three-dimensional structure (46, 54). It functions as a cytoskeletal element mediating the invagination of the septum.

*E. coli* PBP2 of subclass B2, RodA (an integral membrane protein similar to FtsW), and the monofunctional DD-carboxypeptidase PBP5, whose genes are located at the 14-min




FIG. 9. Hierarchical analysis of the n-PB cores and PB cores of the multimodular class B PBPs. PBP codes are given in Table 1.

TABLE 2. Pairwise comparison of the amino acid sequences of inserts of multimodular PBPs

Subclass	PBP <sup>a</sup>	Similarity (%)
A2	Eco1b D88-D198/Cfr1b D88-D198	91
	/Hin1b D44-E150	32
B1	Efam5 E33-R168/Ehi3r E38-R173	97
	/Ehi5 E38-R173	65
	/Efas5 E43-R175	35
	/Bsu3 D25-R159	27
	Ehi5 E38-R173/Efas5 E43-R175	37
	/Bsu3 D25-R159	28
	Efas5 E43-R175/Bsu3 D25-R159	23
	Sau2a D27-R151/Ssc2 E31-R150	71
	/Efam5 E33-R168	17
	/Ehi3r E38-R173	17
/Ehi5 E38-R173	17	
/Efas5 E43-R175	17	
/Bsu3 D25-R159	18	
A1	Eco1a N526-K716/Hin1a N501-K717	48
	/Pae1a N522-K695	31
	Hin1a N501-K717/Pae1a N522-K695	31
A1	Nme1/Ngo1 Q298-L428/Eco1a Q296-I432	25
	/Hin1a Q286-L408	26
	/Pae1a Q296-L428	26
	/Aaemrca Q291-I399	21
	/Bbu3 Q316-I428	6
	Eco1a Q296-I432/Hin1a Q286-L408	45
	/Pae1a Q296-L428	40
	/Aaemrca Q291-I399	16
	/Bbu3 Q316-I428	4
	Hin1a Q286-L408/Pae1a Q296-L428	32
	/Aaemrca Q291-I399	14
	/Bbu3 Q316-I428	9
	Pae1a Q296-L428/Aaemrca Q291-I399	18
	/Bbu3 Q316-I428	10
	Aaemrca Q291-I399/Bbu3 Q316-I428	15

<sup>a</sup> The PBP codes are given in Table 1. PBPs Nme1 and Ngo1 have 98% identity.

<sup>b</sup> Similarity is expressed as percent identity.

region of the chromosome, are involved in wall expansion and shape maintenance (35, 61). PBP2, RodA, and ribosomal activities are coordinated by a chain of interacting elements, one of which is regulated by the nucleotide guanosine 5'-diphosphate 3'-diphosphate ppGpp.

*E. coli* PBP2 is not required for septum synthesis. Loss of PBP2 results in a block of cell division; however, in the absence of PBP2, cell division and viability are restored by increasing the pool of ppGpp or the level of FtsQAZ, showing that the cell septation and cell shape networks are interconnected (35). A fascinating story is now beginning to unfold involving the description of these networks in terms of interaction between the constitutive components. In spite of these advances, the questions of how these proteins work together and how the bacterial cell choreographs the interplay remain before us.

### Class B PBPs of Gram-Positive Bacteria

The gram-positive bacteria possess two class B PBPs of subclasses B4 and B5, and some of them possess one additional PBP of subclass B1 (Fig. 9). The gram-positive PBPs of subclass B4 and the gram-negative PBPs of subclass B3 are almost certainly orthologs involved in cell division. Inactivation of *Enterococcus hirae* PBP3s of subclass B4 results in a block of septum formation (11), and a 10-kb segment of the *E. hirae* chromosome that contains the PBP3s-encoding gene also contains genes that code for proteins similar to MraW, FtsL, FtsQ, FtsW, FtsA, and FtsZ (18), a situation comparable to that of the *E. coli* *dcw* cluster. *B. subtilis* has a *dcw* cluster with a similar organization (7). However, this *dcw* cluster contains two PBP-encoding genes. PBP2b, which is involved in the metabolism of the vegetative cells, belongs to the gram-positive PBPs of subclass B4. PBPVD, which is involved in sporulation, belongs to the gram-negative PBPs of subclass B3.

The gram-positive PBPs of subclass B5 are of unknown function, but they are important. In *Streptococcus pneumoniae*, PBP2b of subclass B5 and PBP2x of subclass B4 cannot substitute for each other (40). They are paralogs. PBP2b and PBP2x are the first PBPs to be affected during selection of laboratory mutants with a reduced affinity for cefotaxime and piperacillin, respectively, and the modified PBP2b and PBP2x each confer resistance upon transformation (32).

The gram-positive PBPs of subclass B1 are endowed with unique properties. They represent an important mechanism of resistance to penicillin. The enterococcal and staphylococcal PBPs of subclass B1 have a very low affinity for the drug. They allow the strains that (over)produce them to grow in the presence of penicillin at concentrations sufficient to inactivate the PB modules of all the other multimodular PBPs of class A and class B (4, 19, 58, 60). To all appearances, the low-affinity PBPs of subclass B1 (including, perhaps, *B. subtilis* PBP3) can perform the basic functions required for wall peptidoglycan assembly in a cell-cycle-dependent fashion in conjunction, presumably, with the transglycosylase n-PB module of class A PBPs or with monofunctional transglycosylases (64).

### Class B PBPs in Cell Morphogenesis

The question of how the class B PBPs function in vivo is left open. However, the unprecedented amino acid sequence signature of their n-PB modules (Fig. 7) and the three-dimensional structure of *S. pneumoniae* PBP2x of subclass B4 (57) are worthy of reflection.

The n-PB module of PBP2x is shaped like a pair of sugar tongs whose head fits in a noncatalytic groove of the PB module. Remarkably, motifs 1 to 4 of the n-PB module are located in the head of the "sugar tongs" in interaction with the PB module, and the 101-amino-acid residue polypeptide stretch that extends between motifs 1 and 2 is well exposed at the surface of the protein. Other class B PBPs probably adopt the same basic folded structure but with subclass- and species-specific variations. Key elements of the amino acid sequence-folding information for *E. coli* PBP3 of subclass B3 reside in the sequence G57 to W110, which contains motif 1, and in motif 3. Alterations of E193 of motif 3 results in the production of misfolded, unstable protein mutants that are rapidly degraded (29).

*S. pneumoniae* PBP2x and *E. coli* PBP3 each catalyze peptide bond formation from properly structured thiolester carbonyl donors and amino acceptors (1). Based on this observation, the acyl serine transferase-PB module of the class B PBPs probably prescribes species-specific traits related to peptidoglycan cross-linking, and this activity might be regulated by

the associated n-PB module itself in interaction with other components of the morphogenetic networks. These speculations illustrate how little we know of the biochemistry of the class B PBPs.

### AN AMAZING PANOPLY OF MULTIMODULAR PBPs

As shown in Table 1, *B. subtilis* possesses one PBP each in subclasses A3, A4, and A5 and one PBP each in subclasses B1, B3, B4, and B5 (one of them is involved in sporulation). *S. pneumoniae* possesses one PBP each in subclasses A3, A4, and A5 and one PBP each in subclasses B4 and B5. *E. coli* and *Haemophilus influenzae* possess one PBP each in subclasses A1 and A2 (not counting *E. coli* PBP1c) and one PBP each in subclasses B2 and B3. *A. aeolicus* possesses one PBP of subclass A1 and one PBP each in subclasses B2 and B3. *Synechocystis* strain PCC6803 possesses three PBPs of subclass A1 and one PBP of subclass B3. *Helicobacter pylori* possesses one PBP of subclass A2 and two mixed class B PBPs whose PB cores belong to subclasses B2 and B3, respectively.

The differentiation of multimodular PBPs into classes A and B is almost certainly an ancient evolutionary event. The question of which properties of the wall peptidoglycan assembly machinery determine the variable, species-specific assortment of multimodular PBPs is left open.

### PENICILLIN-ORIENTED EVOLUTION

Diverging evolution gave rise to many different groups, classes, and subclasses of penicilloyl serine transferases. The use of  $\beta$ -lactam antibiotics functions to fuel the emergence of  $\beta$ -lactamases with widely varying spectra against  $\beta$ -lactam antibiotics. In recent years, more than 50 variants of the class A TEM  $\beta$ -lactamase (8) and many variants of the class A SHV  $\beta$ -lactamase have been identified in clinical isolates. They each arose by alteration of a limited number of amino acid residues in the corresponding wild-type enzymes. Another result of this directional evolution is the emergence and the spread of multimodular PBPs with a low or decreased affinity for the drug among important bacterial pathogens.

Enterococci, although related to streptococci, are 10- to 100-fold more resistant to penicillin because they produce one or two low-affinity PBPs of subclass B1. *Enterococcus faecium* is the most resistant species, and new populations of clinical isolates have appeared in different countries (44, 74). The gene encoding the low-affinity PBP5 is chromosomal in *E. faecium* and *E. hirae* R40. In contrast, *E. hirae* S185R (a derivative of a swine isolate) possesses a large plasmid that carries several copies of the gene encoding the low-affinity PBP3r (preceded by a *psr*-like negative regulatory gene), several IS1216 insertion sequences, and the determinants of resistance to streptomycin and erythromycin (60). These elements form a transposon-like structure, a situation eminently favorable for the horizontal spread of resistance to these antibiotics. Because *pbp3r* and *psr3r* are on a 1.3-kb DNA segment whose sequence is 99% homologous to that present in the *E. faecium* D63R chromosome, a DNA fragment might have been excised from *E. faecium* and inserted in a plasmid of *E. hirae* or in a plasmid of *E. faecium* which was then transferred to *E. hirae*.

In staphylococci, acquired resistance is also caused by the acquisition of a low-affinity PBP of subclass B1, known as PBP2a or PBP2'. The encoding *mecA* is chromosomal, and its environment is somewhat similar to that of the low-affinity PBP-encoding genes in streptococci (4). IS257, which is similar to IS1216, is probably involved in the integration of *mecA* in the chromosome of MRSA (methicillin-resistant *S. aureus*)

strains, in the amplification phenomena observed in highly resistant staphylococcal mutants, in the evolution of the *mec* locus, and, perhaps, in the clustering of additional antibiotic resistance determinants. Strains close to the squirrel *S. sciuri* might be the sources of *mecA* in human staphylococcal pathogens (72).

In *Streptococcus pneumoniae*, *N. meningitidis*, and *N. gonorrhoeae*, acquired resistance is caused by the acquisition of altered forms of wild-type class A and class B PBPs. These PBPs of decreased affinity for the drug are the products of mosaic genes in which sensitive sequence blocks are replaced by homologous, resistant sequence blocks from related species by recombinational events (17). Of particular concern is this type of resistance in pneumococci, in which it is increasing worldwide. Mosaic structures have been described for the *S. pneumoniae* genes that encode PBP1a, PBP2a, and PBP1b of subclasses A3, A4, and A5, PBP2x of subclass B4, and PBP2b of subclass B5 (33). More than 20 variants of PBP2x have been identified in clinical isolates (32). Resistant blocks diverge from the sensitive ones by about 20% in nucleotides, and the mosaic PBPs diverge from the sensitive ones by about 15% in amino acid residues. The origin of the foreign DNA sequences is unknown, but two to four different sources may be involved (32).

### TUNING UP THE CATALYTIC TWO-STROKE ENGINE

Diverging evolution, penicillin-oriented evolution, and site-directed mutagenesis (which allows protein mutants derived from extant genes to be produced through a mode unexplored, or explored but not retained, by natural selection) each show that the ability of the serine  $\beta$ -lactamases and the PB modules of the multimodular PBPs to adapt their catalytic properties to new situations is limitless. As a corollary, the ability of the proton abstraction-donation machinery of the acyl serine transferase catalytic center (Fig. 2A) to acquire new properties in response to structural changes in the enzyme polypeptide backbone is also limitless.

Michaelis complexes are formed by the noncovalent binding of carbonyl donors to the enzyme catalytic center. Dense hydrogen-bonding networks, comparable to three-dimensional cobwebs, connect the side chains and functional groupings of the ligand, the side chains of amino acid residues of the active-site-defining motifs, and several water molecules, one of which is hydrogen bonded to the  $\gamma$ OH of the essential serine residue of motif Sx<sub>2</sub>K.

Formation of the Michaelis complex launches catalysis. At this level of the investigation (the angstrom level), the methods of quantum chemistry only highlight the underlying mechanism. As derived from theoretical studies first carried out on chymotrypsin (14–16), the simplest model of the engine that performs the required proton shuttle for catalysis is a six-membered ring in which a water molecule bridges the hydrogen atom of the serine  $\gamma$ OH and the X atom of the scissile CO—X bond of the carbonyl donor (Fig. 2B).

Assuming that the engine which is created upon binding of the ligand to the catalytic center is in perfect tune, two backbone NH groups polarize the carbonyl of the CO—X bond, the proton of the serine  $\gamma$ OH is transmitted to the X atom via the water molecule, and the serine residue is acylated (step 1). In turn, the serine-ester-linked acyl enzyme adopts the required conformation for the creation of a six-membered ring which, upon entry of the acceptor HY molecule, achieves enzyme deacylation (step 2). In the penicilloyl serine transferases, the backbone NH groups belong to the active-site serine itself and to the amino acid residue immediately downstream from motif [K/H][T/S]G on strand  $\beta$ 3.

Admittedly, the efficacy with which the two-stroke engine overcomes the free energy barrier of the reactions via the transition states depends on many parameters. These parameters include the freedom of the catalytic water molecule itself, the ease with which the donor and the acceptor molecules can undergo deformation, and the ease with which the enzyme polypeptide backbone can undergo relaxation. Moreover, the six-membered ring model is an oversimplification. Additional water molecules, the imidazole moiety of the histidine residue of the catalytic triad of chymotrypsin, the  $\epsilon$  amino group of the conserved lysine residue of motif S<sub>x</sub>K of the penicilloyl serine transferases, the  $\gamma$  carboxylate of the glutamic acid residue of motif Ex<sub>2</sub>LN of class A  $\beta$ -lactamases, and side chains of other amino acid residues at the boundary of the catalytic center shape and orient the two-stroke engine and/or are involved in proton transmission.

To understand the effects of these subtle changes, potential energy hypersurfaces (the dimensionality of which is  $3N - 6$ , where  $N$  is the number of atoms of the system) can be explored. They describe faithfully the geometric rearrangements and the electronic redistributions that the interacting partners, the enzyme catalytic center and the ligands, undergo along the reaction pathway (28). Minute structural changes in the enzyme and/or the ligand with which it is reacting may alter the entire hydrogen-bonding conformation of the Michaelis complex. As a corollary, the characteristics of the potential energy hypersurfaces and the reaction pathways are both ligand and enzyme specific.

One conclusion of these studies is that mutations that affect the polypeptide chain of a  $\beta$ -lactamase or a multimodular PBP may result in the creation of a two-stroke engine of increased catalyzed hydrolysis of a  $\beta$ -lactam antibiotic by the  $\beta$ -lactamase or of decreased affinity of the PBP for the drug (without alteration of its physiological function). In the context of penicillin-oriented evolution, these mutations are selected and the encoded  $\beta$ -lactamase and PBP mutants are maintained. There are no signs indicating that the resistant bacteria which produce these protein mutants suffer from decreased fitness and could not compete with the sensitive ones in a penicillin-free world. The absence of penicillin is not a selective pressure. Resistance is probably a road of no return (52, 62).

## CONCLUSIONS AND FUTURE DIRECTIONS

The simplest conceivable event at the level of the gene can result in the emergence of penicillin (and other antibiotics) resistance determinants. The bacterial world behaves as an enormous organism whose cells exchange their genes with great ease, and the opportunities for the exchange of genetic material in nature are considerable. The antibiotics are societal drugs: a resistance gene which appears somewhere can spread far and fast. The use of current antibiotics in ways that would prevent the worldwide prevalence of resistant bacterial strains is a preemptory necessity.

Direct approaches to the drug resistance problem would be to design new antibiotics that bacteria have never seen before. One approach rests upon the knowledge of the structure and functioning of old targets at the most fundamental level. Multimodular PBPs which are assigned to different functional classes and subclasses and enzymes involved in the synthesis of lipid II are in front of the stage. Another approach rests upon the identification of new targets. The non-PB cell cycle proteins of the morphogenetic networks are good places to look. The ultimate goal, however, will be never reached, due to the evolutionary characteristics of the biological systems.

## ACKNOWLEDGMENTS

This work was supported in part by the Belgian programme on Interuniversity Poles of Attraction initiated by the Belgian State, Prime Minister's Office, Services fédéraux des affaires scientifiques, techniques et culturelles (PAI no. P4/03) and the Fonds de la Recherche Fondamentale Collective (contract 2.4534.95). C.G. is Chercheur qualifié of the Fonds National de la Recherche Scientifique, Brussels, Belgium.

We thank Martine Nguyen-Distèche, Claudine Fraipont, Jacques Coyette, and Georges Dive, Centre d'Ingénierie des Protéines, Regine Hakenbeck (Kaiserslautern University, Germany), Jean van Heijenoort (CNRS, Orsay, France), and two anonymous referees for their authoritative comments and suggestions.

## REFERENCES

- Adam, M., C. Fraipont, N. Rhazi, M. Nguyen-Distèche, B. Lakaye, J. M. Frère, B. Devreese, J. Van Beeumen, Y. van Heijenoort, J. van Heijenoort, and J. M. Ghuyesen. 1997. The bimodular G57-V577 polypeptide chain of the class B penicillin-binding protein 3 of *Escherichia coli* catalyses peptide bond formation from thioesters and does not catalyze glycan chain polymerization from lipid II intermediate. *J. Bacteriol.* **179**:6005-6009.
- Barton, G. J. 1990. Protein multiple sequence alignment and flexible pattern matching. *Methods Enzymol.* **183**:403-428.
- Basu, J., S. Mahapatra, M. Kundu, S. Mukhopadhyay, M. Nguyen-Distèche, P. Dubois, B. Joris, J. Van Beeumen, S. T. Cole, P. Chakrabarti, and J. M. Ghuyesen. 1996. Identification and overexpression in *Escherichia coli* of a *Mycobacterium leprae* gene, *ponI*, encoding high-molecular-mass class A penicillin-binding protein PBP1. *J. Bacteriol.* **178**:1707-1711.
- Berger-Bächi, B. 1994. Expression of resistance to methicillin. *Trends Microbiol.* **2**:389-393.
- Blattner, F. R., et al. 1997. The complete genome sequence of *Escherichia coli* K12. *Science* **277**:1453-1462.
- Boyle, D. S., M. M. Khattar, S. G. Addinall, J. Lutkenhaus, and W. D. Donachie. 1997. *ftsW* is an essential cell-division gene in *Escherichia coli*. *Mol. Microbiol.* **24**:1263-1273.
- Buchanan, C. E., A. O. Henriques, and P. S. Piggot. 1994. Cell wall changes during bacterial endospore formation, p. 167-183. *In* J. M. Ghuyesen and R. Hakenbeck (ed.), *Bacterial cell wall*. Elsevier Science B.V., Amsterdam, The Netherlands.
- Bush, K., and G. Jacoby. 1997. Nomenclature of TEM  $\beta$ -lactamases. *J. Antimicrob. Chemother.* **39**:1-3.
- Carrion, M., M. J. Gómez, and J. A. Ayala. 1995. Molecular analysis of the gene *mrwW* at the *dcw* cluster of *Escherichia coli*. Workshop on Structure, Function and Controls in Microbial Division. Ist. Juan March Estud. Investig. **42**:17-18.
- Carson, M. J., J. Barondess, and J. Beckwith. 1991. The FtsQ protein of *Escherichia coli*: membrane topology, abundance, and cell division phenotypes due to overproduction and insertion mutants. *J. Bacteriol.* **173**:2187-2195.
- Coyette, J., A. Somzé, J. J. Briquet, J. M. Ghuyesen, and R. Fontana. 1983. Function of penicillin-binding protein 3 in *Streptococcus faecium*, p. 523-530. *In* R. Hakenbeck, J. V. Höltje, and H. Labischinski (ed.), *The target of penicillin*. Walter de Gruyter & Co., Berlin, Germany.
- Deckert, G., et al. 1998. The complete genome of the hyperthermophilic bacteria *Aquifex aeolicus*. *Nature* **392**:353-358.
- DiBernardino, M., A. Dijkstra, D. Stüber, W. Keck, and M. Gubler. 1996. The monofunctional glycosyltransferase of *Escherichia coli* is a member of a new class of peptidoglycan-synthesizing enzymes. Overexpression and determination of the glycan-polymerizing activity. *FEBS Lett.* **392**:184-188.
- Dive, G., D. Dehareng, and L. Ghosez. 1998. Catalytic reaction pathways approached by quantum chemistry: a challenge. *Cell. Mol. Life Sci.* **54**:378-382.
- Dive, G., D. Dehareng, and J. M. Ghuyesen. 1994. A detailed study of a molecule into a molecule: the *N*-acetyl-L-tryptophanamide in active site model of the  $\alpha$ -chymotrypsin. *J. Am. Chem. Soc.* **116**:2548-2556.
- Dive, G., D. Dehareng, and D. Peeters. 1996. Proposition for the acylation mechanism of serine proteases: a one-step process? *Int. J. Quantum Chem.* **58**:85-107.
- Dowson, C. G., T. J. Coffey, and B. G. Spratt. 1994. Origin and molecular epidemiology of penicillin-binding protein-mediated resistance to  $\beta$ -lactam antibiotics. *Trends Microbiol.* **2**:361-366.
- Duez, C., I. Thamm, F. Sapunarić, J. Coyette, and J. M. Ghuyesen. The division and cell wall gene cluster of *Enterococcus hirae* S185. DNA Sequence, in press.
- El Kharroubi, A., P. Jacques, G. Piras, J. Coyette, J. Van Beeumen, and J. M. Ghuyesen. 1991. The *Enterococcus hirae* R40 penicillin-binding protein 5 and the methicillin-resistant *Staphylococcus aureus* penicillin-binding protein 2' are homologs. *Biochem. J.* **280**:463-469.
- Fleischmann, R. D., et al. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496-512.
- Frazer, C. M., et al. 1998. Genomic sequence of a Lyme disease spirochaete,

- Borrelia burgdorferi*. Nature **390**:580–586.
22. Frère, J. M. 1995.  $\beta$ -Lactamases and bacterial resistance to antibiotics. Mol. Microbiol. **16**:385–395.
  23. Ghuysen, J. M. 1968. Use of bacteriolytic enzymes in determination of wall structure and their role in cell metabolism. Bacteriol. Rev. **32**:425–464.
  24. Ghuysen, J. M. 1991. Serine  $\beta$ -lactamases and penicillin-binding proteins. Annu. Rev. Microbiol. **45**:37–67.
  25. Ghuysen, J. M. 1994. Molecular structures of penicillin-binding proteins and  $\beta$ -lactamases. Trends Microbiol. **2**:372–380.
  26. Ghuysen, J. M. 1997. Penicillin-binding proteins. Wall peptidoglycan assembly and resistance to penicillin: facts, doubts and hopes. Int. J. Antimicrob. Agents **8**:45–60.
  27. Ghuysen, J. M., P. Charlier, J. Coyette, C. Duez, E. Fonzé, C. Fraipont, C. Goffin, B. Joris, and M. Nguyen-Distèche. 1996. Penicillin and beyond: evolution, protein fold, multimodular polypeptides, and multiprotein complexes. Microb. Drug Resist. **2**:163–175.
  28. Ghuysen, J. M., and G. Dive. 1994. Biochemistry of the penicilloyl serine transferases, p. 103–130. In J. M. Ghuysen and R. Hakenbeck (ed.), Bacterial cell wall. Elsevier Science B.V., Amsterdam, The Netherlands.
  29. Goffin, C., C. Fraipont, J. Ayala, M. Terrak, M. Nguyen-Distèche, and J. M. Ghuysen. 1996. The non-penicillin-binding module of the tripartite penicillin binding protein 3 of *Escherichia coli* is required for folding and/or stability of the penicillin-binding module and the membrane-anchoring module confers cell septation activity on the folded structure. J. Bacteriol. **178**:5402–5409.
  30. Guzman, L. M., J. J. Barondess, and J. Beckwith. 1992. FtsL, an essential cytoplasmic membrane protein involved in cell division in *Escherichia coli*. J. Bacteriol. **174**:7716–7728.
  31. Guzman, L. M., D. S. Weiss, and J. Beckwith. 1997. Domain swapping analysis of FtsI, FtsL, and FtsQ: bitopic membrane proteins essential for cell division in *Escherichia coli*. J. Bacteriol. **179**:5094–5103.
  32. Hakenbeck, R. 1995. Target-mediated resistance to  $\beta$ -lactam antibiotics. Biochem. Pharmacol. **50**:1121–1127.
  33. Hakenbeck, R., A. König, I. Kern, M. van der Linden, W. Keck, D. Billot-Klein, R. Legrand, B. Schoot, and L. Gutmann. 1998. Acquisition of five high-Mr penicillin-binding protein variants during transfer of high-level  $\beta$ -lactam resistance from *Streptococcus mitis* to *Streptococcus pneumoniae*. J. Bacteriol. **180**:1831–1840.
  34. Hölte, J. V. 1998. Growth of the stress-bearing and shape-maintaining murein sacculus of *Escherichia coli*. Microbiol. Mol. Biol. Rev. **62**:181–203.
  35. Joseleau-Petit, D., D. Thévenet, and R. D'Ari. 1994. ppGpp concentration, growth without PBP2 activity, and growth-rate control in *Escherichia coli*. Mol. Microbiol. **13**:911–917.
  36. Kaneko, T., et al. 1996. Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. DNA Res. **3**:109–136.
  37. Kato, J. I., S. Hideho, and Y. Hirota. 1985. Dispensability of either penicillin-binding protein-1a or -1b involved in the essential process for cell elongation in *Escherichia coli*. Mol. Gen. Genet. **200**:272–277.
  38. Kelly, J. A., A. P. Kuzin, P. Charlier, and E. Fonzé. 1991. X-ray studies of enzymes that interact with penicillins. Cell. Mol. Life Sci. **54**:353–358.
  39. Knox, J. R. 1995. Extended-spectrum and inhibitor-resistant TEM-type  $\beta$ -lactamases: mutations, specificity, and three-dimensional structure. Antimicrob. Agents Chemother. **39**:2593–2601.
  40. Krauss, J., M. van der Linden, T. Grebe, and R. Hakenbeck. 1996. Penicillin-binding proteins 2x and 2b as primary PBP targets in *Streptococcus pneumoniae*. Microb. Drug Resist. **2**:183–186.
  41. Kunst, F., et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. Nature **390**:249–256.
  42. Lefèvre, F., M. H. Rémy, and J. M. Masson. 1997. Topographical and functional investigation of *Escherichia coli* penicillin-binding protein 1b by alanine stretch scanning mutagenesis. J. Bacteriol. **179**:4761–4767.
  43. Lepage, S., P. Dubois, T. Ghosh, B. Joris, S. Mahapatra, M. Kundu, J. Basu, P. Chakrabarti, S. T. Cole, M. Nguyen-Distèche, and J. M. Ghuysen. 1997. Dual multimodular class A penicillin-binding proteins in *Mycobacterium leprae*. J. Bacteriol. **179**:4627–4630.
  44. Ligozzi, M., F. Pittaluga, and R. Fontana. 1996. Modification of penicillin-binding protein 5 associated with high-level ampicillin resistance in *Enterococcus faecium*. Antimicrob. Agents Chemother. **40**:354–357.
  45. Livingstone, G. D., and G. J. Barton. 1996. Identification of functional residues and secondary structure from protein multiple sequence alignment. Methods Enzymol. **266**:497–512.
  46. Löwe, J., and L. A. Amos. 1998. Crystal structure of the bacterial cell-division protein FtsZ. Nature **391**:203–206.
  47. Lutkenhaus, J., and S. G. Addinall. 1997. Bacterial cell division and the Z ring. Annu. Rev. Biochem. **66**:93–116.
  48. Matagne, A., J. Lamotte-Brasseur, and J. M. Frère. 1998. Catalytic properties of class A  $\beta$ -lactamases: efficiency and diversity. Biochem. J. **330**:581–598.
  49. Massova, I., and S. Mobashery. 1998. Kinship and diversification of bacterial penicillin-binding proteins and  $\beta$ -lactamases. Antimicrob. Agents Chemother. **42**:1–17.
  50. Matsuhashi, M. 1994. Utilization of lipid-linked precursors and the formation of peptidoglycan in the process of cell growth and division: membrane enzymes involved in the final steps of peptidoglycan synthesis and the mechanisms of their regulation, p. 55–72. In J. M. Ghuysen and R. Hakenbeck (ed.), Bacterial cell wall. Elsevier Science B.V., Amsterdam, The Netherlands.
  51. Mollerach, M., P. Partoune, J. Coyette, and J. M. Ghuysen. 1996. Importance of the E46-D160 polypeptide segment of the non-penicillin-binding module for the stability of the low-affinity, multimodular class B penicillin-binding protein 5 of *Enterococcus hirae*. J. Bacteriol. **178**:1774–1775.
  52. Morell, V. 1997. Antibiotic resistance: road of no return. Science **278**:575–576.
  53. Nanninga, N. 1998. Morphogenesis of *Escherichia coli*. Microbiol. Mol. Biol. Rev. **62**:110–129.
  54. Nogales, E., K. H. Downing, L. A. Amos, and J. Löwe. 1998. Tubulin and FtsZ form a distinct family of GTPases. Nat. Struct. Biol. **5**:451–458.
  55. Paik, J., D. Jendrossek, and R. Hakenbeck. 1997. A putative monofunctional glycosyltransferase is expressed in *Ralstonia eutropha*. J. Bacteriol. **179**:4061–4065.
  56. Paradkar, A. S., K. A. Aidoo, A. Wong, and S. E. Jensen. 1996. Molecular analysis of a  $\beta$ -lactam resistance gene encoded within the cephamycin gene cluster of *Streptomyces clavuligerus*. J. Bacteriol. **178**:6266–6274.
  57. Pares, S., N. Mouz, Y. Pétillot, R. Hakenbeck, and O. Dideberg. 1996. X-ray structure of *Streptococcus pneumoniae* PBP2x, a primary penicillin target enzyme. Nat. Struct. Biol. **3**:284–289.
  58. Piras, G., D. Raze, A. El Kharroubi, D. Hastir, S. Englebert, J. Coyette, and J. M. Ghuysen. 1993. Cloning and sequencing of the low affinity penicillin-binding protein 3'-encoding gene of *Enterococcus hirae* S185. Modular design and structural organization of the protein. J. Bacteriol. **175**:2844–2852.
  59. Popham, D. L., and P. Setlow. 1996. Phenotypes of *Bacillus subtilis* mutants lacking multiple class A high-molecular-weight penicillin-binding proteins. J. Bacteriol. **178**:2079–2085.
  60. Raze, D., O. Dardenne, S. Hallut, M. Martínez-Bueno, J. Coyette, and J. M. Ghuysen. 1998. The low-affinity penicillin-binding protein 3R-encoding gene of *Enterococcus hirae* S185R is borne on a plasmid carrying other antibiotic resistance determinants. Antimicrob. Agents Chemother. **42**:534–539.
  61. Reddy, P. S., A. Raghavan, and D. Chatterjee. 1995. Evidence of a ppGpp-binding site in *Escherichia coli* RNA polymerase: proximity relationship with the rifampicin-binding domain. Mol. Microbiol. **15**:255–265.
  62. Salyers, A. A., and C. F. Amabile-Cuevas. 1997. Why are antibiotic resistance genes so resistant to elimination? Antimicrob. Agents Chemother. **41**:2321–2325.
  63. Sanchez, M., A. Valencia, M. J. Ferrandiz, C. Sander, and M. Vicente. 1994. Correlations between the structure and biochemical activities of FtsA, an essential cell division protein of the actin family. EMBO J. **13**:4919–4925.
  64. Spratt, B. G., J. Zhou, M. Taylor, and M. J. Merrick. 1996. Monofunctional biosynthetic peptidoglycan transglycosylases. Mol. Microbiol. **19**:639–640.
  65. Tomb, J. F., et al. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. Nature **388**:539–547.
  66. van Heijenoort, J. 1996. Murein synthesis, p. 1025–1034. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C.
  67. Vicente, M., and J. Errington. 1996. Structure, function and controls in microbial division. Mol. Microbiol. **20**:1–7.
  68. Ward, J. B., and H. R. Perkins. 1973. The direction of glycan synthesis in a bacterial peptidoglycan. Biochem. J. **135**:721–728.
  69. Weston, A., J. B. Ward, and H. R. Perkins. 1977. Biosynthesis of peptidoglycan in wall plus membrane preparations from *Micrococcus luteus*: direction of chain extension, length of chains and effect of penicillin on cross-linking. J. Gen. Microbiol. **99**:171–181.
  70. Wilson, E. O. 1993. The diversity of life. W. W. Norton & Co., New York, N.Y.
  71. Wu, E. C. Y., W. E. Alborn Jr., J. E. Flokowsch, J. Hoskins, S. Unal, L. C. Blaszcak, D. A. Preston, and P. L. Skatrud. 1994. Site-directed mutagenesis of the *mecA* gene from methicillin-resistant strains of *Staphylococcus aureus*. J. Bacteriol. **176**:443–449.
  72. Wu, S., C. Piscitelli, H. de Lencastre, and A. Tomasz. 1996. Tracking the origin of an antibiotic-resistance gene: methicillin-susceptible strains of the animal species *Staphylococcus sciuri* carry as a native gene a homologue of the *Staphylococcus aureus* methicillin resistance determinant *mecA*. Microb. Drug Resist. **2**:435–441.
  73. Young, K. D., S. A. Denome, P. K. Elf, and T. A. Henderson. 1997. Use of a comprehensive set of PBP mutants to investigate peptidoglycan synthesis in *Escherichia coli*, p. 41. In The Bacterial Cell Cycle Workshop in Chorin, Germany 13 to 17 September 1997.
  74. Zorzi, W., X. Y. Zhou, O. Dardenne, J. Lamotte, D. Raze, J. Pierre, L. Gutmann, and J. Coyette. 1996. Structure of the low-affinity penicillin-binding protein 5 PBP5fm in wild-type and highly penicillin-resistant strains of *Enterococcus faecium*. J. Bacteriol. **178**:4948–4957.