

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

Natural and projectively equivariant quantizations

Fabian Radoux

Przemysl

1 May 2007

Introduction

- At the origin : $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with $P = \partial_x$; $Q = x$.

Introduction

- At the origin : $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with $P = \partial_x$; $Q = x$.

Problems : It depends on the order

Introduction

- At the origin : $qp \rightarrow QP$

$$qp \rightarrow \frac{1}{2}(QP + PQ)$$

with $P = \partial_x$; $Q = x$.

Problems : It depends on the order

It depends on the coordinates system

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- Quantization : $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

- Quantization : $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

Q : linear bijection

- Quantization : $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

Q : linear bijection

preserves the principal symbol

- Quantization : $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

Q : linear bijection

preserves the principal symbol

- $\#Q : Q(\Phi^* S) = \Phi^* Q(S) \forall$ local diffeomorphism Φ

- Quantization : $Q : \mathcal{S}(M) \mapsto \mathcal{D}_{\frac{1}{2}}(M)$

Q : linear bijection

preserves the principal symbol

- $\#Q : Q(\Phi^* S) = \Phi^* Q(S) \forall$ local diffeomorphism Φ
 $\#Q : Q(L_X S) = L_X Q(S) \forall$ vector field X

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$
 $\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$
 $\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$
- $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

- $\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$

- $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$

- $X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^*$ vector field on \mathbb{R}^m

"Flat" case :

- $\exists Q : Q(\Phi_g^* S) = \Phi_g^* Q(S) \forall g \in G$

$$\exists Q : Q(L_h^* S) = L_h^* Q(S) \forall h \in \mathfrak{g}$$

- $PGL(m+1, \mathbb{R})$ acts on $\mathbb{R}P^m$

$$X \in \mathfrak{sl}(m+1, \mathbb{R}) \mapsto X^* \text{ vector field on } \mathbb{R}^m$$

$$\exists Q : L_X Q(S) = Q(L_X S) \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

- Method of the Casimir operator :

$$\mathcal{C} : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

- Method of the Casimir operator :

$$C : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition : $\exists Q$ such that if $C(S) = \alpha S$, then

- Method of the Casimir operator :

$$C : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition : $\exists Q$ such that if $C(S) = \alpha S$, then

$$C(Q(S)) = \alpha Q(S)$$

- Method of the Casimir operator :

$$\mathcal{C} : \mathcal{S}(\mathbb{R}^m) \mapsto \mathcal{S}(\mathbb{R}^m) \quad ; \quad \mathcal{C} : \mathcal{D}(\mathbb{R}^m) \mapsto \mathcal{D}(\mathbb{R}^m)$$

Condition : $\exists Q$ such that if $\mathcal{C}(S) = \alpha S$, then

$$\mathcal{C}(Q(S)) = \alpha Q(S) \text{ and } \sigma(Q(S)) = S$$

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

"Curved" case :

- $Q(\nabla) : \mathcal{S}^3(M) \mapsto \mathcal{D}^3(M)$

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- Conjecture : $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- Conjecture : $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$ natural

- Conjecture : $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$ natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

- Conjecture : $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$ natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

$$\phi_t^* Q(\nabla^0)(S) = Q(\phi_t^* \nabla^0)(\phi_t^* S)$$

- Conjecture : $Q(\nabla) : \mathcal{S}(M) \mapsto \mathcal{D}(M)$ natural and

$$Q(\nabla) = Q(\nabla') \text{ if } \nabla' = \nabla + \alpha \vee id$$

$$\phi_t^* Q(\nabla^0)(S) = Q(\phi_t^* \nabla^0)(\phi_t^* S)$$

$$L_X Q(\nabla^0)(S) = Q(\nabla^0)(L_X S) \quad \forall X \in \mathfrak{sl}(m+1, \mathbb{R})$$

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- M. Bordemann method :

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- M. Bordemann method : $M \mapsto \tilde{M}$;

• M. Bordemann method : $M \mapsto \tilde{M}$;

$$\nabla \mapsto \tilde{\nabla};$$

• M. Bordemann method : $M \mapsto \tilde{M}$;

$$\nabla \mapsto \tilde{\nabla};$$

$$S \mapsto \tilde{S};$$

- M. Bordemann method : $M \mapsto \tilde{M}$;

$$\nabla \mapsto \tilde{\nabla};$$

$$S \mapsto \tilde{S};$$

$$Q(\widetilde{\nabla})(\tilde{S})(f) = \tau(\tilde{\nabla})(\tilde{S})(\tilde{f})$$

Natural and projectively equivariant quantizations

Fabian Radoux

Introduction

Cartan fiber bundles and connections

The case of the densities

Other differential operators

- Questions :

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

The case of the
densities

Other differential
operators

- Questions : Critical values of δ

- Questions : Critical values of δ

Explicit formula

- Questions : Critical values of δ

Explicit formula

Uniqueness

- Questions : Critical values of δ

Explicit formula

Uniqueness

Other differential operators

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

**Cartan fiber
bundles and
connections**

The case of the
densities

Other differential
operators

- Cartan fiber bundles and connections

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \times G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \times G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

- Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \times G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

• Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \times G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

$$f \text{ } G_0\text{-equivariant} \Rightarrow \nabla^\omega f$$

G_0 -equivariant

• Cartan fiber bundles and connections

$$[\nabla] \rightarrow P \mapsto M$$

$$[\nabla] \rightarrow \omega : TP \mapsto \mathfrak{g}$$

$$\omega_u : T_u P \mapsto \mathfrak{g} \text{ bijection } \forall u \in P$$

$$T \rightarrow p^* T \in \mathcal{C}^\infty(P, V)_H$$

$$\text{with } H = G_0 \times G_1, \mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$$

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{h}, \mathfrak{g}_{-1} \cong \mathbb{R}^m$$

$$\omega \rightarrow \nabla^\omega(e_i) = L_{\omega^{-1}(e_i)}, e_i \in \mathfrak{g}_{-1}$$

$$f \text{ } G_0\text{-equivariant} \Rightarrow \nabla^\omega f$$

G_0 -equivariant

$$f \text{ } G_1\text{-equivariant} \not\Rightarrow \nabla^\omega f$$

G_1 -equivariant

Natural and
projectively
equivariant
quantizations

Fabian Radoux

Introduction

Cartan fiber
bundles and
connections

**The case of the
densities**

Other differential
operators

- The case of the densities (P. Mathonet, R.) :

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(e_i)}$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

$$\text{Condition : } L_{h^*} Q(p^* S)(p^* f) = 0 \quad \forall h \in \mathfrak{g}_1$$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

Condition : $L_{h^*} Q(p^* S)(p^* f) = 0 \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$ not G_1 -equivariant !

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

Condition : $L_{h^*} Q(p^* S)(p^* f) = 0 \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$ not G_1 -equivariant !

One adds terms of lower orders in $p^* f \dots$

- The case of the densities (P. Mathonet, R.) :

$$S \rightarrow p^* S \in \mathcal{C}^\infty(P, S_\delta^k(\mathbb{R}^m))_H$$

$$f \rightarrow p^* f \in \mathcal{C}^\infty(P, \Delta^\lambda(\mathbb{R}^m))_H$$

$$\omega \rightarrow \text{Div}^\omega = \sum_i i(\epsilon^i) L_{\omega^{-1}(\epsilon_i)}$$

Condition : $L_{h^*} Q(p^* S)(p^* f) = 0 \quad \forall h \in \mathfrak{g}_1$

$\langle p^* S, \nabla_s^{\omega^k} p^* f \rangle$ not G_1 -equivariant !

One adds terms of lower orders in $p^* f \dots$

One finds then :

$$Q_M(\nabla, S)(f) = p^{*-1} \left(\sum_{l=0}^k C_{k,l} \langle \text{Div}^{\omega^l} p^* S, \nabla_s^{\omega^{k-l}} p^* f \rangle \right),$$

$$\text{with } C_{k,l} = \frac{(\lambda + \frac{k-1}{m+1}) \dots (\lambda + \frac{k-l}{m+1})}{\gamma_{2k-1} \dots \gamma_{2k-l}} \binom{k}{l}, \quad \forall l \geq 1, \quad C_{k,0} = 1$$

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

"Curved" case

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization Q_{Aff} :
 ∂_i

"Curved" case

"Affine" quantization Q_ω :
 $L_{\omega^{-1}(e_i)}$

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization $Q_{Aff} :$
 ∂_i

Application $\gamma :$

$$\mathcal{L}_{X^h} Q_{Aff}(S)(f) = Q_{Aff}((L_{X^h} + \gamma(h))S)(f)$$

"Curved" case

"Affine" quantization $Q_\omega :$
 $L_{\omega^{-1}(e_i)}$

Application $\gamma :$

$$\mathcal{L}_{h^*} Q_\omega(S)(f) = Q_\omega((L_{h^*} + \gamma(h))S)(f), \quad h \in \mathfrak{g}_1$$

- Other differential operators (P. Mathonet, R.) : method of the Casimir operator :

"Flat" case

Affine quantization $Q_{Aff} :$
 ∂_i

Application $\gamma :$

$$\mathcal{L}_{X^h} Q_{Aff}(S)(f) = Q_{Aff}((L_{X^h} + \gamma(h))S)(f)$$

Casimirs C and C
 ∂_i

"Curved" case

"Affine" quantization $Q_\omega :$
 $L_{\omega^{-1}(e_i)}$

Application $\gamma :$

$$\mathcal{L}_{h^*} Q_\omega(S)(f) = Q_\omega((L_{h^*} + \gamma(h))S)(f), h \in \mathfrak{g}_1$$

"Casimirs" C^ω and C^ω
 $L_{\omega^{-1}(e_i)}$

Quantization :

$Q_{Aff}(Q(S))$, $Q(S)$ such that
if $C(S) = \alpha S$, then

$\mathcal{C}(Q(S)) = \alpha Q(S)$ and

"head" of $Q(S) = S$

Quantization :

$Q_{\omega}(Q(S))$, $Q(S)$ such that
if $C^{\omega}(S) = \alpha S$, then

$\mathcal{C}^{\omega}(Q(S)) = \alpha Q(S)$ and

"head" of $Q(S) = S$

Quantization :

$Q_{Aff}(Q(S))$, $Q(S)$ such that
if $C(S) = \alpha S$, then

$$\mathcal{C}(Q(S)) = \alpha Q(S) \text{ and}$$

"head" of $Q(S) = S$

Then :

$$\mathcal{L} \circ Q = Q \circ L$$

because

$$[\mathcal{C}, \mathcal{L}] = 0 \text{ and } [C, L] = 0$$

Quantization :

$Q_\omega(Q(S))$, $Q(S)$ such that
if $C^\omega(S) = \alpha S$, then

$$\mathcal{C}^\omega(Q(S)) = \alpha Q(S) \text{ and}$$

"head" of $Q(S) = S$

Then :

$$(L_{h^*} + \gamma(h)) \circ Q = Q \circ L_{h^*}$$

because $[\mathcal{C}^\omega, L_{h^*} + \gamma(h)] = 0$

$$\text{and } [\mathcal{C}^\omega, L_{h^*}] = 0$$

Quantization :

$Q_{Aff}(Q(S))$, $Q(S)$ such that
if $C(S) = \alpha S$, then

$$\mathcal{C}(Q(S)) = \alpha Q(S) \text{ and}$$

"head" of $Q(S) = S$

Then :

$$\mathcal{L} \circ Q = Q \circ L$$

because

$$[\mathcal{C}, \mathcal{L}] = 0 \text{ and } [C, L] = 0$$

• Conclusion : "Flat" case
 ∂_i

Quantization :

$Q_\omega(Q(S))$, $Q(S)$ such that
if $C^\omega(S) = \alpha S$, then

$$\mathcal{C}^\omega(Q(S)) = \alpha Q(S) \text{ and}$$

"head" of $Q(S) = S$

Then :

$$(L_{h^*} + \gamma(h)) \circ Q = Q \circ L_{h^*}$$

because $[C^\omega, L_{h^*} + \gamma(h)] = 0$

$$\text{and } [C^\omega, L_{h^*}] = 0$$

"Curved" case
 $L_{\omega^{-1}(e_j)}$