


This article was downloaded by:[SCD - Universite Bordeaux 2]
On: 17 October 2007
Access Details: [subscription number 773444277]
Publisher: Informa Healthcare
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Enzyme Inhibition and Medicinal Chemistry

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title-content=t713642123>

Quantitative structure-activity relationships studies of antioxidant hexahydropyridoindoles and flavonoid derivatives

Anne-Catherine Durand ^a; Amaury Farce ^a; Pascal Carato ^a; Sebastien Dilly ^a;
Said Yous ^a; Pascal Berthelot ^a; Philippe Chavatte ^a
^a Faculté des Sciences Pharmaceutiques et Biologiques, Laboratoire de Chimie
Thérapeutique, Lille Cedex, France

Online Publication Date: 01 October 2007

To cite this Article: Durand, Anne-Catherine, Farce, Amaury, Carato, Pascal, Dilly, Sebastien, Yous, Said, Berthelot, Pascal and Chavatte, Philippe (2007)

'Quantitative structure-activity relationships studies of antioxidant hexahydropyridoindoles and flavonoid derivatives',
Journal of Enzyme Inhibition and Medicinal Chemistry, 22:5, 556 - 562

To link to this article: DOI: 10.1080/14756360701425238

URL: <http://dx.doi.org/10.1080/14756360701425238>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Quantitative structure-activity relationships studies of antioxidant hexahydropyridoindoles and flavonoid derivatives

ANNE-CATHERINE DURAND, AMAURY FARCE, PASCAL CARATO, SEBASTIEN DILLY, SAID YOUS, PASCAL BERTHELOT, & PHILIPPE CHAVATTE

Faculté des Sciences Pharmaceutiques et Biologiques, Laboratoire de Chimie Thérapeutique, EA1043, 3 rue du Professeur Laguesse, BP 83, 59006 Lille Cedex, France

(Received 16 October 2006; accepted 8 February 2007)

Abstract

In order to predict the antioxidant activity of 22 pinoline derivatives (1,2,3,4-tetrahydro- β -carbolines), two dimensional quantitative-structure activity relationships (2D-QSAR) analysis of 19 hexahydropyridoindoles and 12 flavonoids was realized. Five statistically significant models were obtained from randomly constituted training sets (21 compounds) and subsequently validated with the corresponding test sets (10 compounds). Antioxidant activity (pIC_{50}) was correlated with 5 molecular descriptors calculated with the software DRAGON. The best predictive model ($n = 21$, $q^2 = 0.794$, $N = 2$, $r^2 = 0.888$, $s = 0.157$) could offer structural insights into the features conferring a strong antioxidant activity to compounds built from a pinoline scaffold prior to their synthesis.


Keywords: pinoline, β -carbolines, melatonin, hexahydropyridoindoles, flavonoids, 2D-QSAR, antioxidant

Introduction


N-acetyl-5-methoxytryptamine (melatonin) is a neurohormone synthesized in the pineal gland during the dark period in all species [1]. Besides its effects on circadian and seasonal rhythms [2] it has been found to be one of the most efficient free radical scavenger for hydroxyl radical [3,4] and peroxy radical. In addition, several *in vivo* studies report that melatonin protects cells against oxidative agents such as safrole [5,6], paraquat [7] and kainic acid [8]. The antioxidant properties of melatonin are now well documented [9], and recent studies suggest that structurally related indole-based derivatives and their cyclic analogues such as β -carbolines may have a similar antioxidant capacity [10,11]. Because of the protective effects of melatonin against lipid peroxidation, several melatonin structural analogues have been synthesized, screened and a QSAR analysis with the low-density lipoprotein oxidation model was made

in the laboratory [12,13]. Among the structural analogues of melatonin studied, some derivatives of a minor melatonin metabolite called pinoline (6-methoxy-1,2,3,4-tetrahydro- β -carboline) have been studied for identifying more efficient antioxidant compounds [14,15,16]. Pinoline has been isolated from the nervous system of mammals [17,18] and can be performed under physiological conditions from 5-methoxytryptamine [19] or as a minor metabolite of melatonin [20]. Pähkila et al. first reported the antioxidant activity of pinoline and showed that this compound is five fold less effective than melatonin in scavenging hydroxyl radical. However, considering that the hydroxyl radical scavenging ability of melatonin is much higher than that of well-known antioxidants such as glutathione and mannitol, pinoline can still be considered as a good hydroxyl radical scavenger [21]. Other studies showed that melatonin is slightly more potent than pinoline in inhibiting lipid


Correspondence: P. Chavatte, Faculté des Sciences Pharmaceutiques et Biologiques, Laboratoire de Chimie Thérapeutique, EA1043, 3 rue du Professeur Laguesse, BP 83, 59006 Lille Cedex, France. Tel: 33 3 20 96 40 20. Fax: 33 3 20 96 43 61. E-mail: philippe.chavatte@univ-lille2.fr

Table I. Structures and antioxidant activities (pIC₅₀) of hexahydropyridoindoles derivatives

Compounds	R ₁	R ₂	R ₃	R ₄	R ₅	pIC ₅₀
1	CH ₃	H	H	CH ₃	H	4.60
2	CH ₃	NO ₂	H	CH ₃	H	4.58
3	CH ₃	Br	H	CH ₃	H	4.80
4	CH ₃	NH ₂	H	CH ₃	H	4.90
5	CH ₃	N(CH ₃) ₂	H	CH ₃	H	4.66
6	CH ₃	H	H	OH	H	4.68
7	CH ₃	H	H	OCH ₃	H	4.48
8	CO-O- <i>i</i> -Bu	H	H	OCH ₃	H	5.12
9	CO-O-CH ₃	Br	CH ₃	OCH ₃	CH ₃	4.71
10	CO-O-CH ₃	H	CH ₃	OCH ₃	CH ₃	4.71
11	CO-O-Et	H	CH ₃	OCH ₃	CH ₃	4.70
12	CO-O-Pr	H	CH ₃	OCH ₃	CH ₃	5.16
13	CO-O-Bu	H	CH ₃	OCH ₃	CH ₃	5.35
14	CO-O- <i>i</i> -Bu	H	CH ₃	OCH ₃	CH ₃	5.19
15	CO-O-CH ₂ -Ph	H	CH ₃	OCH ₃	CH ₃	5.48
16	CO-O-CH ₂ -P	H	H	OCH ₃	H	5.49
17	CO-O-CH ₂ -P	CH ₃	H	OCH ₃	H	5.29
18	CO-O-CH ₂ -P	CH ₃	H	CH ₃	H	5.36
19	CO-O-CH ₂ -P	CH ₃	CH ₃	Br	H	5.53

Table II. Structures and antioxidant activities (pIC₅₀) of flavonoid derivatives

Compounds	R ₁	R ₂	OH position	pIC ₅₀
20	C ₆ H ₃ -3,4'-OH	OH	5,7	4.43
21	C ₆ H ₄ -4'-OH	H	5,7	4.16
22	C ₆ H ₄ -3'-OH	H	5,7	4.28
23	C ₆ H ₄ -4'-OCH ₃	H	6	3.86



Compounds	R ₁	R ₂	OH position	pIC ₅₀
24	C ₆ H ₃ -3',4'-OH	OH	3,5,7	4.93
25	C ₆ H ₃ -2',4'-OH	OH	3,5,7	4.24
26	C ₆ H ₅	OH	3,5,7	3.92
27	C ₆ H ₅	H	6	3.72
28	C ₆ H ₅	H	7,8	4.22
29	C ₆ H ₅	H	5,7	4.17
30	C ₆ H ₅	H	5,6,7	3.80
31	H	C ₆ H ₄ -4'-OH	7	4.10





Figure 1. Distribution of pIC_{50} of the compounds.

peroxidation *in vitro*, whereas the ability of pinoline to protect rat brain homogenates against H_2O_2 -induced lipid peroxidation was higher than that of melatonin [22,23,24]. Pinoline was also found to be more active than melatonin in reducing nitric oxide-induced lipid peroxidation in rat retinal homogenates [25], and to protect DNA against chromium(III) and H_2O_2 oxidative damages [26]. On the other hand, both pinoline and melatonin participate in the stabilization of hepatic microsomal membranes owing to their capacity to inhibit membrane lipid peroxidation [27]. Therefore it appears that pinoline is endowed with antioxidant activity although this depends on the nature of the radical to scavenge and the tissue to protect. In previous works [14,15,16], some 1-aryl-1,2,3,4-tetrahydro- β -carbolines have been synthesized and their ability to prevent low density lipoprotein (LDL) copper-induced peroxidation in comparison with melatonin and pinoline using quantitative structure-activity relationships has been investigated [15]. The best compound of the series, identified as [6-ethyl-1-(3-methoxyphenyl)-2-propyl-1,2,3,4-tetrahydro- β -carboline] chlorhydrate (GWC22), displays an ethyl group in the 6 position of the β -carboline, that increased the antioxidant activity [16]. This work aimed to draw two dimensional quantitative structure-activity relationships on the effects of different substituents on the nitrogen atom in position 2 and the aromatic ring, onto the antioxidant activity. In order to build a predictive model of the antioxidant activity of these compounds, we used hexahydropyridoindoles [28] and flavonoids [29] reported from the literature. The chemical structure of the 19 hexahydropyridoindoles studied by Rackova et al. is very close to that of our β -carbolines and the introduction of flavonoids



$R_1, R_2 =$ alkyl groups.




Figure 2. Distributions of chemical families (a) and biological activities (pIC_{50}) (b) *versus* number of compounds for the training (black) and test (grey) sets of model 1.

in this model can allow us to obtain a wider spectrum of prediction.

Materials and methods

Selection of compounds

Rackova et al. synthesized 31 compounds. Their antioxidant efficacy against lipid peroxidation of liposomal membranes was studied in a biphasic system comprising suspension of dioleoylphosphatidylcholine (DOPC) liposomes and peroxy radicals continually generated from 2,2-azobis(2-amidinopropane) hydrochloride (AAPH). The antioxidant activity was expressed in terms of pIC_{50} or $\log(1/IC_{50})$ (Tables I and II). The 31 molecules selected belong to two different chemical families: hexahydropyridoindoles [28] (Family A: 19 compounds) and flavonoids [29] (Family B: 12 compounds). A homogenous repartition of the antioxidant activity between the two chemical families is a prerequisite if meaningful results are to be obtained from a quantitative structure-activity relationships study (Figure 1 and 2). A multi-model approach was used to assess the predictive power of the final model. Approximately two thirds of the 31 compounds were divided into 5 individual training sets of 21 compounds each and the remaining third was used as test sets with 10 compounds each. Compounds were randomly split between training and test sets by following Oprea et al. suggestions [30].

Molecular modeling

Molecular modeling studies were performed using SYBYL 6.9.1 [31] running on Silicon Graphics workstations. Three-dimensional models of compounds

Table III. Statistical results for the five 2D-QSAR models.

	Model 1	Model 2	Model 3	Model 4	Model 5
q^2 ^a	0.794	0.876	0.890	0.898	0.880
N ^b	2	5	2	3	2
s_{cv} ^c	0.213	0.209	0.203	0.185	0.209
r^2 ^d	0.888	0.929	0.931	0.934	0.929
s ^e	0.157	0.158	0.161	0.148	0.161
F ^f	71.708	39.245	121.992	80.705	117.327
r_{pred}^2 ^g	0.952	0.918	0.907	0.901	0.939

^aCrossvalidation correlation coefficient; ^bNumber of components; ^cStandard error of prediction; ^dCorrelation coefficient; ^eStandard error of estimate; ^fF-ratio; ^gPredicted correlation coefficient.

were built from a standard fragments library and their geometry was subsequently optimized with the semi-empirical MOPAC 6.0 package using the hamiltonian AM1 (keywords: PRECISE, NOMM, PARASOK, XYZ) and Coulson atomic charges were calculated using the same method [32].

Selection of the descriptors

The software DRAGON 2.1 [33] was employed to select the best descriptors among 18 families of descriptors. 35 descriptors were retained due to their usefulness in a correct correlation with the pIC_{50} of the compounds. They belong to constitutional, topological, charge, empirical descriptors, aromaticity indices, functional groups and properties. Among them, only 5 descriptors were chosen based on their high correlation with the biological activity (Table V).

χ_2 : the 2nd order connectivity index, usually known as Kier-Hall connectivity indice (topological descriptors). It is a variant of the Randic connectivity index and can be interpreted as the contribution of one molecule to the biomolecular interaction arising from the encounters of bonds of two identical molecules.

IC_5 : the 5th order information content index or neighbourhood information content is based on neighbour degrees and edge multiplicity, and is an indice of neighbourhood symmetry (topological descriptors).

ICR: the radial centric information index quantifies the degree of compactness of molecules by distinguishing between molecular structures organized differently with respect to their centres, i.e. long linear chains or short ramified substitution pattern (molecular descriptor).

nCT: total number of tertiary carbons (functional group descriptors).

nCO: total number of aliphatic ketones (functional group descriptors).

These five descriptors were chosen because they took into account the structural differences of the studied

compounds and gave the highest crossvalidated q^2 with the smallest number of components N . Although electronic, energetic and lipophilic parameters such as Hammett sigma (σ^+), hydration energy (E_{HYDR}), enthalpy of formation (ΔH), energy of the lowest occupied molecular orbital $\epsilon(LUMO)$, energy of the highest occupied molecular orbital $\epsilon(HOMO)$ or the partition coefficient $ClogP$ are often used in the calculation of the antioxidant activity in QSAR studies, they did not show an increase of q^2 value in this case, probably because they did not accurately handle the differences into each family of compounds.

2D-QSAR studies

The 2D-QSAR equation was derived by using a linear regression analysis. The partial least square (PLS) method implemented in the QSAR module of SYBYL was used to build and validate the models. The optimal number of components N retained for the final PLS analysis was defined as the one that yielded the highest crossvalidated q^2 value and which normally had the smallest standard error of prediction s_{cv} . The robustness of the model was internally evaluated by calculating r^2 , s and F values from the training sets and was externally validated by calculating r_{pred}^2 from the test set.

Presentation of the results

The results obtained for the best model are expressed as a representation of the predicted values of pIC_{50} versus experimental values of pIC_{50} for the training and test sets.

Results and discussion

In order to obtain the best predictive 2D-QSAR model we used a multi-model approach.

Training and test sets

A great attention was paid to the distribution of biological activities and structural classification of compounds in both training and test sets. All models show homogeneity in the distribution of antioxidant activity and structural characteristics of their compounds, which were split between training and test sets randomly as presented for model 1 in Figure 2.

D-QSAR models

Five different training sets were used to derive five separate 2D-QSAR models. The results obtained are represented in Table III. Although the predictivity of a model can only be accessed validating it using a test set, the results indicate the robustness of the models, which yield crossvalidated correlation coefficients q^2

Table IV. Predicted values *versus* actual values for the test set of model 1.

Compounds	Experimental pIC ₅₀	Predicted pIC ₅₀	Difference
5	4.66	4.68	0.02
13	5.35	5.26	0.09
14	5.19	5.27	0.08
18	5.36	5.36	0.00
19	5.53	5.36	0.17
22	4.28	4.10	0.18
23	3.86	4.00	0.14
27	3.72	3.91	0.19
28	4.22	4.00	0.22
30	3.80	4.00	0.20

(from 0.794 to 0.898) with reasonable respective standard errors of prediction s_{cv} (from 0.185 to 0.203). A q^2 value of 0.3 corresponds to a confidence limit greater than 95%, which minimizes the risk of finding correlation just by mere chance [34]. Our five models yielded high conventional r^2 (from 0.888 to 0.934) with relatively low standard errors of estimate s (from 0.148 to 0.161) by using the optimal number of components (2).

Predictivity of the models

To validate our models, we attempted to predict the activities for the 10 compounds of the test sets. The calculated correlation coefficient r_{pred}^2 are given in Table IV. Among the five models, model 1 yielding a good r_{pred}^2 (0.952) appears to be the best predictive one. Low differences between the experimental pIC₅₀ and the predicted pIC₅₀ are obtained for the test set of

the model 1 (Table IV) and all compounds were predicted with an error lower than 0.3. More over the predicted pIC₅₀ values *versus* the experimental pIC₅₀ values for both training and test sets of the model 1 are linear and without outliers (Figure 3).

The 2D-QSAR equation obtained for model 1 is the following:

$$\begin{aligned} \text{pIC}_{50} = & 2.698 + 0.093 \chi_2 - 0.183 \text{ICR} \\ & + 0.400 \text{IC}_5 + 0.405 \text{nCT} - 0.675 \text{nCO} \\ n = & 21, r^2 = 0.888, s = 0.157, F_{2,18} = 71,708 \end{aligned} \quad (1)$$

Table V. Values of the 5 descriptors chosen.

Compounds	χ_2	ICR	IC ₅	nCT	nCO
1	6.00	1.00	4.00	0.00	0.00
2	8.00	2.00	4.00	0.00	0.00
3	7.00	1.00	4.00	0.00	0.00
4	7.00	1.00	4.00	0.00	0.00
5	8.00	2.00	4.00	0.00	0.00
6	6.00	1.00	4.00	0.00	0.00
7	7.00	2.00	4.00	0.00	0.00
8	9.00	2.00	4.00	1.00	0.00
9	9.00	2.00	4.00	0.00	0.00
10	9.00	2.00	4.00	0.00	0.00
11	9.00	2.00	4.00	0.00	0.00
12	9.00	2.00	5.00	0.00	0.00
13	10.00	2.00	5.00	0.00	0.00
14	10.00	2.00	4.00	1.00	0.00
15	11.00	2.00	5.00	0.00	0.00
16	10.00	2.00	5.00	0.00	0.00
17	11.00	2.00	5.00	0.00	0.00
18	11.00	2.00	5.00	0.00	0.00
19	11.00	2.00	5.00	0.00	0.00
20	10.00	2.00	5.00	0.00	1.00
21	10.00	2.00	5.00	0.00	1.00
22	10.00	2.00	5.00	0.00	1.00
23	8.00	2.00	4.00	0.00	1.00
24	7.00	2.00	4.00	0.00	1.00
25	8.00	2.00	4.00	0.00	1.00
26	8.00	2.00	4.00	0.00	1.00
27	8.00	2.00	4.00	0.00	1.00
28	9.00	2.00	4.00	0.00	1.00
29	9.00	2.00	4.00	0.00	1.00
30	8.00	2.00	4.00	0.00	1.00
31	8.00	2.00	4.00	0.00	1.00




Figure 3. Predicted values *versus* experimental values for the training (a) and test (b) sets of model 1. General structure of 1,2,3,4-tetrahydro- β -carbolines R₁, R₂ = alkyl groups.

Coefficients obtained for this equation show that the 2nd order connectivity index, the 5th order neighbourhood information content and the total number of tertiary carbons increase pIC₅₀. That means that the more ramified the substitution pattern and the higher the number of tertiary carbons are, the higher the antioxidant activity will be. On the contrary, the radial centric information index and the total number of aliphatic ketones decrease pIC₅₀, what means that compact molecules with aliphatic ketones must be avoided to keep a good antioxidant activity. This last point agrees with the experimental values of pIC₅₀ for the 12 flavonoids. Most of them display a pIC₅₀ inferior to the worst of hexahydropyridoindoles (see Tables I and II).

Conclusion

In conclusion, a series of 19 hexahydropyridoindoles and 12 flavonoids, reported from the literature, were used in the present study to generate and validate 2D-QSAR models. Topological descriptors (χ_2 , IC₅), molecular descriptors (ICR) and functional group parameters (nCT, nCO) showed the highest predictive power and a highly significant predictive correlation ($r_{\text{pred}}^2 = 0.952$) was obtained, whereas electronic, energetic or lipophilic parameters were expected to give the best results but failed. This demonstrates that although it is important to take account of electronic and energetic parameters and lipophilicity for an antioxidant activity, topological, molecular and functional groups parameters are also very important and can sometimes give a better view of the structural differences of various families of compounds. It highlights that the antioxidant activity can be governed by topological, molecular and functional groups parameters. This model could be further applied for the prediction of the antioxidant activity of 22 1,2,3,4-tetrahydro- β -carbolines designed in our laboratory but it will be finally validated only when the designed compounds are synthesized and evaluated.

References

- [1] Vijayalaxmi TCR, Thomas CRJr, Reiter RJ, Herman TS. Melatonin: From basic research to cancer treatment clinics. *J Clin Oncol* 2002;20:2575–2601.
- [2] Delagrangé P, Guardiola-Lemaitre B, Delagrangé P, Guardiola-Lemaitre B. Melatonin, its receptors, and relationships with biological rhythm disorders. *Clin Neuropharmacol* 1997;20:482–510.
- [3] Tan DX, Chen LD, Poeggeler B, Manchester LC, Reiter R. Melatonin: A potent endogenous hydroxyl radical scavenger. *Endocrinol J* 1993;1:215–218.
- [4] Stasica P, Ulanski P, Rosiak JM. Melatonin as a hydroxyl radical scavenger. *J Pineal Res* 1998;25:65–66.
- [5] Tan DX, Poeggeler B, Reiter RJ, Chen LD, Chen S, Manchester LC, Barlow-Walden LR. The pineal hormone melatonin inhibits DNA-adduct formation induced by the chemical carcinogen safrole in vivo. *Cancer Lett* 1993;70:65–71.
- [6] Tan DX, Reiter RJ, Chen LD, Poeggeler B, Manchester LC, Barlow-Walden LR. Both physiological and pharmacological levels of melatonin reduce DNA adduct formation induced by the carcinogen safrole. *Carcinogenesis* 1994;15:215–218.
- [7] Melchiorri D, Reiter RJ, Attia AM, Hara M, Burgos A, Nistico G. Potent protective effect of melatonin on in vivo paraquat-induced oxidative damage in rats. *Life Sci* 1995;56:83–89.
- [8] Melchiorri D, Reiter RJ, Severynek E, Chen LD, Nistico G. Melatonin reduces kainate-induced lipid peroxidation in homogenates of different brain regions. *FASEB J* 1995;9:1205–1210.
- [9] Reiter RJ, Tan DX, Cabrera J, D'Arpa D. Melatonin and tryptophan derivatives as free radical scavengers and antioxidants. *Adv Exp Med Biol* 1999;467:379–387.
- [10] Matuszak Z, Reszak KJ, Chignell CF. Reaction of melatonin and related indoles with hydroxyl radicals: EPR and spin trapping investigations. *Free Radical Bio Med* 1997;23:367–372.
- [11] Stolc S. Indole derivatives as neuroprotectants. *Life Sci* 1999;65:1943–1950.
- [12] Gozzo A, Lesieur D, Duriez P, Fruchart JC, Teissier E. Structure-activity relationships in a series of melatonin analogues with the low-density lipoprotein oxidation model. *Free Radical Bio Med* 1999;26:1538–1543.
- [13] Tailleux A, Gozzo A, Torpier G, Martin-Nizard F, Bonnefont-Rousselot D, Lemdani M, Furman C, Foricher R, Chevè G, Yous S, Micard F, Bordet R, Gardès-Albert M, Lesieur D, Teissier E, Fruchart JC, Fiévet C, Duriez P. Increased susceptibility of low-density lipoprotein to ex vivo oxidation in mice transgenic for human apolipoprotein B treated with 1 melatonin-related compound is not associated with atherosclerosis progression. *J Cardiovasc Pharm* 2005;46:241–249.
- [14] Bonnefont-Rousselot D, Chevè G, Gozzo A, Tailleux A, Guilloz V, Caisey S, Teissier E, Fruchart JC, Delattre J, Jore D, Lesieur D, Duriez P, Gardès-Albert M. Melatonin related compounds inhibit lipid peroxidation during copper or free radical-induced LDL oxidation. *J Pineal Res* 2002;33:107–117.
- [15] Chevè G, Duriez P, Fruchart JC, Teissier E, Poupaert J, Lesieur D. Antioxidant activity of pinolines analogues in the LDL oxidation model. *Med Chem Res* 2003;11:361–379.
- [16] Mekhloufi J, Bonnefont-Rousselot D, Yous S, Lesieur D, Couturier M, Thérond P, Legrand A, Jore D, Gardès-Albert M. Antioxidant activity of melatonin and a pinoline derivative on linoleate model system. *J Pineal Res* 2005;39:27–33.
- [17] Peura P, Johnson JV, Yost RA, Faull KF. Concentrations of tryptoline and methtryptoline in rat brain. *J Neurochem* 1989;52:847–852.
- [18] Kari I. 6-methoxy-1,2,3,4-tetrahydro(-)carboline in pineal gland of chicken and cock. *FEBS Lett* 1981;127:277–280.
- [19] Callaway JC, Ghynther J, Poso A, Vepsäläinen J, Airaksinen MM. The Pictet-Spengler reaction and biogenic tryptamines: Formation of tetrahydro(-)carbolines at physiological pH. *J Het Chem* 1994;31:431–435.
- [20] Airaksinen MM, Kari I. (-)carbolines, psychoactive compounds in the mammalian body. Part I: Occurrence, origin and metabolism. *Med Biol* 1981;59:21–34.
- [21] Pähkla R, Zilmer M, Kullisaar T, Rågo L. Comparison of the antioxidant activity of melatonin and pinoline in vitro. *J Pineal Res*; 1998;24:96–101.
- [22] Pless G, Frederiksen TJ, Garcia JJ, Reiter RJ. Pharmacological aspects of N-acetyl-5-methoxytryptamine (melatonin) and 6-methoxy-1,2,3,4-tetrahydro-beta-carboline (pinoline) as antioxidants: Reduction of oxidative damage in brain region homogenates. *J Pineal Res* 1999;26:236–246.
- [23] Pless G, Frederiksen TJP, Garcia JJ, Reiter RJ. Pharmacological aspects of N-acetyl-5-methoxytryptamine (melatonin) and

- 6-methoxy-1,2,3,4-tetrahydro(-carboline (pinoline) as antioxidants: Reduction of oxidative damage in brain region homogenates. *J Pineal Res* 1999;26:236–246.
- [24] Garcia JJ, Martinez-Ballain E, Robinson M, Allué JL, Reiter RJ, Osuna C, Acuña-Castroviejo D. Protective effect of β -carbolines and other antioxidants on lipid peroxidation due to hydrogen peroxide in rat brain homogenates. *Neurosci Lett* 2000;29:1–4.
- [25] Siu AW, Reiter RJ, To CH. Pineal indoleamines and vitamin E reduce nitric oxide-induced lipid peroxidation in rat retinal homogenates. *J Pineal Res* 1999;27:122–128.
- [26] Qi W, Reiter RJ, Tan DX, Manchester LC, Siu AW, Garcia JJ. Increased levels of oxidatively damaged DNA induced by chromium(III) and H_2O_2 : Protection by melatonin and related molecules. *J Pineal Res* 2000;29:54–61.
- [27] Garcia JJ, Reiter RJ, Pie J, Ortiz GG, Cabrera J, Sainz RM, Acuña-Castroviejo D. Role of pinoline and melatonin in stabilizing hepatic microsomal membranes against oxidative stress. *J Bioenerg Biomembr* 1999;31:609–616.
- [28] Rackova L, Snirc V, Majekova M, Majek P, Stefek M. Free radical scavenging and antioxidant activities of substituted hexahydropyridoindoles. Quantitative structure-activity relationships. *J Med Chem* 2006;49:2543–2548.
- [29] Rackova L, Firakova S, Kostalova D, Stefek M, Sturdik E, Majekova M. Oxidation of liposomal membrane suppressed by flavonoids: Quantitative structure-activity relationships. *Bioorg Med Chem* 2005;13:6477–6484.
- [30] Oprea TI, Waller CL, Marshall GR. Three-dimensional quantitative structure-activity relationship of human immunodeficiency virus (I) protease inhibitors. 2. Predictive power using limited exploration of alternate binding modes. *J Med Chem* 1994;37:2206–2215.
- [31] SYBYL, 6.9 ed., Tripos Associates Inc., 1699 South Hanley Road, St. Louis, MO 63144
- [32] Stewart JJP. MOPAC: A semi-empirical molecular orbital program. *J Comput Aid Mol Des* 1990;4:1–103.
- [33] Todeschini R, Consonni V. Handbook of molecular descriptors. Weinheim: Wiley-VCH; 2000. p 667.
- [34] Clark M, Cramer V, III RD. Validation of the general purpose tripos 5.2 force field. *Quant Struct-Act Rel* 1993;12:137–145.