

ENZYMATIC SYNTHESIS OF NOVEL CARBOHYDRATE SURFACTANTS FOR WATER/SUPERCRITICAL CO₂ EMULSIONS.

Cédric Boyère¹, Audrey Favrelle,¹ Magali Deleu,² Guy Broze,¹ Michel Paquot,² Christophe Blecker,³ Christine Jérôme,^{1*} Antoine Debuigne¹

¹ Center for Education and Research on Macromolecules, University of Liège. Chemistry Institute, Sart-Tilman (B6a) 4000 Liège, Belgium; ² Laboratory of Industrial Biological Chemistry, University of Liège. Gembloux Agro-Bio Tech (GxABT), 5030 Gembloux, Belgium; ³ Laboratory of Food Technology, University of Liège. Gembloux Agro-Bio Tech (GxABT), 5030 Gembloux, Belgium

“Superzym” project is supported by the French Community of Belgium under the scientific “Concerted Research Actions- ARC” program.

The use of aqueous dispersed environments, such as emulsions and miniemulsions, for chemical transformations and polymerization reactions has many advantages over solution processes, i.e. limited environmental impact, ease of products recovery and increased reaction rate. Processes in dispersed media are generally implemented from a water/organic solvent mixture with a surfactant. Nevertheless, supercritical carbon dioxide (Sc-CO₂) (Pc = 74 bars; Tc = 31°C) constitutes an interesting alternative to the traditional organic solvents because it is inexpensive, non-toxic, non-inflammable and environmentally friendlier.^{1,2}

In this context, we develop a novel class of surfactants able to stabilize water/Sc-CO₂ emulsions, i.e. fluorinated and silicone modified carbohydrates. The hydrophilic head of the surfactant consists in a sugar moiety whereas a fluorinated or polysiloxane tail will be specifically located in the Sc-CO₂ phase. The general strategy for the synthesis of these carbohydrates esters surfactants relies on lipase-catalyzed modifications³ of sugars (mannose, glucose), which prevents many protection/deprotection steps. The synthesis and preliminary assessments of these surfactant properties will be presented. Ultimately, we plan to use these compounds as stabilizers for polymerization reactions carried out under water/Sc-CO₂ emulsion conditions.

References:

- 1) Wood C., Tan B., Zhang H., Cooper A., Ed. T.M. Letcher, “Supercritical carbon dioxide as a green solvent for polymer synthesis”, in *Thermodynamics, Solubility and Environmental Issues*, **2007**, Chap 21, 383-396
- 2) Eastoe J., Gold S., Steyler D., Surfactants for CO₂, *Langmuir*, **2006**, 22, 9832-9842
- 3) Degn P., Pedersen L., Duus J., Zimmermann W., Lipase-catalysed synthesis of glucose fatty acid esters in tert-butanol, *Biotechnology Letters*, **1999**, 21, 275-280

Acknowledgements: The authors are grateful to P. Laurent, J.-P. Wathelet, K. Nott, G. Richard and all the partners of the “SUPERZYM” ARC program.