

## CENTA as a Chromogenic Substrate for Studying $\beta$ -Lactamases

CARINE BEBRONE,<sup>1</sup> CATHERINE MOALI,<sup>1</sup> FLORENCE MAHY,<sup>1</sup> SANDRINE RIVAL,<sup>2</sup> JEAN DENIS DOCQUIER,<sup>3</sup> GIAN MARIA ROSSOLINI,<sup>3</sup> JACQUES FASTREZ,<sup>4</sup> REX F. PRATT,<sup>5</sup> JEAN-MARIE FRÈRE,<sup>1</sup> AND MORENO GALLENI<sup>1</sup>

*Centre for Protein Engineering, University of Liège, Institut de Chimie B6, Sart Tilman, 4000 Liège,<sup>1</sup> and Laboratoire de Biochimie Physique et des Biopolymères, Université Catholique de Louvain, Bâtiment Lavoisier, 1348 Louvain-la-Neuve,<sup>2</sup> Belgium; Fachrichtung 8.8 Biochemie, Universität des Saarlandes, D-66123 Saarbrücken, Germany<sup>3</sup>; Dipartimento di Biologia Molecolare, Sezione di Microbiologia, Università di Siena, I-53100 Siena, Italy<sup>4</sup>; and Department of Chemistry, Wesleyan University, Middletown, Connecticut 06459<sup>5</sup>*

### Abstract

CENTA, a chromogenic cephalosporin, is readily hydrolyzed by  $\beta$ -lactamases of all classes except for the *Aeromonas hydrophila* metalloenzyme. Although it cannot practically be used for the detection of  $\beta$ -lactamase-producing strains on agar plates, it should be quite useful for kinetic studies and the detection of the enzymes in crude extracts and chromatographic fractions.


Nitrocefin and, to a lesser extent, PADAC have been used as chromogenic substrates of  $\beta$ -lactamases. Such substrates, whose hydrolysis can be directly monitored in the visible wavelength range, are of particular interest for the kinetic characterization of  $\beta$ -lactamases. Nitrocefin has, for instance, been widely used as a reporter substrate in the study of the inactivation of  $\beta$ -lactamases or of their interactions with poor substrates (3). It also allows the rapid identification of active fractions during  $\beta$ -lactamase purification. However, the price of nitrocefin has recently been increased significantly, and PADAC is no longer commercially available. Synthesis of compounds is also rather tedious. It is thus surprising that a third chromogenic cephalosporin, CENTA (Fig. 1), which can be prepared from the commercially available drug cephalothin, has not received more attention, although it was shown to be sensitive to many  $\beta$ -lactamases (11). In the study described in this report, we determined the kinetic parameters characterizing the interactions between CENTA and a representative set of  $\beta$ -lactamases and some penicillin-binding proteins (PBPs). Although CENTA cannot be used for the direct detection of  $\beta$ -lactamase-producing colonies on agar plates, it still represents an interesting alternative to nitrocefin for the kinetic characterization of  $\beta$ -lactamases.

CENTA was prepared as follows: 3-carboxyl-4-nitrothiophe-nol (TNB) was obtained by dissolving 5.05 mmol (2 g) of 5,5'-dithio-bis-(2-nitrobenzoic acid) in 100 ml of an aqueous solution of 0.5 M Tris base, and the pH was adjusted to 8.0 by addition of 6 M HCl. Dithiothreitol (7.1 mmol, 1.1 g) was added, and the solution turned orange-red. The mixture was stirred for 10 min at 22°C and extracted six times with 25 ml of ethyl acetate before being acidified to pH 1.5 by addition of 6 M HCl. The residual ethyl acetate was eliminated by bubbling nitrogen through the solution, which was thereafter left overnight at 4°C. The precipitate (TNB) was collected by filtration, washed, and dried. The sodium salt of cephalothin (1 g, 2.4 mmol) and 1 equivalent of TNB (478 mg) were dissolved in 19 ml of H<sub>2</sub>O, and the pH was adjusted to 7.0 with 1 M NaOH. The solution was stirred for 6 h at 65°C. The cooled solution was extracted with 10 ml of ethyl acetate, acidified to pH 2.0 with 1 M HCl, and extracted three times with 15 ml of ethyl acetate. The organic phase was washed three times with 15 ml of water, dried over MgSO<sub>4</sub>, and evaporated to dryness in vacuo. The sodium salt of CENTA was obtained by dissolving the dry residue in 25 ml of water containing 1 equivalent of NaHCO<sub>3</sub>, and the solution was freeze-dried, yielding 1.36 g of a light brown solid. Filtration through a Sephadex G-10 column did not modify the yield or improve the purity of the compound, as demonstrated by infrared spectroscopy (KBr) or nuclear magnetic resonance imaging at 200 MHz. The final yield was 1.31 g (94%, with respect to cephalothin).

CENTA was tested as a substrate for class A (TEM-1 [17], NMCA [19], SHV-1 [1], TOHO-1 [10], *Mycobacterium tuberculosis* [22], *Staphylococcus aureus* [21]), class B (*Bacillus cereus* [2], CphA [9], IMP-1 [12], CfiA [16], BlaB [18], VIM-1 [5]), class C (*Enterobacter cloacae* 908R [7], *Pseudomonas aeruginosa*, *Citrobacter freundii*, and ACT-1), and class D (OXA-10 [1], OXA-2 [13])  $\beta$ -lactamases and as an inactivator of

the soluble form of the *Escherichia coli* PBP 5 (20), *Streptomyces* sp. R61 DD-peptidase (8), *Actinomadura* sp. R39 DD-peptidase (8), and *Streptomyces* sp. K15 DD-transpeptidase (14). All kinetic experiments with  $\beta$ -lactamases and PBP 5 were performed at 30°C in 50 mM sodium phosphate (pH 7.0) with 100  $\mu$ M ZnSO<sub>4</sub> added for the *B. cereus*, IMP-1, and CfiA class B metallo- $\beta$ -lactamases. The hydrolysis of CENTA was monitored by continuously recording the absorbance variation at 346 nm ( $\Delta\epsilon = -2,500 \text{ M}^{-1} \text{ cm}^{-1}$ ) or 405 nm ( $\Delta\epsilon = +6,400 \text{ M}^{-1} \text{ cm}^{-1}$ ). The  $k_{\text{cat}}$  and  $K_m$  values were derived from initial rate measurements with the help of the Hanes linearization of the Henri-Michaelis equation and direct fitting on the hyperbolic equation by nonlinear regression or from complete time courses (3). The lowest  $K_m$  values obtained with SHV-1, CfiA, OXA-10, and OXA-2 were verified by using CENTA as a competitive inhibitor versus 100  $\mu$ M nitrocefin for the first three enzymes and versus 300  $\mu$ M cefaclor for the fourth one. The other experimental conditions are detailed in Table 1.

**FIG. 1.** Structure of CENTA.



Inactivation of the R61, R39, and K15 DD-peptidases was monitored by incubating the enzymes with various concentrations of CENTA and measuring the residual activity of an aliquot after increasing periods of time. Activity was determined by measuring the production of D-alanine from  $N^{\alpha},N^{\epsilon}$ -diacetyl-L-lysyl-D-alanyl-D-alanine by the D-amino acid oxidase method (6). In the case of the K15 enzyme, the assay mixture also contained 10 mM glycylglycine. The buffers were as follows: for R39, 50 mM Tris (pH 8.0) plus 1 mM MgCl<sub>2</sub>; for R61 and K15, 50 mM Tris (pH 8.0).

Table 1 summarizes the kinetic constants obtained with CENTA and the various  $\beta$ -lactamases and compares them with those obtained with nitrocefin. CENTA was a relatively good substrate of all the enzymes with the sole exception of the CphA enzyme, which is very specific for carbapenems and similarly exhibits very poor activity against nitrocefin. The kinetic parameters of OXA-10 and SHV-1 with nitrocefin were determined in the present work. When compared to nitrocefin, the initial rates recorded at the same 100  $\mu$ M concentration are of the same order of magnitude with the exceptions of those for the OXA-10 and SHV enzymes. One should be reminded, however, that the  $\Delta\epsilon$  value of CENTA (+6,400  $\text{M}^{-1} \text{ cm}^{-1}$ ) is significantly lower than that of nitrocefin (+ 17,500  $\text{M}^{-1} \text{ cm}^{-1}$ ). Nonetheless, CENTA can easily be used for monitoring the presence of all the enzymes (excepted CphA) in chromatographic fractions and as a reporter substrate for detailed kinetic studies even with enzymes (*M. tuberculosis* and OXA-10) which exhibit rather low levels of activity against this compound. The maximum change of absorbance of the leaving group is different from that for most  $\beta$ -lactam antibiotics. Interestingly, and in contrast to nitrocefin, hydrolysis of CENTA by the OXA-2 class D enzyme does not exhibit the burst phenomenon which precludes the use of the latter substrate in competition and reporter substrate experiments.

Although it was a poor substrate and inactivator of PBP 5 ( $k_{\text{cat}}/K_m = 22 \text{ M}^{-1} \text{ s}^{-1}$ ), the  $k_{\text{cat}}$  value was unexpectedly high for a PBP ( $5 \times 10^{-3} \text{ s}^{-1}$ ), reflecting a relatively rapid deacylation step ( $>5 \times 10^{-3} \text{ s}^{-1}$ ). The  $K_m$  value was, accordingly, rather high (220  $\mu$ M). With the *Streptomyces* sp. R61 and *Actinomadura* sp. R39 enzymes, the second-order inactivation rate constants were of the same order of magnitude as those observed with cephalothin ( $2,000 \pm 200$  and  $90,000 \pm 10,000 \text{ M}^{-1} \text{ s}^{-1}$ , respectively). Deacylation was very slow in both cases ( $<10^{-4} \text{ s}^{-1}$ ). Finally, the K15 enzyme was not sensitive to CENTA ( $k_2/K' < 0.1 \text{ M}^{-1} \text{ s}^{-1}$ ), a result which reflects the low sensitivity of this enzyme to cephalothin ( $k_2/K' = 2 \text{ M}^{-1} \text{ s}^{-1}$  [14]).

**TABLE 1.** Kinetic parameters for interaction between active-site serine and zinc  $\beta$ -lactamases and CENTA of nitrocefin<sup>a</sup>

Enzyme group and enzyme	CENTA						Nitrocefin			$(v_0)_N/(v_0)_C$ <sup>b</sup>
	Methods	[CENTA] (mM)	[Enz] (nM)	$k_{cat}$ (s <sup>-1</sup> )	$K_m$ ( $\mu$ M)	$k_{cat}/K_m$ ( $\mu$ M <sup>-1</sup> s <sup>-1</sup> )	$k_{cat}$ (S <sup>-1</sup> )	$K_m$ ( $\mu$ M)	$k_{cat}/K_m$ ( $\mu$ M <sup>-1</sup> s <sup>-1</sup> )	
<b>Class A <math>\beta</math>-lactamases</b>										
TEM-1	$v_0$	0.025-0.3	11.5	110	70	1.6	930	52	18	17
NMCA	$v_0$	0.25-1.5	7	340	100	3.4	88	104	0.84	19
SHV-1	CTC + RS	0.25-1	1.4-170	11	27	0.4	88	104	0.84	1
TOHO 1	$v_0$	0.25-1.5	1.8	2,0 00	180	11	520	60	9	10
<i>M. tuberculosis</i>	$v_0$	0.25-1.5	310	5	120	0.04	31	80	0.4	22
<i>S. aureus</i>	$v_0$	0.01-0.1	77	2	10	0.2	16	1.5	10	4
<b>Class C <math>\beta</math>-lactamases</b>										
908R	CTC	0.03	10	110	10	11	780	23	34	7
<i>P. aeruginosa</i>	CTC	100-200	25-50	390	63	6.3	740	27	27	7
<i>C. freundii</i>	CTC	100-200	50-90	75	25	3	330	12	28	7
ACT-1	CTC	100-200	1-2	690	80	8.6	ND	ND	ND	ND
<b>Class D <math>\beta</math>-lactamases</b>										
OXA-10	CTC + RS	0.1-1	14- 1,000	0.4	9	0.04	3,000	450	6.6	13
OXA-2	CTC + RS	0.025-0.5	4.8	95	13	7.3	Burst complex kinetic	Burst complex kinetic	Burst complex kinetic	13
<b>Class B <math>\beta</math>-lactamases</b>										
BeII	$v_0$	0.25-1.5	14	50	330	0.15	45	70	0.64	16
CphA	$v_0$	0.1	6,000	NH	NH	NH	0.31	100	0.0003	9
IMP-1	$v_0$	0.25-1.5	1	400	200	2	63	27	2.3	12
CfiA	CTC + RS	0.025-0.5	2.5-10	40	7	5.7	200	16	12.5	23
BlaB	$v_0$	0.05-0.4	14	4.5	31	0.15	17	66	0.26	3
VIM-1	$v_0$	0.05-2.2	1.8	430	600	0.7	95	17	5.6	5


<sup>a</sup> Complete time courses (CTC) were recorded over 3- to 15 min periods; initial rate ( $v_0$ ) values were recorded over 1 min. With CphA, no hydrolysis (NH) of 100  $\mu$ M CENTA could be detected after 30 min (initial rate,  $< 5 \mu$ M h<sup>-1</sup>). The low  $K_m$  values were measured as i $\zeta$ s with the help of a reporter substrate (RS) (usually 100  $\mu$ M nitrocefin, with the exception of the OXA-2 enzyme, with which 300  $\mu$ M cefaclor was used). Standard deviation values did not exceed 10%.

<sup>b</sup> Kinetic data for nitrocefin were found in the cited references.

<sup>c</sup> The ratio between the expected initial rates for the hydrolysis of 100  $\mu$ M nitrocefin [ $(v_0)_N$ ] and 100  $\mu$ M CENTA [ $(v_0)_C$ ] by the same amount of enzyme.

Since the appearance of the chromophore is related to the expulsion of the C-3' leaving group which is not concomitant with the opening of the  $\beta$ -lactam ring (4), it was important to verify that this expulsion was sufficiently rapid so that no artifact would be introduced in the measurement of initial rates or complete time courses. To do so, CENTA (30 to 60  $\mu$ M) was hydrolyzed at 30°C with increasing concentrations of the *E. cloacae* 908R class C  $\beta$ -lactamase (0.03 to 20  $\mu$ M). The variation in the absorbance was monitored at 260 nm (hydrolysis of the endocyclic amide bond) or 405 nm (appearance of the expelled chromophore) for 1 to 80 s on a stopped-flow spectrophotometer (Biologic SFM-3; Grenoble, France). Figure 2 shows that a clear lag becomes detectable in the curve for 405 nm when the reaction is completed within about 1 s. From these data, it can be estimated that the first-order rate constant characterizing the expulsion step is  $\geq 5 \text{ s}^{-1}$  and that, as a consequence, the rate of this reaction is unlikely to influence the values of the rate constants derived from complete time courses recorded over at least 1 min or from initial rate measurements performed over at least 30 s. Accordingly, the increase in the  $A_{405}$  value was linear under conditions in which about 5% of a 200  $\mu$ M CENTA solution was hydrolyzed in 1 min, and within the limits of experimental errors, the initial rates were the same when derived from measurements obtained when the absorbance was monitored at 405 or 260 nm.

**FIG. 2.** Complete hydrolysis of 60  $\mu$ M CENTA by 20  $\mu$ M *E. cloacae* 908R  $\beta$ -lactamase. The light line shows the increase in the  $A_{405}$ , and the heavy line shows the decrease in the  $A_{260}$ . The ordinate on the right ( $A_{260}$ ) has been inverted to facilitate the comparison. The optical pathway of the cell was 1 cm.



The chemical properties of CENTA are also favorable. In contrast to nitrocefin, whose stock solution must be prepared in dimethyl sulfoxide or dimethylformamide, CENTA is highly soluble in aqueous buffers. At pH 7 and in 200 mM sodium phosphate buffer, it was soluble up to a final concentration of 60 mg/ml. The stability of a 100  $\mu$ M solution was analyzed at a pH range of 4 to 12 at 25°C. The following buffers were used: for pH 4 and 5, 50 mM sodium acetate-acetic acid; for pH 6, 10 mM sodium cacodylate-HCl; for pH 7, 50 mM sodium phosphate; for pH 8, 10 mM Tris-HCl and 10 mM HEPES-NaOH; for pH 9 and 10, 50 mM CAPSO-NaOH; for pH 11, 50 mM Na<sub>2</sub>HPO<sub>4</sub>-NaOH; for pH 12, 50 mM KCl-NaOH. The absorbance at 405 nm was determined after increasing periods of time. When the incubation was performed below pH 7, the reading was made after adjustment of the pH of an aliquot to pH 7. Up to pH 9, no significant spontaneous hydrolysis of CENTA could be detected after a 60-min incubation. At higher pH values, the compound was less stable and the hydrolysis rates constants were  $0.7 \times 10^{-5} \text{ s}^{-1}$  at pH 11 and  $1 \times 10^{-3} \text{ s}^{-1}$  at pH 12. At the latter pH, the rate of hydrolysis of nitrocefin was  $2.6 \times 10^{-3} \text{ s}^{-1}$ . Incubation of 100  $\mu$ M CENTA in rabbit serum diluted fourfold at pH 7 and 30°C did not result in significant hydrolysis after 1 h, whereas the half-life of nitrocefin was 13 min under the same conditions (15). Similarly, at 30°C, substantial aminolysis of CENTA was observed in the presence of 300 mM Tris-HCl (pH 8.0; half-life, 19 min [the half-life for nitrocefin is 38 min]), but no detectable degradation occurred in the same buffer at a 10 mM concentration, in which nitrocefin exhibited a half-life of 380 min.

These experiments demonstrate that CENTA is a readily obtained chromogenic substrate which can conveniently be used in kinetic studies of  $\beta$ -lactamases and for the detection of these enzymes in bacterial crude extracts or in chromatographic fractions during enzyme purification. It can also be easily used in high-throughput screening tests for the selection of new  $\beta$ -lactamase inactivators. Unfortunately, the absorption spectrum of the leaving group is such that the contrast is not sufficient for the direct detection of  $\beta$ -lactamase-producing colonies on agar plates or on paper strip tests or for the localization of  $\beta$ -lactamases after gel isoelectric focusing.

### Acknowledgements

This work was supported by the Belgian government (grant PAI P 4/03), a TMR contract with the European Union (grant CT98-0232), and a grant from the Fonds National de la Recherche Scientifique (Brussels, Belgium) for the purchase of the stopped-flow apparatus. C.B. was the beneficiary of Comoth (University of Liège) and FRIA (Brussels) predoctoral fellowships.

### References

1. Bouillenne, F., A. Matagne, B. Joris, and J.-M. Frère. 2000. Technique for a rapid and efficient purification of the SHV-1 and PSE-2 beta-lactamases. *J. Chromatogr. B Biomed. Sci. Appl.* 737:261-265.
2. Carfi, A., S. Pares, E. Duée, M. Galleni, C. Duez, J. M. Frère, and O. Dideberg. 1995. The 3-D structure of a zinc metallo-beta-lactamase from *Bacillus cereus* reveals a new type of protein fold. *EMBO J.* 14:4914-4921.
3. De Meester, F., B. Joris, G. Reckinger, C. Bellefroid-Bourguignon, J. M. Frère, and S. G. Waley. 1987. Automated analysis of enzyme inactivation phenomena. Application to beta-lactamases and DD-peptidases. *Biochem. Pharmacol.* 36:2393-2403.

4. Faraci, W. S, and R. F. Pratt. 1985. Mechanism of inhibition of the PCI beta-lactamase of *Staphylococcus aureus* by cephalosporins: importance of the 3'-leaving group. *Biochemistry* 24:903-910.
5. Franceschini, N., B. Caravelli, J. D. Docquier, M. Galleni, J. M. Frère, G. Amicosante, and G. M. Rossolini. 2000. Purification and biochemical characterization of the VIM-1 metallo-β-lactamase. *Antimicrob. Agents chemother.* 44:3003-3007.
6. Frère, J. M., M. Leyh-Bouille, J. M. Ghuyse, M. Nieto, and H. R. Perkins. 1976. Exocellular DD-carboxypeptidases-transpeptidases from *Streptomyces*. *Methods Enzymol.* 45:610-636.
7. Galleni, M., and J. M. Frère. 1988. A survey of the kinetic parameters of class C beta-lactamases. *Biochem. J.* 255:119-122.
8. Granier, B., M. Jamin, M. Adam, M. Galleni, B. Lakaye, W. Zorzi, J. Granchamps, J. M. Wilkin, C. Fraipont, B. Joris, C. Duez, M. Nguyen-Distèche, J. Coyette, M. Leyh-Bouille, J. Dusart, L. Christiaens, J.-M. Frère, and J.-M. Ghuyse. 1994. Serine-type D-Ala-D-Ala peptidases and penicillin-binding proteins. *Methods Enzymol.* 244:249-266.
9. Hernandez Valladares, M., A. Felici, G. Weber, H. W. Adolph, M. Zeppeza-uer, G. M. Rossolini, G. Amicosante, J.-M. Frère, and M. Galleni. 1997. Zn(II) dependence of the *Aeromonas hydrophila* AE036 metallo-beta-lactamase activity and stability. *Biochemistry* 36:11534-11541.
10. Ishii, Y., A. Ohno, H. Taguchi, S. Imajo, M. Ishiguro, and H. Matsuzawa. 1995. Cloning and sequence of the gene encoding a cefotaxime-hydrolyzing class A beta-lactamase isolated from *Escherichia coli*. *Antimicrob. Agents Chemother.* 39:2269-2275.
11. Jones, R. N., H. W. Wilson, W. J. Novick, Jr., A. L. Barry, and C. Thornsberry. 1982. In vitro evaluation of CENTA, a new beta-lactamase-susceptible chromogenic cephalosporin reagent. *J. Clin. Microbiol.* 15:954-958.
12. Laraki, N., N. Franceschini, G. M. Rossolini, P. Santucci, C. Meunier, E. de Pauw, G. Amicosante, J.-M. Frère, and M. Galleni. 1999. Biochemical characterization of the *Pseudomonas aeruginosa* 101/1477 metallo-beta-lactamase IMP-1 produced by *Escherichia coli*. *Antimicrob. Agents Chemother.* 43: 902-906.
13. Ledent, P., X. Raquet, B. Joris, J. Van Beeumen, and J. M. Frère. 1993. A comparative study of class-D beta-lactamases. *Biochem. J.* 292(Pt 2):555-562.
14. Leyh-Bouille, M., M. Nguyen-Distèche, S. Pirlot, A. Veithen, C. Bourguignon, and J. M. Ghuyse. 1986. *Streptomyces* K15 DD-peptidase-catalysed reactions with suicide beta-lactam carbonyl donors. *Biochem. J.* 235:177-182.
15. O'Callaghan, C. H., A. Morris, S. M. Kirby, and A. H. Shingler. 1972. Novel method for detection of beta-lactamases by using a chromogenic cephalosporin substrate. *Antimicrob. Agents Chemother.* 1:283-288.
16. Paul-Soto, R., M. Hernandez-Valladares, M. Galleni, R. Bauer, M. Zeppe-zauer, J.-M. Frère, and H.-W. Adolph. 1998. Mono- and binuclear Zn-beta-lactamase from *Bacteroides fragilis*: catalytic and structural roles of the zinc ions. *FEBS Lett.* 438:137-140.
17. Raquet, X., J. Lamotte-Brasseur, E. Fonzé, S. Goussard, P. Courvalin, and J. M. Frère. 1994. TEM beta-lactamase mutants hydrolysing third-generation cephalosporins. A kinetic and molecular modelling analysis. *J. Mol. Biol.* 244:625-639.
18. Rossolini, G. M., N. Franceschini, M. L. Riccio, P. S. Mercuri, M. Perilli, M. Galleni, J. M. Frère, and G. Amicosante. 1998. Characterization and sequence of the *Chryseobacterium (Flavobacterium) meningosepticum* carbapenemase: a new molecular class B beta-lactamase showing a broad substrate profile. *Biochem. J.* 332:145-152.
19. Swarén, P., L. Maveyraud, X. Raquet, S. Cabantous, C. Duez, J.-D. Pédelacq, S. Mariotte-Boyer, L. Mourey, R. Labia, M.-H. Nicolas-Chanoine, P. Nordmann, J.-M. Frère, and J.-P. Samama. 1998. X-ray analysis of the NMC-A beta-lactamase at 1.64 Å resolution, a class A carbapenemase with broad substrate specificity. *J. Biol. Chem.* 273:26714-26721.

20. Van der Linden, M. P. G., L. de Haan, O. Dideberg, and W. Keck. 1994. Site-directed mutagenesis of proposed active-site residues of penicillin-binding protein 5 from *Escherichia coli*. *Biochem. J.* 303:357-362.
21. Virden, R., A. F. Bristow, and R. H. Pain. 1975. The active site of penicillinase from *Staphylococcus aureus* PC1. Isolation of a specific covalent complex with the substrate quinacillin. *Biochem. J.* 149:397-401.
22. Voladri, R. K., D. L. Lakey, S. H. Hennigan, B. F. Menzies, K. M. Edwards, and D. S. Kernodle. 1998. Recombinant expression and characterization of the major beta-lactamase of *Mycobacterium tuberculosis*. *Antimicrob. Agents Chemother.* 42:1375-1381.
23. Yang, Y., B. A. Rasmussen, and K. Bush. 1992. Biochemical characterization of an imipenem hydrolysing metallo  $\beta$ -lactamase from *Bacteroides fragilis* TAL3636. *Antimicrob. Agents Chemother.* 36:1155-1157.