





Prediction of cow pregnancy status using conventional and novel mid-infrared predicted milk traits

H. Hammami^{1,2}, C. Bastin¹, A. Gillon¹, V.M.-R. Arnould^{1,3}, J. Stoll³, H. Soyeurt^{1,2}, and N. Gengler^{1,2}

¹ Animal Science Unit, Gembloux Agro-Bio Tech (GxABT), University of Liège, Gembloux, Belgium
² National Fund for Scientific Research (FNRS), Brussels, Belgium
³ Convis Herdbuch, Ettelbruck, Luxembourg




Context: Management Indicator Traits (MIT)

Fertility

- Economics:
 - Main mover affecting the economical return
- Genetics:
 - Traits with low heritabilities

➔ Innovative tools based on **robust MIT's** are of interest for:

- Profitability and sustainability
 - ✓ Reduction of production costs
 - ✓ Increase of incomes

Context: Pregnancy & milk composition

- Diagnosis methods: costly, labor and investments
- Negative energy balance (NEB) ➔ delayed first ovulation and pregnancy rates
- Association "energy balance - milk composition":
 - Variation in fat and protein / fat:protein
 - *de novo* synthesis of fatty acids (C6:0 to C14:0)
 - Body fat mobilisation (C16:0 and C18:0)
- No study looked into the associations between milk composition and probability of conception

Objective

Ability of cow milk characteristics to predict the cow pregnancy status once inseminated

- Using only conventional milk component (fat, protein, lactose, and SCC)
- Extended to fatty acids

Data

- **Data sets** "repro check program" CONVIS, Luxembourg
 - AI records (9,996) and diagnosis results (2,826)
 - Test-day records (40,548)
 - Spectral data (35,555)
 - ➔ 6,147 lactations from 4,674 cows in 169 herds
- **Pregnant cow is defined as**
 - Positively checked
 - If no check (based on new registered calving)
 - Otherwise the cow was discarded from analysis

Methods

- **Predictors** (Milk components)
 - Conventional
 - Modified best prediction method (Gillon *et al.*, 2010)
 - ✓ Yields at specific DIM
 - ✓ Cumulated yields at specific DIM
 - ✓ Peaks, minimum
 - ✓ Ratios
 - Fatty acids
 - MIR equations (Soyeurt *et al.*, 2011)
 - ✓ Yields at the nearest TD to last AI
 - ✓ Cumulated yields at specific DIM
 - ✓ Ratios

Methods

- Separate **logistic regression** models
 - 3 periods
 - 35 to 44 days from last AI (DAI)
 - 45 to 60 DAI
 - 60 to 90 DAI
 - Lactation number (1, 2, 3 and plus)
 - Holsteins
- **Calibration** dataset (n=1,346 cows)
- **Validation** dataset (n=733 cows)

Predictive power "Holsteins 1st lactation"

Associations between predicted and observed probabilities of pregnancy at 3 periods



Only conventional

DAI	35-44	45-60	60-90
R ² (calibration)	0.96	0.98	0.96
R ² (validation)	0.76	0.83	0.76

Conventional + FA



DAI	35-44	45-60	60-90
R ² (calibration)	0.97	0.99	0.96
R ² (validation)	0.79	0.85	0.83

Predictive power "Holsteins - 1st lact. 45-60 DAI"




The dots represent the average predicted probabilities of 10 groups of cows that are plotted against their respective average observed probabilities

Sensitivity and specificity



Expert system



Conclusions

- Logistic regression model was able to predict the pregnancy status using combination of predictors based on milk routine analysis (**even if the cow number was limited**).
- FA predictors added to conventional milk component measurements improved slightly the prediction ability of studied models.
- Farmers could be able to identify pregnant cows and limit diagnosis to only problematic cows.

Perspectives

- ❑ Need more data and **cows with spectral data** to validate the final models
- ❑ Apply multi-level logistic regression models (multi-lactations, breeds, production systems,...)
- ❑ Evaluate the potentiality of models fits and prediction power when additionally using extra-data (health, feeding, BCS,...)

Acknowledgments



Thanks a lot for your attention



Corresponding author's e-mail: hedi.hammami@ulg.ac.be