

## In Vitro Activities of *Strychnos* Alkaloids and Extracts against *Plasmodium falciparum*

MICHEL FREDERICH<sup>1,2\*</sup>, MARIE-PIERRE HAYETTE<sup>2</sup>, MONIQUE TITS<sup>1</sup>, PATRICK DE MOL<sup>2</sup>, AND LUC ANGENOT<sup>1</sup>

<sup>1</sup>Laboratoire de Pharmacognosie, Institut de Pharmacie, CHU du Sart Tilman, Tour 4, B36 and

<sup>2</sup>Laboratoire de Microbiologie Médicale, Institut de Pathologie, CHU du Sart Tilman, Tour 3, B23 Liège, Belgium

Corresponding author. Laboratoire de Pharmacognosie, Institut de Pharmacie, Avenue de l'Hôpital 1, CHU - Sart Tilman, Tour 4, B36, 4000 Liège, Belgium.

### ABSTRACT


The in vitro antimalarial activities of 46 alkaloids and extracts from *Strychnos* species were evaluated. Two types of quasidimeric alkaloids exhibit high and selective activities against *Plasmodium*. Strychnopentamine and isostrychnopentamine were active against chloroquine-sensitive and-resistant strains (50% inhibitory concentration [IC<sub>50</sub>] ≈ 0.15 μM), while dihydrousambarensine exhibited a 30-fold higher activity against the chloroquine-resistant strain (IC<sub>50</sub> = 0.03 μM) than it did against the chloroquine-sensitive strain.

Malaria is the major parasitic infection in many tropical and subtropical regions. An increase in the resistance of *Plasmodium falciparum* to conventional treatments is a worldwide problem, and few alternative drugs are under development, necessitating urgent efforts to identify new classes of antimalarial drugs. The clinical utility of quinine and quinidine, isolated from *Cinchona* tree bark, and the Chinese discovery of artemisinin from the herb

*Artemisia annua* have stimulated much interest in plants as potential sources of new antimalarial drugs.

The in vitro antiplasmodial, antiameobic, and cytotoxic activities of alkaloids isolated from *Strychnos usambarensis* have been previously reported (5, 12, 17). In this article, we report the antiplasmodial activities against two additional *P. falciparum* strains of 19 total extracts, 21 new *Strychnos* alkaloids, and 6 alkaloids which were previously tested against the K1 strain of *P. falciparum* (17).

**Figure 1.** Chemical structures of *Strychnos* alkaloids.



Dihydrousambarensine, usambarensine, Nb-methylusambarensine, 10'-hydroxyusambarensine, malindine, usambarine, dihydrousambarine, strychnopentamine, isostrychnopent-amine, isostrychnofoline, strychnophylline, 10-hydroxyusambarine, 11-hydroxyusambarine, 10-hydroxydihydrousambarine, 11-hydroxydihydrousambarine, and tetradehydrolongicaudatine Y were isolated from *Strychnos usambarensis* 'Gilg' root bark (1, 7, 10), leaves (2), or stem bark (9). The root bark and leaves of *Strychnos icaja* "Baill." were the sources of 14-hydroxyepoxynovacine, epoxynovacine, icajine, sungucine, and bisnordihydrotoxiferine (11) and of vomicine and novacine (4), respectively. Holstiine and diaboline, strychnochromine, and guianensine were obtained from *Strychnos henningsii* "Gilg"

root bark, *Strychnos gossweileri* “Exell.” root bark, and *Strychnos guianensis* “Aubl.” stem bark, respectively (3, 6, 14). The chemical structures of most alkaloids are shown in Fig. 1. The total plant extracts were prepared by maceration of powdered material (5 g) with ethanol (EtOH) or ethyl acetate (EtOAc) (4 3 50 ml). The extracts were assembled and concentrated to dryness under vacuum. Voucher specimens of the different plants used in this study were deposited in the herbarium of the Pharmaceutical Institute at Liège, Belgium. *P. falciparum* strains were continuously maintained in culture by the method of Trager and Jensen (16) and as described previously (10). Tests for antimalarial activity were adapted from methods described earlier (8, 10, 13). Chloroquine diphosphate (Sigma) and quinine base (Aldrich) were used as antimalarial references. Parasite growth was estimated by [<sup>3</sup>H]hypoxanthine incorporation. The results are expressed as percentage of growth inhibition. The sigmoid dose-response curve was used to derive 50% inhibitory concentration values (IC<sub>50</sub>s) as the means of two experiments. The human cancer cell lines KB and HeLa were cultured, and the IC<sub>50</sub>s were determined as previously described (5).

The in vitro antimalarial activities of extracts and alkaloids against the two strains are shown in Table 1. The different total extracts displayed a wide range of antiplasmodial activities. Higher activities (IC<sub>50</sub> < 1 µg/ml) were obtained with root and leaf extracts from *S. usambarensis* and with root extracts from *S. icaja*. The EtOAc extract from *S. icaja* root was about five times more active than the EtOH extract. The stem bark extract of the liana form of *S. usambarensis* also had an IC<sub>50</sub> of <1 µg/ml, while the tree form extract had an IC<sub>50</sub> of about 25 µg/ml. This can be explained by the difference in alkaloid compositions between these two forms (15). Four *Strychnos* batches were found to exhibit moderate activities, with IC<sub>50</sub>s ranging between 10 and 30 µg/ml, namely, *Strychnos variabilis* root bark, *Strychnos angolensis* leaves, *S. angolensis* root, and *Strychnos memecyloides* leaf.

A large number of alkaloids from *S. usambarensis* were highly active against the chloroquine-sensitive FCA 20 strain. Dihydrousambarensine (IC<sub>50</sub> = 0.857 µM), 11-hydroxy-usambarine (0.487 µM), strychnopentamine (0.117 µM), and isostrychnopentamine (0.120 µM) were found to be the most active alkaloids, with IC<sub>50</sub>s of <1 µM.

**TABLE 1.** *In vitro* activities of crude extracts and alkaloids from some *Strychnos* species against two *P. falciparum* strains<sup>a</sup>

Compound N°	FCA 20 Ghana (chloroquine-sensitive strain)			W2 Indochina (chloroquine-resistant strain)		
	IC <sub>50</sub>	IC <sub>90</sub>	n <sup>b</sup>	IC <sub>50</sub>	IC <sub>90</sub>	n <sup>b</sup>
<b>Total extracts</b>						
<i>S. usambarensis</i> roots, EtOAc extract	(0.457 ± 0.005)	(2.345)	2	(0.039 ± 0.0029)	(0.597)	3
<i>S. usambarensis</i> leaves, EtOH extract	(0.287 ± 0.022)	(1.977)	3	(0.970 ± 0.141)	(4.853)	2
<i>S. usambarensis</i> bark liana variety, EtOAc extract	(0.766 ± 0.022)	(2.378)	2	(0.226 ± 0.039)	(2.897)	2
<i>S. usambarensis</i> bark tree variety, EtOH extract	(23.507)	(47.855)	1	(24.832)	(46.854)	1
<i>S. scheffleri</i> bark, EtOH extract	Inactive at (30)		1	ND <sup>c</sup>		
<i>S. matogrossensis</i> roots, EtOH extract	Inactive at (30)		1	ND		
<i>S. henningii</i> leaves, EtOH extract	(≅40)	(≅200)	1	ND		
<i>S. longicaudata</i> roots, EtOH extract	Inactive at (50)		1	ND		
<i>S. variabilis</i> bark, EtOH extract	(17.702)	(≅50)	1	ND		
<i>S. camptoneura</i> roots, EtOH extract	(≅50)		1	ND		
<i>S. innocua</i> roots, EtOH extract	Inactive at (40)		1	ND		
<i>S. angolensis</i> leaves, EtOH extract	(29.935)	(≅80)	1	ND		
<i>S. angolensis</i> roots, EtOH extract	(20.895)	(≅60)	1	ND		
<i>S. gossweileri</i> roots, EtOH extract	(43.505)	(≅400)	1	ND		
<i>S. icaja</i> leaves, EtOH extract	(40.421)	(94.159)	1	ND		
<i>S. icaja</i> roots, EtOH extract	(1.433 ± 0.068)	(4.834)	5	(1.002 ± 0.054)	(4.628)	2
<i>S. icaja</i> roots, EtOAc extract	(0.286 ± 0.061)	(1.381)	3	(0.303 ± 0.022)	(1.998)	2
<i>S. spinosa</i> bark, EtOH extract	Inactive at (80)		1	ND		
<i>S. memecyloides</i> leaves, EtOH extract	(13.489)	(48.75)	1	ND		
<b>Alkaloids</b>						
<i>S. usambarensis</i>						
Dihydrousambarensine	0.857 ± 0.061 (0.371)	2.486	5	0.032 ± 0.002 (0.013)	4.676	4
Usambarensine <sup>d</sup>	1.516 ± 0.031 (0.655)	4.412	6	0.594 ± 0.052 (0.257)	4.831	3
10'-Hydroxyusambarensine	1.071 ± 0.031 (0.480)	3.619	3	0.357 ± 0.035 (0.160)	6.851	3
Nb-Methylusambarensine <sup>d</sup>	5.045 (2.436)	9.835	1	3.919 ± 0.083 (1.893)	≅20	2
Strychnopentamine <sup>d</sup>	0.117 ± 0.033 (0.064)	0.443	4	0.145 ± 0.020 (0.079)	2.982	4
Isostrychnopentamine	0.120 ± 0.042 (0.066)	0.450	2	0.152 ± 0.009 (0.070)	0.628	2
Usambarine <sup>d</sup>	2.501 (1.125)	9.207	1	2.358 ± 0.015 (1.061)	7.294	2
Dihydrousambarine <sup>d</sup>	1.247 (0.673)	6.051	1	1.408 ± 0.013 (0.636)	5.515	2
11-Hydroxyusambarine	0.487 ± 0.015 (0.227)	1.770	5	1.538 ± 0.109 (0.716)	4.535	3
10-Hydroxyusambarine	1.134 ± 0.012 (0.528)	3.384	2	1.958 ± 0.115 (0.912)	5.803	2
Strychnophylline	1.009 ± 0.035 (0.570)	4.491	2	2.523 ± 0.134 (1.425)	9.310	2
Isostrychnofoline	5.657 ± 0.126 (3.2923)	25.959	2	2.598 ± 0.420 (1.519)	15.745	2
10-Hydroxydihydrousambarine	0.937 ± 0.044 (0.438)	5.522	3	4.699 ± 0.373 (2.199)	35.687	3
11-Hydroxydihydrousambarine	2.864 ± 0.046 (1.340)	7.023	2	5.449 ± 0.019 (2.550)	15.772	2
Malindine	162.63 (49.43)	≅800	1	Inactive at 60 (20)		1
Tetradehydrologicaudatine Y	1.236 (0.823)	7.970	1	0.958 ± 0.129 (0.543)	12.087	2
<i>Strychnos icaja</i>						
Vomicine	Inactive at 30 (12)		1	ND		
Novacine	Inactive at 90 (40)		1	ND		
14-Hydroxyepoxynovacine	Inactive at 50 (20)		1	ND		
Epoxynovacine	Inactive at 50 (20)		1	ND		
Icajine	Inactive at 80 (30)	198.16	2	95 ± 5 (34.7)	≅28	2
Sungucine	2.292 ± 0.049 (1.435)	11.182	2	1.659 ± 0.089 (1.051)	12.327	2
Bisnordihydrotoxiciferine	3.826 (2.112)	22.30	1	4.480 ± 0.092 (2.470)	13.123	2
Strychnine	Inactive at 20 (7)		1	23 (8.5)	≅50	1
<b>Other <i>Strychnos</i> species</b>						
Holstiine	≅80 (30)		1	34 (12.9)	≅100	1
Diaboline	Inactive at 33 (12)		1	ND		
Strychnochromine	40.046 (11.93)	≅200	1	16 (4.82)	≅70	1
Guianensine	5.920 (3.694)	≅20	1	7309 ± 486 (4.561)	24.765	2
<b>Reference compounds</b>						
Chloroquine	0.020 ± 0.002 (0.006)	0.119	9	0.284 ± 0.017 (0.091)	1.745	6
Quinine	0.269 ± 0.006 (0.087)	1.913	3	0.413 ± 0.011 (0.134)	1.718	2

<sup>a</sup> Data are expressed as mean micromolar concentrations ± standard deviations. Values in parentheses are given in micrograms per milliliter.

<sup>b</sup> n, number of independent experiments. All experiments were realized in duplicate.

<sup>c</sup> ND, not determined.

<sup>d</sup> These six alkaloids were previously tested against *P. falciparum* with essentially the same IC<sub>50</sub> as that for the W2 strain (17).

**TABLE 2.** Cytotoxic activity toward human cancer cell lines (KB and HeLa) and antiprotozoal selectivity index<sup>a</sup> of some *Strychnos* alkaloids

Compound	IC <sub>50</sub> ( $\mu$ M)	Selectivity index <sup>a</sup>	
		FCA	W2
Dihydrousambarensine (KB)	12	14	375
Usambarensine (KB)	9.7	6	16
10'-Hydroxyusambarensine (HeLa)	20	19	56
Strychnopentamine (KB)	11.3	96	78
Tetradehydrolongicaudatine Y (HeLa)	>50 <sup>b</sup>	>40	>52

<sup>a</sup> Selectivity index is defined as the ratio of cytotoxicity (IC<sub>50</sub>) to antiplasmodial activity (IC<sub>50</sub>).

<sup>b</sup> This compound has no effect at 50  $\mu$ M.

Eight other alkaloids (usambarensine, 10'-hydroxyusambarensine, usambarine, dihydrousambarine, 10- and 11-hydroxyusambarines, strychnophylline, and tetradehydrolongicaudatine Y) possess IC<sub>50</sub>s between 1 and 2  $\mu$ M. The activities against the W2 strain were of the same order as those observed with the FCA strain. However, two compounds (usambarensine and 10'-hydroxyusambarensine) were more than twice as active against the resistant clone than the susceptible clone, and the IC<sub>50</sub> for one compound (dihydrousambarensine) was 30-fold lower for the chloroquine-resistant strain (32 nM). Its 90% inhibitory concentration (IC<sub>90</sub>), however, remained higher, with a value of 4.6  $\mu$ M. The IC<sub>50</sub> of strychnopentamine for both strains was the same (0.15  $\mu$ M), and its isomer, isostrychnopentamine, exhibited the same activity but with a lower IC<sub>90</sub> for strain W2 (0.6  $\mu$ M). On the other hand, some alkaloids, like strychnophylline and the hydroxyusambarines, were notably less active against the W2 strain.

Among the known alkaloids of *S. icaja* roots, only sungucine and bisnordihydrotoxiferine were slightly active, with an IC<sub>50</sub> of 2 to 4  $\mu$ M ( $\approx$ 1 and 2  $\mu$ g/ml). However, these activities could not explain the high IC<sub>50</sub> (0.3  $\mu$ g/ml) of the EtOAc extract. Further investigations to find the compounds responsible for this activity are actually in progress.

The most active compounds were tested for cytotoxicity against the human cell lines KB and HeLa. All of these compounds exhibited a 6- to 400-fold higher activity against *Plasmodium* than against human cells, thus indicating some selectivity (Table 2). The most selective compounds were strychnopentamine and tetradehydrolongicaudatine Y, with, respectively, 70 to 100 and

>40 to 50 times higher activities against the two *Plasmodium* strains. Dihydrousambarensine showed a good selectivity for the W2 strain only (400 times higher activity).

All active compounds were tertiary dimers. There is no clear relationship which allows us to associate some structural features (stereochemistry or substitutions, etc.) with an increase in activity. Nevertheless, the results of the present study confirm the antimalarial activities of *Strychnos* bisindole alkaloids and more particularly of two alkaloid types. The usambarensine skeleton is linked with an important rise in the activity on chloroquine-resistant strains, more particularly for the 5',6'- dihydro derivative. Strychnopentamine and its isomer demonstrate high and selective activities against the two strains. It would therefore be interesting to test closely related compounds that possess these kinds of structures.

This research was supported by the Belgian National Fund for Scientific Research (grant 3451997 and a fellowship to M.F.).

We thank J. Boniver (Anatomie et Cytologie Pathologiques, University of Liège) for liquid scintillation measurements, M. C. DePauw (Cytologie et Histologie, University of Liège) for cytotoxicity testing,

M. Wéry (Tropical Medicine Institute, Antwerp, Belgium), and J. Le Bras (Hôpital Bichat-Claude Bernard, Laboratoire de Parasitologie, Paris, France) for providing the *P. falciparum* strains.

## REFERENCES

1. Angenot, L., and N. G. Bisset. 1971. *Isolement et structure de nouveaux alcaloïdes extraits du Strychnos usambarensis GILG du Rwanda. J. Pharm. Belg.* 26:585–588.
2. Angenot, L., C. Coune, and M. Tits. 1978. *Nouveaux alcaloïdes des feuilles du Strychnos usambarensis. J. Pharm. Belg.* 33:11–23.
3. Angenot, L., and M. Tits. 1981. *Isolement d'un nouvel alcaloïde et d'un triterpénoïde à partir de Strychnos henningsii du Zaïre. Planta Med.* 41:240–243.
4. Bisset, N. G., B. Das, and J. Parello. 1973. *Alkaloids from the leaves of Strychnos icaja Baill. Tetrahedron* 29:4137–4148.
5. Bonjean K., M. C. De Pauw, J. Quetin-Leclercq, L. Angenot, and R. Bassleer. 1996. *In vitro cytotoxic activity of two potential anticancer drugs isolated from Strychnos: strychnopentamine and usambarensine. Anticancer Res.* 16: 1129–1138.
6. Bosly, J. 1951. *Etude des alcaloïdes du Strychnos Holstii. J. Pharm. Belg.* 6:150.

7. Caprasse, M., D. Tavernier, M. Anteuinis, and L. Angenot. 1984. Isolement de Nb-méthylantirhine, malindine et isomalindine à partir du *Strychnos usambarensis*. *Planta Med.* 44:27–30.
8. Desjardins, R., C. Canfield, J. Haynes, and J. Chulay. 1979. Quantitative assessment of antimalarial activity *in vitro* by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.* 16:710–718.
9. Frédéricich, M., J. Quetin-Leclercq, R. GadiBiala, V. Brandt, J. Penelle, M. Tits, and L. Angenot. 1998. 3',4',5',6' Tetradehydrolongicaudatine Y, an anhydronium base from *Strychnos usambarensis*. *Phytochemistry* 48:1263–1266.
10. Frédéricich, M., M. Tits, M. P. Hayette, V. Brandt, J. Penelle, P. De Mol, G. Llabrés, and L. Angenot. 1999. 10'-Hydroxyusambarensine, a new bisindole alkaloid from the roots of *Strychnos usambarensis*. *J. Nat. Prod.* 62:619–621.
11. Kambu, K., C. Coune, and L. Angenot. 1979. Nouveaux alcaloïdes des racines du *Strychnos icaja*. *Planta Med.* 37:161–164.
12. Leclercq J., M. C. DePauw, R. Bassleer, and L. Angenot. 1986. Screening of cytotoxic activities of *Strychnos* alkaloids. *J. Ethnopharmacol.* 15:305–316.
13. Mirovsky, P., F. Gay, D. Bustos, D. Mazier, and M. Gentilini. 1990. Cloning of a fresh isolate of *Plasmodium falciparum* and drug sensitivity of the clones. *Trans. R. Soc. Trop. Med. Hyg.* 84:511–515.
14. Quetin-Leclercq, J., G. Llabrés, R. Warin, M. Belem-Pinheiro, H. MavarManga, and L. Angenot. 1995. Guianensine, a zwitterionic alkaloid from *Strychnos guianensis*. *Phytochemistry* 40:1557–1559.
15. Quetin-Leclercq, J., M. Tits, L. Angenot, and N. G. Bisset. 1991. Alkaloids of *Strychnos usambarensis* stem bark. *Planta Med.* 57:501–502.
16. Trager, W., and J. B. Jensen. 1976. Human malaria parasites in continuous cultures. *Science* 193:673–675.
17. Wright, C. W., D. Allen, Y. Cai, Z. Chen, J. D. Phillipson, G. Kirby, D. Warhurst, M. Tits, and L. Angenot. 1994. Selective antiprotozoal activity of some *Strychnos* alkaloids. *Phytother. Res.* 8:149–152.